



# PIZZA SALES

BY RENUKALIMBHARE





# RETRIEVE THE TOTAL NUMBER OF ORDERS PLACED

```
select count(order_id) as total_orders from orders;
```

total_orders
21350



# CALCULATE THE TOTAL REVENUE GENERATED FROM PIZZA SALES.

```
SELECT
    ROUND(SUM(order_details.quantity * pizzas.price),
          2) AS total_sales
FROM
    order_details
    JOIN
    pizzas ON pizzas.pizza_id = order_details.pizza_id
```

total_sales
817860.05



# IDENTIFY THE HIGHEST-PRICED PIZZA.

```
SELECT
    pizza_types.name, pizzas.price
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
ORDER BY pizzas.price DESC
LIMIT 1;
```

name	price
The Greek Pizza	35.95



# IDENTIFY THE MOST COMMON PIZZA SIZE ORDERED.

```
SELECT
    pizzas.size,
    COUNT(order_details.order_details_id) AS order_count
FROM
    pizzas
    JOIN
    order_details ON pizzas.pizza_id = order_details.pizza_id
GROUP BY pizzas.size
ORDER BY order_count DESC;
```

size	order_count
L	18526
M	15385
S	14137
XL	544
XXL	28



# LIST THE TOP 5 MOST ORDERED PIZZA TYPES ALONG WITH THEIR QUANTITIES.

```
SELECT
    pizza_types.name, SUM(order_details.quantity) AS quantity
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY quantity DESC
LIMIT 5;
```

name	quantity
The Classic Deluxe Pizza	2453
The Barbecue Chicken Pizza	2432
The Hawaiian Pizza	2422
The Pepperoni Pizza	2418
The Thai Chicken Pizza	2371



# JOIN THE NECESSARY TABLES TO FIND THE TOTAL QUANTITY OF EACH PIZZA CATEGORY ORDERED

```
SELECT
    pizza_types.category,
    SUM(order_details.quantity) AS quantity
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.category
ORDER BY quantity DESC;
```

category	quantity
Classic	14888
Supreme	11987
Veggie	11649
Chicken	11050



# DETERMINE THE DISTRIBUTION OF ORDERS BY HOUR OF THE DAY.

```
SELECT
    HOUR(order_time) AS hour, COUNT(order_id) AS order_count
FROM
    orders
GROUP BY HOUR(order_time);
```

hour	order_count
11	1231
12	2520
13	2455
14	1472
15	1468



# JOIN RELEVANT TABLES TO FIND THE CATEGORY-WISE DISTRIBUTION OF PIZZAS

```
SELECT  
    category, COUNT(name)  
FROM  
    pizza_types  
GROUP BY category;
```

category	count(name)
Chicken	6
Classic	8
Supreme	9
Veggie	9



# GROUP THE ORDERS BY DATE AND CALCULATE THE AVERAGE NUMBER OF PIZZAS ORDERED PER DAY.

```
SELECT  
    ROUND(AVG(quantity), 0)  
FROM  
    (  
        SELECT  
            (orders.order_date), SUM(order_details.quantity) AS quantity  
        FROM  
            orders  
        JOIN order_details ON orders.order_id = order_details.order_id  
        GROUP BY orders.order_date) AS order_quantity;
```

avg_pizza_ordered
138



# DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE.

```
SELECT
    pizza_types.name,
    SUM(order_details.quantity * pizzas.price) AS revenue
FROM
    pizza_types
    JOIN
    pizzas ON pizzas.pizza_type_id = pizza_types.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY revenue DESC
LIMIT 3;
```

name	revenue
The Thai Chicken Pizza	43434.25
The Barbecue Chicken Pizza	42768
The California Chicken Pizza	41409.5



# CALCULATE THE PERCENTAGE CONTRIBUTION OF EACH PIZZA TYPE TO TOTAL REVENUE

```
SELECT
    pizza_types.category,
    ROUND(SUM(order_details.quantity * pizzas.price) / (SELECT
        ROUND(SUM(order_details.quantity * pizzas.price),
            2) AS total_sales
        FROM
            order_details
            JOIN
                pizzas ON pizzas.pizza_id = order_details.pizza_id) * 100,
        2) AS revenue
FROM
    pizza_types
    JOIN
        pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
        order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.category
ORDER BY revenue DESC;
```

category	revenue
Classic	26.91
Supreme	25.46
Chicken	23.96
Veggie	23.68



# ANALYZE THE CUMULATIVE REVENUE GENERATED OVER TIME.

```
select order_date,  
sum(revenue) over (order by order_date) as cum_revenue  
from  
(select orders.order_date,  
sum(order_details.quantity*pizzas.price) as revenue  
from order_details join pizzas  
on order_details.pizza_id= pizzas.pizza_id  
join orders  
on orders.order_id = order_details.order_id  
group by orders.order_date) as sales ;
```

order_date	cum_revenue
2015-01-01	2713.850000000000004
2015-01-02	5445.75
2015-01-03	8108.15
2015-01-04	9863.6
2015-01-05	11929.55



# DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE FOR EACH PIZZA CATEGORY.

```
select name, revenue from
(select category, name, revenue,
rank() over(partition by category order by revenue desc) as rn
from
(select pizza_types.category, pizza_types.name,
sum((order_details.quantity) * pizzas.price) as revenue
from pizza_types join pizzas
on pizza_types.pizza_type_id = pizzas.pizza_type_id
join order_details
on order_details.pizza_id = pizzas.pizza_id
group by pizza_types.category, pizza_types.name) as a) as b
where rn <= 3;
```

name	revenue
The Thai Chicken Pizza	43434.25
The Barbecue Chicken Pizza	42768
The California Chicken Pizza	41409.5
The Classic Deluxe Pizza	38180.5
The Hawaiian Pizza	32273.25





THANK YOU!

