Table S3. Exploratory X1/X2: group summaries of the per-cell Δ for every analysis, arm, comparator and group.

Arm	Comparator	Group	Cells	Mean Δ (95% interval)	Cells with Δ < 0
X1: eraser (LEACE) refitted on the 158-case analysis region					
FM-P−C	step-6 FM-P−C	P1b-FM	48	−0.033 (−0.042 to −0.023)	34
FM-P−C	step-6 FM-P−C	P2-FM	12	−0.002 (−0.007 to 0.005)	6
FM-P−C	step-6 FM-P−C	slice	12	−0.010 (−0.022 to 0.002)	9
FM-P−C	step-6 FM-P	P1b-FM	48	0.561 (0.549 to 0.573)	0
FM-P−C	step-6 FM-P	P2-FM	12	−0.007 (−0.013 to −0.001)	7
FM-P−C	step-6 FM-P	slice	12	−0.003 (−0.014 to 0.009)	6
FM-P−C	step-6 NV	P1b-FM	48	−1.250 (−1.277 to −1.224)	48
FM-P−C	step-6 NV	P2-FM	12	−0.024 (−0.032 to −0.013)	7
FM-P−C	step-6 NV	slice	12	−0.378 (−0.392 to −0.363)	11
FM-S−C	step-6 FM-S−C	P1b-FM	48	0.023 (0.014 to 0.033)	18
FM-S−C	step-6 FM-S−C	P2-FM	12	0.000 (−0.005 to 0.006)	7
FM-S−C	step-6 FM-S−C	slice	12	0.000 (−0.012 to 0.012)	7
FM-S−C	step-6 FM-S	P1b-FM	48	0.343 (0.333 to 0.353)	1
FM-S−C	step-6 FM-S	P2-FM	12	−0.004 (−0.009 to 0.002)	7
FM-S−C	step-6 FM-S	slice	12	0.006 (−0.006 to 0.019)	4
FM-S−C	step-6 NV	P1b-FM	48	−1.247 (−1.270 to −1.222)	46
FM-S−C	step-6 NV	P2-FM	12	−0.019 (−0.028 to −0.009)	7
FM-S−C	step-6 NV	slice	12	−0.384 (−0.400 to −0.369)	11
REF−C	step-6 REF−C	P1b-FM	48	0.004 (−0.006 to 0.012)	18
REF−C	step-6 REF−C	P2-FM	12	0.002 (−0.003 to 0.008)	5
REF−C	step-6 REF−C	slice	12	0.000 (−0.012 to 0.011)	5
REF−C	step-6 REF	P1b-FM	48	0.496 (0.484 to 0.508)	1
REF−C	step-6 REF	P2-FM	12	−0.006 (−0.013 to −0.001)	6
REF−C	step-6 REF	slice	12	0.004 (−0.008 to 0.016)	5
REF−C	step-6 NV	P1b-FM	48	−1.280 (−1.307 to −1.256)	47
REF−C	step-6 NV	P2-FM	12	−0.022 (−0.030 to −0.012)	7
REF−C	step-6 NV	slice	12	−0.406 (−0.421 to −0.392)	11
X2: linear-only nuisance learners					
FM-P−C	step-6 FM-P−C	P1b-FM	48	0.011 (0.002 to 0.020)	22
FM-P−C	step-6 FM-P−C	P2-FM	12	0.001 (−0.004 to 0.008)	7
FM-P−C	step-6 FM-P−C	slice	12	−0.003 (−0.014 to 0.008)	7

Arm	Comparator	Group	Cells	Mean Δ (95% interval)	Cells with Δ < 0
FM-P−C	step-6 FM-P	P1b-FM	48	0.605 (0.593 to 0.617)	0
FM-P−C	step-6 FM-P	P2-FM	12	−0.004 (−0.010 to 0.002)	7
FM-P−C	step-6 FM-P	slice	12	0.004 (−0.008 to 0.016)	3
FM-P−C	step-6 NV	P1b-FM	48	−1.206 (−1.233 to −1.179)	47
FM-P−C	step-6 NV	P2-FM	12	−0.021 (−0.030 to −0.012)	7
FM-P−C	step-6 NV	slice	12	−0.371 (−0.387 to −0.355)	11
FM-S−C	step-6 FM-S−C	P1b-FM	48	−0.003 (−0.012 to 0.006)	27
FM-S−C	step-6 FM-S−C	P2-FM	12	−0.0005[a] (−0.006 to 0.005)	6
FM-S−C	step-6 FM-S−C	slice	12	−0.001 (−0.013 to 0.011)	6
FM-S−C	step-6 FM-S	P1b-FM	48	0.317 (0.306 to 0.328)	0
FM-S−C	step-6 FM-S	P2-FM	12	−0.004 (−0.010 to 0.002)	7
FM-S−C	step-6 FM-S	slice	12	0.004 (−0.008 to 0.017)	3
FM-S−C	step-6 NV	P1b-FM	48	−1.273 (−1.299 to −1.248)	47
FM-S−C	step-6 NV	P2-FM	12	−0.020 (−0.029 to −0.010)	7
FM-S−C	step-6 NV	slice	12	−0.386 (−0.401 to −0.370)	11
REF−C	step-6 REF−C	P1b-FM	48	−0.0003[a] (−0.009 to 0.008)	21
REF−C	step-6 REF−C	P2-FM	12	−0.001 (−0.006 to 0.005)	7
REF−C	step-6 REF−C	slice	12	0.002 (−0.010 to 0.013)	7
REF−C	step-6 REF	P1b-FM	48	0.492 (0.480 to 0.504)	0
REF−C	step-6 REF	P2-FM	12	−0.009 (−0.015 to −0.003)	7
REF−C	step-6 REF	slice	12	0.005 (−0.006 to 0.018)	5
REF−C	step-6 NV	P1b-FM	48	−1.285 (−1.310 to −1.258)	47
REF−C	step-6 NV	P2-FM	12	−0.024 (−0.033 to −0.015)	7
REF−C	step-6 NV	slice	12	−0.405 (−0.420 to −0.389)	11

Exploratory (outside the frozen protocol; plan registered on OSF, osf.io/5c627); no verdict. Δ = (|bias| X arm − |bias| step-6 arm) / mean truth per cell, always X arm − step-6 arm (negative: the X arm has the smaller |bias|); mean Δ = unweighted mean over the group's cells; 95% interval: paired bootstrap percentile interval (2,000 resamples of the 200 synthetic replicates within each cell; cells fixed); simulation uncertainty, not real-data sampling uncertainty.

Groups: P1b-FM, the 48 W = 0 cells with P = 5 or 30; P2-FM, the 12 cells with K = 1, P = 0, α > 0; slice, the 12 main-text slice cells (W = 30, α = 0.5, γ = 1, φ_A = 0); cell 15 belongs to P2-FM and the slice. −C, linearly site-removed; NV, no video.

Ceiling (fact base §8): exploratory; no verdict; cannot restore P1b-FM/P2-FM; nonlinear site information remains after X1; X2 changes do not show that erasure caused a difference.

[a]Mean Δ shown with 4 decimals so that its sign is visible (3 decimals would display −0.000); all other values have 3 decimals.