Table S4. Exploratory X1: E3-site, E2 and E3-state retention on the 158-case analysis region, next to step 5.

(A) E3-site: case-level site balanced accuracy after removal

Representation	Probe	Step 5: BA (95% interval)	Step 5: one-sided 95% upper bound	X1: BA (95% interval)	X1: one-sided 95% upper bound
FM-P	linear	0.416 (0.352 to 0.481)	0.472	0.250 (0.250 to 0.250)	0.250
FM-P	MLP	0.702 (0.640 to 0.763)	0.755	0.577 (0.506 to 0.647)	0.637
FM-S	linear	0.393 (0.346 to 0.444)	0.435	0.250 (0.250 to 0.250)	0.250
FM-S	MLP	0.428 (0.368 to 0.489)	0.476	0.309 (0.255 to 0.366)	0.357
REF	linear	0.448 (0.387 to 0.518)	0.505	0.250 (0.250 to 0.250)	0.250
REF	MLP	0.620 (0.543 to 0.687)	0.680	0.458 (0.392 to 0.530)	0.517

(B) E2 primary: ΔAUC for S_t of base + R_t over base

Representation	Condition	ΔAUC (95% CI)	E2 criterion
FM-P	step 5, before removal	0.050 (0.019 to 0.080)	met
FM-P	step 5, after protocol removal	0.044 (0.012 to 0.074)	not met
FM-P	X1, after X1 removal	0.051 (0.019 to 0.083)	met
FM-S	step 5, before removal	0.002 (−0.017 to 0.024)	not met
FM-S	step 5, after protocol removal	0.003 (−0.018 to 0.025)	not met
FM-S	X1, after X1 removal	−0.0005[a] (−0.022 to 0.021)	not met
REF	step 5, before removal	0.024 (−0.0005[a] to 0.049)	not met
REF	step 5, after protocol removal	0.024 (−0.0004[a] to 0.051)	not met
REF	X1, after X1 removal	0.024 (−0.002 to 0.052)	not met

(C) E3-state retention (post / pre ΔAUC ≥ 0.50)

Representation	Step 5	X1
FM-P	retained fraction 0.868	retained fraction 1.019
FM-S	not applicable (E2 primary failed)	not applicable (E2 primary failed)
REF	not applicable (E2 primary failed)	not applicable (E2 primary failed)

Exploratory (X1; plan registered on OSF, osf.io/5c627); no verdict. X1 refits the LEACE eraser on the 158-case analysis region (training folds) and evaluates on held-out folds of the same population; step-5 values are the protocol LEACE fitted on the 278-case set. The linear E3-site value of X1 is a check of the procedure and does not gate X1. E3-site boundary 0.35 (one-sided upper bound); E2 criterion ΔAUC ≥ 0.05 and CI excluding 0. Intervals: case-level (cluster) bootstrap stratified by site (2,000 resamples).
Ceiling (fact base §8): exploratory; no verdict; cannot restore P1b-FM/P2-FM; nonlinear site information remains after X1; X2 changes do not show that erasure caused a difference. BA, balanced accuracy; MLP, two-layer multilayer perceptron probe.

[a]Shown with 4 decimals so that its sign is visible (3 decimals would display −0.000); all other values have 3 decimals.