


Supplementary Figure X (X1 vs step-6 k−C (same arm, original features)). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.




Supplementary Figure X (X1 vs step-6 FM-k (site retained)). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.




Supplementary Figure X (X1 vs step-6 NV). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.



Supplementary Figure X (X2 vs step-6 k−C (same arm, original features)). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.



Supplementary Figure X (X2 vs step-6 FM-k (site retained)). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.



Supplementary Figure X (X2 vs step-6 NV). EXPLORATORY (outside the frozen protocol). Per-cell Δ = (|bias_X| − |bias_step-6|) / mean truth with 95 % paired bootstrap intervals (B = 2,000); negative = X arm smaller |bias|. All 71 cells.

