

Last time : Relations (Sections 9.1 and 9.3)

A relation on A is a subset of $A \times A$

Reflexive: $\forall a \in A: aRa$


Symmetric: $\forall a, b \in A: (aRb \Rightarrow bRa)$

Antisymmetric: $\forall a, b \in A: (aRb \wedge a \neq b \Rightarrow b \not R a)$

Transitive: $\forall a, b, c \in A: (aRb \wedge bRc \Rightarrow aRc)$

Ex: $R = \{(1,3), (2,1), (2,2), (2,3), (3,1), (3,3)\}$
relation on $\{1,2,3,4\}$

$$\begin{bmatrix} 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$




Neither reflexive, symmetric, antisymmetric,
nor transitive.


Relations can be composed like functions

$$\text{Ex: } R = \{ (1,3), (2,1), (2,2), (2,3), (3,1), (3,3) \}$$

$$S = \{ (2,1), (3,1), (4,2), (4,4) \}$$




$$R \circ S = \{ (2,3), (3,3), (4,1), (4,2), (4,3) \}$$



$$S^2 = S \circ S = \{ (4,1), (4,2), (4,4) \}$$

$$= S^3 = S^4 = \dots$$



Closures of relations (Section 9.4)

Def. 9.4.1:

For any relation R and any property P , the closure of R w.r.t. P (if it exists) is the set C satisfying

(1) C has property P

(2) $R \subseteq C$

(3) \forall relation S satisfying (1) and (2): $C \subseteq S$

i.e., C is a smallest possible set containing R and having property P .


Ex: Relation on $\{1, 2, 3, 4\}$:

$$R = \{(1, 3), (2, 1), \underline{(2, 2)}, (2, 3), (3, 1), \underline{(3, 3)}\}$$

The reflexive closure of R is

$$r(R) = \{(\underline{1, 1}), (1, 3), (2, 1), (2, 2), (2, 3), (3, 1), (3, 3), (\underline{4, 4})\}$$

$$\begin{bmatrix} \color{red}{1} & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & \color{red}{1} \end{bmatrix}$$



R : relation on A

The reflexive closure of R is

$$r(R) = R \cup \{(a, a) \mid a \in A\}$$

Ex: Relation on \mathbb{N} :

$$R_< = \{(a, b) \mid a < b\}$$


$$\begin{aligned} r(R_<) &= R_< \cup \{(a, a) \mid a \in \mathbb{N}\} \\ &= \{(a, b) \in \mathbb{N} \times \mathbb{N} \mid a \leq b\} \end{aligned}$$

Ex: $R = \{(1,3), \underline{(2,1)}, \underline{(2,3)}, (3,1), (3,3)\}$

The symmetric closure of R is

$$S(R) = \{(1,2), (1,3), \underline{(2,1)}, \underline{(2,3)}, (3,1), \underline{(3,2)}, (3,3)\}$$

$$\begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$



The symmetric closure of a relation R is
 $S(R) = R \cup \{(b,a) \mid (a,b) \in R\}$

Ex:

$$S(R_{<}) = R_{<} \cup R_{>} = R_{\neq}$$

$$S(\text{"same parity"}) = \text{"same parity"}$$

Antisymmetric closure?

Not defined

A relation which is not antisymmetric cannot be made antisymmetric by adding pairs to it.

Ex: $R = \{(1,2), (2,3), (3,4)\}$




To make R transitive, we need to add R^2 :

$$\begin{aligned} R \cup R^2 &= \{(1,2), (2,3), (3,4)\} \cup \{(1,3), (2,4)\} \\ &= \{(1,2), (1,3), (2,3), (2,4), (3,4)\} \end{aligned}$$

But then we also need to add R^3 :

$$\begin{aligned} R \cup R^2 \cup R^3 &= \{(1,2), (1,3), (2,3), (2,4), (3,4)\} \cup \{(1,4)\} \\ &= \{(1,2), (1,3), (1,4), (2,3), (2,4), (3,4)\} \end{aligned}$$

And now, we are done, since $R^4 = \{\}$




Theorem 9.4.2

The transitive closure of any relation R is
 $t(R) = R^*$


Proof:

- $R \subseteq R^*$
- R^* is transitive (by Theorem 9.4.1):
For any paths $a \rightsquigarrow b$ and $b \rightsquigarrow c$,
 G_{R^*} also has an edge $a \rightarrow c$
- For any transitive relation S s.t. $R \subseteq S$,
 $R^* \subseteq S$

$$\text{Ex: } R = \{(1,3), (2,1), (2,2), (3,1), (3,3)\}$$




$$R^2 = \{(1,1), (1,3), (2,1), (2,2), (2,3), (3,3)\}$$



Note : For each path of length > 2 ,
there is a path of length ≤ 2 .

Thus,

$$\begin{aligned} t(R) &= R \cup R^2 \\ &= \{(1,1), (1,3), (2,1), (2,2), (2,3), (3,1), (3,3)\} \end{aligned}$$



Equivalence Relations

Section 9.5

Def 9.5.1

A relation which is

- reflexive
- symmetric
- transitive

is called an **equivalence relation**

Ex :

✓: $=$, same parity

✗: \neq , \leq , $|$

Def. 9.5.2

If R is an **equivalence relation** and $(a,b) \in R$,
then a and b are **equivalent**

Def. 9.5.3

If R is an equivalence relation on A and $a \in A$, then the equivalence class of a is

$$[a]_R = \{ b \mid (a, b) \in R \}$$

$$= \{ b \mid (b, a) \in R \}, \text{ since } R \text{ is symmetric}$$

Ex: Relation on \mathbb{N} :

$$\{ (a, b) \mid a \text{ and } b \text{ have the same parity} \}$$

$$[0]_R = \{ 0, 2, 4, 6, \dots \} = [2]_R = [4]_R = \dots$$


$$[1]_R = \{ 1, 3, 5, 7, \dots \} = [3]_R = [5]_R = \dots$$

Note:

$$[0]_R \cup [1]_R = \mathbb{N} \text{ and}$$

$$[0]_R \cap [1]_R = \emptyset,$$

i.e., the eq classes form a partition of \mathbb{N}



Ex: Relation on \mathbb{N} :


$$\{ (a, b) \mid a = b \}$$

$$[0]_s = \{ 0 \}$$

$$[1]_s = \{ 1 \}$$

$$[2]_s = \{ 2 \}$$

\vdots



Again, the eq. classes form a partition of \mathbb{N} .

This is not a coincidence, according to Thm 9.5.2

In order to prove Theorem 9.5.2, we will first prove Theorem 9.5.1:

Theorem 9.5.1:

If R is an equivalence relation, then

$$\begin{array}{ll} aRb & (i) \\ \Downarrow & \\ [a]_R = [b]_R & (ii) \\ \Downarrow & \\ [a]_R \cap [b]_R \neq \emptyset & (iii) \end{array}$$

The fact that $(iii) \Rightarrow (ii)$ is used in the proof of Theorem 9.5.2).

Proof:

We will prove $(i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i)$, giving a direct proof of each implication.

For simplicity we will write $[a]$ and $[b]$ instead of $[a]_R$ and $[b]_R$.

$$\underline{aRb \Rightarrow [a] = [b]} :$$

$$(i) \Rightarrow (ii)$$

Assume aRb .

Then, bRa , since R is symmetric.

$$\underline{[a] \subseteq [b]} :$$




$$\Downarrow c \in [a]$$

$$\Downarrow aRc, \text{ by def.}$$

$$\Downarrow bRc, \text{ since } bRa \text{ and } R \text{ is transitive}$$

$$\Downarrow c \in [b], \text{ by def.}$$

$$\underline{[b] \subseteq [a]} :$$




$$\Downarrow d \in [b]$$

$$\Downarrow bRd, \text{ by def.}$$

$$\Downarrow aRd, \text{ since } aRb \text{ and } R \text{ is transitive}$$

$$\Downarrow d \in [a], \text{ by def.}$$



Since $[a] \subseteq [b]$ and $[b] \subseteq [a]$, we conclude $[a] = [b]$.



$$\underline{[a] = [b] \Rightarrow [a] \cap [b] \neq \emptyset :} \quad (ii) \Rightarrow (iii)$$

$$\begin{aligned} & [a] = [b] \\ \Downarrow & [a] \cap [b] = [a] \\ \Downarrow & [a] \cap [b] \neq \emptyset, \text{ since } a \in [a] \text{ by reflexivity} \end{aligned}$$



$$\underline{[a] \cap [b] \neq \emptyset \Rightarrow aRb :} \quad (iii) \Rightarrow (ii)$$

$$\begin{aligned} & [a] \cap [b] \neq \emptyset \\ \Updownarrow & \exists c : c \in [a] \wedge c \in [b] \\ \Updownarrow & \exists c : aRc \wedge bRc, \text{ by def.} \\ \Updownarrow & \exists c : aRc \wedge cRb, \text{ by symmetry} \\ \Downarrow & aRb, \text{ by transitivity} \end{aligned}$$



□