


Mandelstam Variables

Consider any kind of a 2 particles \rightarrow 2 particles process



The 4-momenta p_1^μ , p_2^μ , $p_1'^\mu$, and $p_2'^\mu$ of the 2 incoming and 2 outgoing particles are on-shell satisfy 8 constraints: the on-shell conditions for each particle

$$p_1^2 = m_1^2, \quad p_2^2 = m_2^2, \quad p_1'^2 = m_1'^2, \quad p_2'^2 = m_2'^2 \quad (2)$$

and the net 4-momentum conservation

$$p_1 + p_2 - p_1' - p_2' = 0. \quad (3)$$

Consequently, all Lorentz invariant combinations of the 4 external momenta may be expressed in terms of the particles' masses and 3 Mandelstam's variables

$$\begin{aligned} s &= (p_1 + p_2)^2 = (p_1' + p_2')^2, \\ t &= (p_1 - p_1')^2 = (p_2' - p_2)^2, \\ u &= (p_1 - p_2')^2 = (p_1' - p_2)^2. \end{aligned} \quad (4)$$

Moreover, only 2 out of these 3 variables are independent while their sum has a fixed value

$$s + t + u = m_1^2 + m_2^2 + m_1'^2 + m_2'^2. \quad (5)$$

Indeed,

$$\begin{aligned} s + t + u &= (p_1 + p_2)^2 + (p_1 - p_1')^2 + (p_1 - p_2')^2 \\ &= 3p_1^2 + p_2^2 + p_1'^2 + p_2'^2 + 2(p_1 p_2) - 2(p_1 p_1') - 2(p_1 p_2') \\ &= p_1^2 + p_2^2 + p_1'^2 + p_2'^2 + 2p_1 \times (p_1 + p_2 - p_1' - p_2') = 0 \\ &= p_1^2 + p_2^2 + p_1'^2 + p_2'^2 \\ &= m_1^2 + m_2^2 + m_1'^2 + m_2'^2. \end{aligned} \quad (6)$$

In terms of Mandelstam's variables, Lorentz products of momenta are given by

$$\begin{aligned}
2(p_1 p_2) &= (p_1 + p_2)^2 - p_1^2 - p_2^2 = s - m_1^2 - m_2^2, \\
2(p'_1 p'_2) &= (p'_1 + p'_2)^2 - p_1'^2 - p_2'^2 = s - m_1'^2 - m_2'^2, \\
2(p_1 p'_1) &= p_1^2 + p_1'^2 - (p_1 - p'_1)^2 = m_1^2 + m_1'^2 - t, \\
2(p_2 p'_2) &= p_2^2 + p_2'^2 - (p_2 - p'_2)^2 = m_2^2 + m_2'^2 - t, \\
2(p_1 p'_2) &= p_1^2 + p_2'^2 - (p_1 - p'_2)^2 = m_1^2 + m_2'^2 - u, \\
2(p_2 p'_1) &= p_2^2 + p_1'^2 - (p_2 - p'_1)^2 = m_2^2 + m_1'^2 - u.
\end{aligned} \tag{7}$$

In particular, for an elastic scattering of 2 same-mass particles

$$\begin{aligned}
s + t + u &= 4m^2, \\
2(p_1 p_2) &= 2(p'_1 p'_2) = s - 2m^2, \\
2(p_1 p'_1) &= 2(p_2 p'_2) = 2m^2 - t, \\
2(p_1 p'_2) &= 2(p_2 p'_1) = 2m^2 - u.
\end{aligned} \tag{8}$$

For the future reference, let me give you similar formulae for the $e^- e^+ \rightarrow \mu^- \mu^+$ pair-production,

$$\begin{aligned}
s + t + u &= 2M_\mu^2 + 2m_e^2 \approx 2M_\mu^2, \\
2(p_1 p_2) &= s - 2m_e^2 \approx s, \\
2(p'_1 p'_2) &= s - 2M_\mu^2, \\
2(p_1 p'_1) &= 2(p_2 p'_2) = M_\mu^2 + m_e^2 - t \approx M_\mu^2 - t, \\
2(p_1 p'_2) &= 2(p_2 p'_1) = M_\mu^2 + m_e^2 - u \approx M_\mu^2 - u.
\end{aligned} \tag{9}$$

and for the $e^- e^+ \rightarrow \gamma \gamma$ annihilation process $p_- + p_+ \rightarrow k_1 + k_2$,

$$\begin{aligned}
s + t + u &= 2m_e^2, \\
2(p_- p_+) &= s - 2m_e^2, \\
2(k_1 k_2) &= s, \\
2(p_- k_1) &= 2(p_+ k_2) = m_e^2 - t, \\
2(p_- k_2) &= 2(p_+ k_1) = m_e^2 - u.
\end{aligned} \tag{10}$$