Log-Distance Integrability for Dissipative Flows: BV Selections on Fractal Attractors

Lukas Geiger
Independent Researcher, Bernau im Schwarzwald, Germany
ORCID: 0009-0005-7296-1534*

March 2026

MSC 2020: 37L30, 47H09, 46B20, 35Q30, 26A45
Keywords: nearest-point projection, log-Lipschitz regularity, bounded variation, attractor geometry, dissipative PDEs, Navier–Stokes

Abstract

We introduce a regularity framework for nearest-point projections onto compact attractors of dissipative semiflows in Hilbert spaces. The classical Lipschitz projection theorem (Federer, 1959) requires positive reach—a condition incompatible with fractal attractor geometry. We replace it with a *trajectory log-Lipschitz* condition (TLL), which controls the projection modulus at the geometrically natural scale of the attractor distance rather than the partition mesh. Combined with a *log-distance integrability* condition (LDI), this yields bounded-variation selections along absolutely continuous trajectories—precisely the regularity needed for Gronwall-type arguments in conditional regularity criteria. The framework applies to any dissipative semiflow with a compact attractor of finite fractal dimension, including the 3D Navier–Stokes equations, reaction-diffusion systems, and damped wave equations. Two verifiable sufficient conditions for LDI are identified: bounded variation of the log-distance function, and exponential sublevel time-measure decay.

1 Introduction

Let H be a separable Hilbert space and $\mathcal{A} \subset H$ a compact set—the global attractor of a dissipative semiflow $\{S(t)\}_{t\geq 0}$. A fundamental question in the qualitative theory of dissipative PDEs is the regularity of the nearest-point projection

$$P(x) \in \arg\min_{a\in\mathcal{A}} \|x-a\|_H$$

as a function of time along a trajectory $u(t) = S(t)\,u_0$. If the map $t \mapsto P(u(t))$ is absolutely continuous (AC), Gronwall-type energy estimates close automatically; if it is merely measurable, critical error terms remain uncontrolled.

*AI tools (Claude by Anthropic) were used for mathematical formalization, literature verification, and editorial refinement. All mathematical content was developed, verified, and approved by the author.

The classical theory (Federer [9]) guarantees Lipschitz continuity of the nearest-point projection on the tubular neighbourhood of sets with *positive reach*. However, attractors of dissipative PDEs generically have non-integer fractal dimension and reach zero (cf. [19], Chapter V). This rules out the direct application of Federer's theorem.

The purpose of this note is to introduce a weaker regularity class— *trajectory log-Lipschitz* projections—that is compatible with fractal geometry and still suffices for bounded-variation (BV) control of the projected trajectory. Motivated by log-Lipschitz regularity results for dissipative flows (cf. [4]), we observe that the naive global log-Lipschitz condition

$$\|P(x) - P(y)\| \;\le\; C\,\|x-y\|\left(1+\log\tfrac{R}{\|x-y\|}\right) \tag{1}$$

does *not* imply BV of $P \circ u$ for AC curves u: the partition sums $\sum_i \Delta_i(1+\log(R/\Delta_i))$ diverge logarithmically as the mesh refines (Section 3). The correct formulation replaces the increment-scale weight $\log(R/\|x-y\|)$ with the attractor-distance weight $\log(R/d(t))$, where $d(t) = \operatorname{dist}(u(t), \mathcal{A})$. This yields a trajectory-adapted condition whose BV sums converge whenever a log-distance integrability condition (LDI) holds.

2 Setup and Notation

Throughout, H denotes a separable real Hilbert space with norm $\|\cdot\|$ and inner product $\langle\cdot,\cdot\rangle$.

Definition 2.1 (Dissipative semiflow and attractor)**.** A *dissipative semiflow* on H is a family $\{S(t)\}_{t\ge 0}$ of continuous maps $S(t) : H \to H$ satisfying $S(0) = \mathrm{id}$, $S(t+s) = S(t)\circ S(s)$, and possessing a *compact global attractor* $\mathcal{A} \subset H$: a compact, invariant ($S(t)\mathcal{A} = \mathcal{A}$ for all $t \ge 0$) set that attracts every bounded subset of H.

Definition 2.2 (Tubular neighbourhood and distance)**.** For $R > 0$, the *R-tubular neighbourhood* of $\mathcal{A}$ is

$$\mathcal{A}^R \;:=\; \{x \in H : \operatorname{dist}(x,\mathcal{A}) < R\},$$

where $\operatorname{dist}(x,\mathcal{A}) := \inf_{a\in\mathcal{A}}\|x-a\|$. For a curve $u : [0,T] \to \mathcal{A}^R$, we write $d(t) := \operatorname{dist}(u(t),\mathcal{A})$.

Definition 2.3 (Nearest-point selection)**.** A *nearest-point selection* is a measurable map $P : \mathcal{A}^R \to \mathcal{A}$ satisfying $\|x - P(x)\| = \operatorname{dist}(x,\mathcal{A})$ for all $x \in \mathcal{A}^R$, and $P|_{\mathcal{A}} = \mathrm{id}$. Such a selection exists by the Kuratowski–Ryll-Nardzewski theorem applied to the compact-valued multifunction $x \mapsto \arg\min_{a\in\mathcal{A}}\|x-a\|$.

Proposition 2.4 (Minimal metric derivative selection)**.** *Among the nearest-point selections $P : \mathcal{A}^R \to \mathcal{A}$, there exists a* canonical selection *$u^*(t)$ along any absolutely continuous trajectory $u \in \mathrm{AC}([0,T];\,H)$ defined by*

$$u^*(t) \;:=\; \arg\min_{a\in\mathcal{A},\ \|u(t)-a\|=d(t)} |a|'(t), \qquad |a|'(t) \;:=\; \lim_{h\to 0}\frac{\operatorname{dist}(a,\,P(u(t+h)))}{|h|},$$

that minimises the metric derivative $|u^*|'(t) = \lim_{h\to 0}\|u^*(t+h) - u^*(t)\|/|h|$ among all nearest-point selections. This selection is a.e. unique: for a.e. $t \in [0,T]$, the minimiser of the metric derivative over the (compact) nearest-point set $\{a \in \mathcal{A} : \|u(t)-a\| = d(t)\}$ is unique.

Proof. The nearest-point set $\Pi(t) := \{a \in \mathcal{A} : \|u(t) - a\| = d(t)\}$ is compact and non-empty for each t. The metric derivative functional $a \mapsto \limsup_{h\to 0} \|P_a(u(t+h)) - a\|/|h|$ is lower semicontinuous on the compact set $\Pi(t)$, hence attains its minimum. For a.e. t at which $|u'|(t)$ exists, the nearest-point set has a unique element minimising the metric derivative: if two distinct minimisers $a_1, a_2 \in \Pi(t)$ existed on a set of positive measure, convexity of the norm would yield $\|(a_1 + a_2)/2 - u(t)\| < d(t)$ for the midpoint (strict convexity of the Hilbert norm), contradicting $a_1, a_2 \in \mathcal{A}$. Hence uniqueness holds a.e. □

Remark 2.5. The canonical selection u^* is the natural candidate for TLL (Definition 4.1): among all measurable nearest-point selections, it has the smallest metric derivative and hence the best chance of satisfying the log-Lipschitz bound. In the Limitations section, we noted that different selections may have different regularity; the minimal metric derivative selection is the "best" one.

3 Why Global Log-Lipschitz Fails

We first show that the global log-Lipschitz condition (1), even combined with log-distance integrability, does not directly yield BV of $P \circ u$.

Proposition 3.1 (Failure of the global LL bound to control variation)**.** *Let $P : \mathcal{A}^R \to \mathcal{A}$ satisfy (1) with constant $C > 0$. Let $u : [0,1] \to \mathcal{A}^R$ be Lipschitz with $\|u'(t)\| = 1$ a.e. For the uniform partition $t_i = i/n$ $(i = 0, \dots, n)$, the upper bound obtained from (1) satisfies*

$$\sum_{i=0}^{n-1} \|P(u(t_{i+1})) - P(u(t_i))\| \;\le\; C\,(1 + \log(nR)),$$

which diverges as $n \to \infty$. In particular, the global LL condition does not provide a finite a priori *bound on* $\mathrm{Var}(P \circ u)$.

Proof. Each increment satisfies $\Delta_i := \|u(t_{i+1}) - u(t_i)\| = 1/n$. By (1):

$$\sum_{i=0}^{n-1} \|P(u(t_{i+1})) - P(u(t_i))\| \;\le\; C \sum_{i=0}^{n-1} \frac{1}{n}\Big(1 + \log(nR)\Big) \;=\; C\,(1 + \log(nR)).$$

As $n \to \infty$, this grows as $C \log n \to \infty$. Hence the bound derived from (1) diverges with the mesh refinement and does not yield a finite upper bound on $\mathrm{Var}(P \circ u)$. (Whether the variation itself is infinite depends on the specific geometry of $\mathcal{A}$; the point is that the global LL condition is insufficient as a *tool* for establishing BV.) □

Remark 3.2. The divergence is *logarithmic*, not polynomial—much slower than for Hölder maps (where $\sum \Delta_i^\alpha$ converges for $\alpha < 1$). This makes global LL "almost" sufficient for BV, but the logarithmic accumulation over infinitely many partition points defeats it. The trajectory log-Lipschitz condition avoids this by anchoring the log weight at the attractor distance $d(t)$, which does not depend on the partition.

4 The Trajectory Log-Lipschitz Condition

Definition 4.1 (Trajectory log-Lipschitz projection (TLL))**.** Let $u \in \mathrm{AC}([0,T]; H)$ with $u(t) \in \mathcal{A}^R$ for all t, and let $P : \mathcal{A}^R \to \mathcal{A}$ be a nearest-point selection. We say that

P satisfies the *trajectory log-Lipschitz condition* (TLL) along u if there exist constants $C_{\mathrm{TLL}} > 0$ and $h_0 > 0$ such that for all $t \in [0, T]$ with $d(t) > 0$ and all $h \in \mathbb{R}$ with $0 < |h| < h_0$ and $t + h \in [0, T]$ (i.e., both forward and backward increments):

$$\|P(u(t+h)) - P(u(t))\| \;\le\; C_{\mathrm{TLL}}\, \|u(t+h) - u(t)\|\, \Omega(t), \tag{TLL}$$

where $\Omega(t) := 1 + \log(R/d(t))$ is the *log-distance weight*.

On the set $\{t : d(t) = 0\}$ (where $u(t) \in \mathcal{A}$), $P(u(t)) = u(t)$ by idempotence of P. For the transition $d(t) = 0$, $d(t+h) > 0$, we have $\|P(u(t+h)) - u(t)\| = \|P(u(t+h)) - P(u(t))\|$. Since $u(t) \in \mathcal{A}$ is a candidate for the nearest point of $u(t+h)$, the minimality of P gives $\|u(t+h) - P(u(t+h))\| \le \|u(t+h) - u(t)\|$. Together with the triangle inequality:

$$\|P(u(t+h)) - u(t)\| \;\le\; \|P(u(t+h)) - u(t+h)\| + \|u(t+h) - u(t)\| \;\le\; 2\,\|u(t+h) - u(t)\|.$$

This gives $|v'|(t) \le 2\,|u'|(t)$ on $\{d = 0\}$.

Remark 4.2 (Convention $C_{\mathrm{TLL}} \ge 2$). The factor 2 on $\{d = 0\}$ is absorbed by the TLL bound provided $C_{\mathrm{TLL}} \ge 2$ and $\Omega(t) \ge 1$. Throughout, we adopt the convention $C_{\mathrm{TLL}} := \max(C_{\mathrm{TLL}}, 2)$; this changes the BV bound (2) by at most a factor of 2 and does not affect the integrability of (LDI).

Remark 4.3 (Geometric interpretation of TLL). The condition TLL asserts that the nearest-point projection, evaluated along the trajectory u, has a local Lipschitz constant that blows up *at most logarithmically* as the trajectory approaches the attractor:

$$\mathrm{Lip}_{\mathrm{loc}}(P \circ u;\, t) \;\lesssim\; 1 + \log \frac{R}{d(t)}.$$

At distance $d(t) \sim R$ from the attractor, the projection is essentially Lipschitz (the log term is $O(1)$). As $d(t) \to 0$, the log weight grows, reflecting the increasing geometric complexity of the attractor boundary at fine scales. The logarithmic rate is the *critical threshold*: faster growth (e.g., polynomial) would make LDI fail generically, while slower growth (bounded) would imply Lipschitz projection—too strong for fractal geometry.

Remark 4.4 (TLL as Aubin property of the projection correspondence). The TLL condition admits a natural reformulation in the language of variational analysis. Define the *projection correspondence*

$$\Gamma : t \;\mapsto\; \Big\{a \in \mathcal{A} : \|a - u(t)\| = d(t)\Big\}$$

mapping each time to the set of nearest attractor elements. The TLL condition $|d'(t)| \le C/|\log d(t)|$ (i.e., the metric derivative of the distance function is controlled by the inverse logarithm) is equivalent to the *Aubin property* (metric regularity) of the correspondence Γ at rate $1/|\log r|$. Specifically, Γ is metrically regular at rate $\phi(r) = 1/|\log r|$ if

$$\mathrm{dist}\Big(a,\, \Gamma(t')\Big) \;\le\; L\, \phi\Big(\|u(t') - u(t)\|\Big)\, \|u(t') - u(t)\|$$

for $a \in \Gamma(t)$ and t' near t. This reformulation connects the projection regularity theory to the Aubin criterion and metric regularity framework of Rockafellar–Wets [16] and Dontchev–Rockafellar [6]. The key advantage is that Aubin-property criteria are well-developed for set-valued maps and provide verifiable sufficient conditions (e.g., via the Mordukhovich coderivative) that may be applicable to the attractor projection.

Remark 4.5 (TLL vs global LL). The global log-Lipschitz condition (1) does *not* imply TLL. For $h \to 0$ at fixed t:

$$\log \frac{R}{\|u(t+h) - u(t)\|} \;\approx\; \log \frac{R}{h\,|u'(t)|} \;=\; \log \frac{R}{|u'(t)|} + \log \frac{1}{h} \;\to\; \infty,$$

while $\log(R/d(t))$ remains fixed. The global condition produces an *increment-scale* log weight that diverges with the partition mesh; TLL produces a *distance-scale* log weight that depends only on the current geometry. This is the structural reason why TLL yields BV while global LL does not.

5 Log-Distance Integrability

Definition 5.1 (Log-distance integrability (LDI))**.** In the setting of Definition 4.1, we say that *log-distance integrability* holds if

$$\int_{\{d>0\}} \|u'(t)\|\,\Omega(t)\,\mathrm{d}t \;=\; \int_{\{d>0\}} \|u'(t)\| \left(1 + \log \tfrac{R}{d(t)}\right) \mathrm{d}t \;<\; \infty. \tag{LDI}$$

On the set $\{t : d(t) = 0\}$, $v(t) = u(t)$ and the BV contribution is controlled by $\int_{\{d=0\}} \|u'(t)\|\,\mathrm{d}t < \infty$ (since u is AC), so only the region $\{d > 0\}$ is relevant for (LDI). To state the BV bound (2) uniformly over all of $[0, T]$, we extend Ω to $\{d = 0\}$ by setting $\Omega(t) := 1$ there (consistent with the factor-2 bound of Remark 4.2).

Remark 5.2 (Minimality of LDI). The condition LDI is positioned precisely between the known regularity classes:

- **Lipschitz projection** ($\Omega \equiv 1$): LDI reduces to $\int \|u'\| < \infty$, which is automatic for AC curves. This is the Federer/positive-reach regime.
- **Hölder projection** ($\Omega(t) \sim d(t)^{-\beta}$, $\beta > 0$): LDI requires $\int \|u'\|\, d(t)^{-\beta} < \infty$, which generically fails near the attractor. This is why Mané projectors (Hölder inverse) do not yield BV.
- **Log-Lipschitz projection** (TLL): LDI requires $\int \|u'\| \log(1/d) < \infty$, which is the borderline case—logarithmic singularities are integrable under mild dynamical conditions.

The log-Lipschitz/LDI pair thus occupies the *critical* regularity window for BV selections on fractal attractors.

6 Main Result: BV Chain Rule

Theorem 6.1 (BV chain rule for log-Lipschitz projections)**.** *Let $\{S(t)\}_{t\geq 0}$ be a dissipative semiflow on a separable Hilbert space H with compact global attractor $\mathcal{A} \subset H$. Let $u \in \mathrm{AC}([0,T];\, H)$ with $u(t) \in \mathcal{A}^R$ for all t, and let $P : \mathcal{A}^R \to \mathcal{A}$ be a nearest-point selection satisfying TLL (TLL) along u. Set $v(t) := P(u(t))$.*

If LDI (LDI) holds, then $v \in \mathrm{BV}([0,T];\, H)$ with the explicit bound

$$\mathrm{Var}_H(v;\, [0,T]) \;\leq\; C_{\mathrm{TLL}} \int_0^T \|u'(t)\|\,\Omega(t)\,\mathrm{d}t, \tag{2}$$

where $\Omega(t) := 1 + \log(R/d(t))$ on $\{d > 0\}$ and $\Omega(t) := 1$ on $\{d = 0\}$ (Remark 4.2).

Proof. The proof has three steps.

Step 1 (Metric derivative bound). For any continuous curve $v : [0,T] \to H$, the *upper metric derivative* at t is

$$|v'|^+(t) \;:=\; \limsup_{h\to 0} \frac{\|v(t+h)-v(t)\|}{|h|} \;\in\; [0,+\infty].$$

This quantity exists everywhere (as a value in $[0,+\infty]$) and is Borel measurable. For absolutely continuous curves the lim sup is a genuine limit a.e. (cf. [1, Theorem 1.1.2]). Since u is AC, $|u'|(t) := \lim_{h\to 0} \|u(t+h)-u(t)\|/|h|$ exists for a.e. t.

Case $d(t) > 0$: Dividing (TLL) by $|h|$ and taking lim sup:

$$|v'|^+(t) \;\le\; C_{\mathrm{TLL}}\, |u'|(t)\, \Omega(t) \qquad \text{for a.e. } t \text{ with } d(t) > 0. \tag{3}$$

(The right-hand side is a genuine limit; the left-hand side may be a lim sup, which is dominated by the same bound.)

Case $d(t) = 0$ (interior of $\mathcal{Z}$): On $\mathcal{Z} := \{t \in [0,T] : d(t) = 0\}$, $u(t) \in \mathcal{A}$ and $v(t) = P(u(t)) = u(t)$. Since $u(t) \in \mathcal{A}$ is a candidate minimiser for $\mathrm{dist}(u(t+h), \mathcal{A})$, the triangle inequality gives $\|P(u(t+h)) - u(t)\| \le 2\,\|u(t+h) - u(t)\|$ (cf. the bound in Definition 4.1), so $|v'|^+(t) \le 2\,|u'|(t)$.

Boundary points of $\mathcal{Z}$: At $t_0 \in \partial\mathcal{Z}$ with $d(t_0) = 0$, sequences $t_n \to t_0$ with $d(t_n) > 0$ may exist. For continuity of v at t_0: $\|v(t_n) - v(t_0)\| = \|P(u(t_n)) - u(t_0)\| \le d(t_n) + \|u(t_n) - u(t_0)\| \to 0$, since u is continuous and $d = \mathrm{dist}(\cdot, \mathcal{A}) \circ u$ inherits continuity (the distance function is 1-Lipschitz). The metric derivative bound $|v'|^+(t_0) \le 2\,|u'|(t_0)$ remains valid by the same triangle inequality argument as above.

Since $\Omega(t) \ge 1$ and $C_{\mathrm{TLL}} \ge 2$ (Remark 4.2), the bound (3) holds on all of $[0,T]$ (with $|v'|^+$ in place of $|v'|$ where the limit may not exist; this suffices for the BV criterion in Step 3).

Step 2 (Continuity of v). The nearest-point projection on a compact (but non-convex) set is in general only upper semicontinuous as a set-valued map and need not be single-valued everywhere. However, the TLL condition (TLL) directly implies continuity of $v = P \circ u$ along the trajectory: for $d(t) > 0$, $\|v(t+h) - v(t)\| \le C_{\mathrm{TLL}}\, \|u(t+h) - u(t)\|\, \Omega(t) \to 0$ as $h \to 0$; for $d(t) = 0$, $\|v(t+h) - v(t)\| \le 2\,\|u(t+h) - u(t)\| \to 0$ (including boundary points of $\mathcal{Z}$, cf. Step 1). Hence v is continuous on $[0,T]$.

Step 3 (BV bound). We show $\|v(b) - v(a)\| \le \int_a^b g(t)\,\mathrm{d}t$ for all $0 \le a < b \le T$, where $g(t) := C_{\mathrm{TLL}}\, |u'|(t)\, \Omega(t) \in L^1$ (by LDI). Since v is continuous (Step 2), the BV bound $\mathrm{Var}(v) \le \int_0^T g$ then follows by summing over any partition.

Set $f(t) := \|v(t) - v(a)\|$. Then $f : [a,b] \to \mathbb{R}$ is continuous and real-valued, $f(a) = 0$, and the upper Dini derivative satisfies $D^+ f(t) \le |v'|^+(t)$ (since $f(t+h) - f(t) \le \|v(t+h) - v(t)\|$ by the reverse triangle inequality, and division by $h > 0$ followed by lim sup gives $D^+ f \le |v'|^+$). By Step 1, $|v'|^+(t) \le g(t)$ a.e. A classical result on Dini derivatives (see, e.g., [18, Chapter VII, §§ 9–10]; cf. also [2, Chapter 3]) gives: for a continuous function $f : [a,b] \to \mathbb{R}$ with $D^+ f(t) \le g(t)$ a.e. and $g \in L^1$, the inequality $f(b) - f(a) \le \int_a^b g(t)\,\mathrm{d}t$ holds. Therefore

$$\|v(b) - v(a)\| \;=\; f(b) \;\le\; \int_a^b g(t)\,\mathrm{d}t \;\le\; \int_0^T g(t)\,\mathrm{d}t.$$

Summing over any partition $0 = t_0 < \cdots < t_n = T$:

$$\sum_i \|v(t_{i+1}) - v(t_i)\| \;\le\; \sum_i \int_{t_i}^{t_{i+1}} g(t)\,\mathrm{d}t \;=\; \int_0^T g(t)\,\mathrm{d}t \;=\; C_{\mathrm{TLL}} \int_0^T |u'|(t)\,\Omega(t)\,\mathrm{d}t,$$

which gives (2).

The bound (2) follows from Steps 1–3. □

Remark 6.2 (Direct partition-sum verification). For a fixed partition with mesh $< h_0$, the finiteness of the partition sum can be seen directly from TLL without the Dini-derivative machinery: on each subinterval, TLL gives $\|v(t_{i+1}) - v(t_i)\| \le C_{\mathrm{TLL}} \|u(t_{i+1}) - u(t_i)\| \, \Omega(t_i)$ (or the factor-2 bound on $\{d = 0\}$, absorbed by $C_{\mathrm{TLL}} \ge 2$). The resulting partition sum $\sum_i C_{\mathrm{TLL}} \, \Omega(t_i) \int_{t_i}^{t_{i+1}} \|u'\| \, \mathrm{d}t$ is finite for each fixed partition (since Ω is bounded on any compact set disjoint from $\{d = 0\}$, and the $\{d = 0\}$ contribution is controlled by $2 \int \|u'\|$). However, the passage to the supremum over *all* partitions—which is the definition of $\mathrm{Var}(v)$—requires the Dini-derivative argument of Step 3, which sharpens the pointwise $\Omega(t_i)$-weight to the integrated $\Omega(t)$-weight.

Remark 6.3 (Banach space generality). The proof of Theorem 6.1 uses only the norm, compactness of $\mathcal{A}$, and measurable selection—no inner product or Hilbert space structure. The BV chain rule therefore holds verbatim in any separable Banach space. We restrict to Hilbert spaces only for the PDE applications in Section 8.

7 Sufficient Conditions for LDI

The condition LDI (LDI) is a single integrability requirement. We provide two independent sufficient conditions, each verifiable from dynamical information about the semiflow.

7.1 Variant A: Log-distance has bounded variation

Proposition 7.1 (BV log-distance implies LDI)**.** *Suppose* $d(t) > 0$ *for a.e.* $t \in [0, T]$ *and* $\log(R/d(t)) \in \mathrm{BV}([0, T])$. *Then LDI holds.*

Proof. Write $\ell(t) := 1 + \log(R/d(t))$. Since $u' \in L^1(0, T; H)$ (as u is AC) and $\ell \in \mathrm{BV}([0, T]) \subset L^\infty([0, T])$, the product $\|u'(t)\| \, \ell(t)$ is locally integrable:

$$\int_0^T \|u'(t)\| \, \ell(t) \, \mathrm{d}t \;\le\; \|\ell\|_{L^\infty(0,T)} \int_0^T \|u'(t)\| \, \mathrm{d}t \;<\; \infty. \qquad \square$$

Remark 7.2 (When is log-distance BV?). The condition $\log(R/d) \in \mathrm{BV}$ requires that the trajectory does not oscillate infinitely often between different distance scales from the attractor. Formally, it excludes infinite "zickzack frequency" in $d(t)$. For monotonically attracting trajectories ($d(t)$ eventually decreasing), this is automatic. For trajectories with transient oscillations, BV of $\log d$ follows if the total number of scale crossings $\{t : d(t) = \varepsilon\}$ is finite for each $\varepsilon > 0$—a generic property of dissipative semiflows with isolated equilibria.

7.2 Variant B: Sublevel time-measure control

Definition 7.3 (Sublevel-time control (STC))**.** The trajectory u satisfies *sublevel-time control* with exponent $\alpha > 0$ if there exists $C_* > 0$ such that

$$\Big|\{t \in [0, T] : d(t) \le \varepsilon\}\Big| \;\le\; C_* \, \varepsilon^\alpha \qquad \text{for all } \varepsilon \in (0, R]. \tag{STC}$$

Remark 7.4 (STC implies $|\{d = 0\}| = 0$). Taking $\varepsilon \to 0$ in (STC) shows $|\{d = 0\}| = 0$. Hence STC is only applicable to trajectories that do not spend positive time on the attractor. This is the generic case for non-stationary solutions approaching $\mathcal{A}$ asymptotically. For trajectories that reach the attractor in finite time (e.g., convergence to an equilibrium with $u(t) \in \mathcal{A}$ for $t \geq t_1$), Variant A (BV log-distance) is the appropriate sufficient condition.

Proposition 7.5 (STC implies LDI)**.** *If STC* (STC) *holds with some* $\alpha > 0$ *and* $u' \in L^2(0, T; H)$*, then* $\log(R/d) \in L^p(0, T)$ *for every* $p < \infty$*, and LDI holds.*

Proof. **Step 1 (Tonelli representation).** Since $u(t) \in \mathcal{A}^R$, we have $d(t) \leq R$ for all t. On the set $\{t : d(t) = 0\}$ (where $u(t) \in \mathcal{A}$), the integrand $\|u'(t)\| \log(R/d(t))$ is $+\infty$; we set the contribution to zero by convention (on this set, $v(t) = u(t)$ and the BV chain rule bound holds trivially). On $\{t : d(t) > 0\}$, using the layer-cake identity $\log(R/d(t)) = \int_{d(t)}^{R} s^{-1}\,\mathrm{d}s$ and Tonelli:

$$\int_{\{d>0\}} \|u'(t)\| \log \frac{R}{d(t)}\,\mathrm{d}t \;=\; \int_0^R \frac{1}{s}\Big(\int_{\{0<d(t)\leq s\}} \|u'(t)\|\,\mathrm{d}t\Big)\,\mathrm{d}s. \tag{4}$$

Step 2 (Hölder on inner integral). By Hölder's inequality:

$$\int_{\{0<d(t)\leq s\}} \|u'(t)\|\,\mathrm{d}t \;\leq\; \Big(\int_0^T \|u'(t)\|^2\,\mathrm{d}t\Big)^{1/2} \cdot \Big|\{0 < d \leq s\}\Big|^{1/2} \;\leq\; \|u'\|_{L^2}\, C_*^{1/2}\, s^{\alpha/2},$$

where the last step uses $|\{0 < d \leq s\}| \leq |\{d \leq s\}| \leq C_*\, s^\alpha$ from STC.

Step 3 (Integrability near $s = 0$). Substituting into (4):

$$\int_0^R \frac{1}{s}\,\|u'\|_{L^2}\, C_*^{1/2}\, s^{\alpha/2}\,\mathrm{d}s \;=\; \|u'\|_{L^2}\, C_*^{1/2} \int_0^R s^{\alpha/2-1}\,\mathrm{d}s \;=\; \|u'\|_{L^2}\, C_*^{1/2}\, \frac{2}{\alpha}\, R^{\alpha/2} \;<\; \infty.$$

The integral converges because $\alpha > 0$, so $s^{\alpha/2-1}$ is integrable near $s = 0$.

Step 4 (L^p bounds). The same Tonelli argument with Hölder exponents p and q ($1/p + 1/q = 1$) gives $\|\log(R/d)\|_{L^p} \leq C(p, \alpha, R, C_*)$ for every $p < \infty$, so LDI follows from any finite L^q bound on $\|u'\|$. □

Remark 7.6 (Sources of STC in dissipative flows). Sublevel-time control (STC) is the genuine new analytical input. Three mechanisms can generate it:

(i) **Exponential tracking:** If $d(t) \leq C_0\, e^{-\lambda t}$ for $t \geq t_0$, then $\{d \leq \varepsilon\} \subset \{t \geq (\log(C_0/\varepsilon))/\lambda\}$, whose measure is $O(\log(1/\varepsilon))$—much smaller than $C_*\, \varepsilon^\alpha$ for any $\alpha > 0$. So exponential tracking implies STC with *any* exponent.

(ii) **Lyapunov dissipation:** If a Lyapunov functional E satisfies $-\dot{E} \geq \Psi(d)$ with Ψ not too flat near $d = 0$ (e.g. $\Psi(d) \geq c\, d^\beta$ for some β), then sublevel times are controlled by the total energy budget $E(0) - \inf E$.

(iii) **No fast oscillation:** Energetic constraints (finite dissipation integral) prevent arbitrarily rapid back-and-forth motion near the attractor, bounding the number of excursions into $\{d \leq \varepsilon\}$.

Each mechanism yields STC with different exponents α. The STC condition is physically plausible for all standard classes of dissipative PDEs.

Lemma 7.7 (Exponential tracking implies log-integrability)**.** *If $d(t) \le C\,e^{-\lambda t}$ for some $\lambda > 0$ and $C > 0$, then $\log(1/d) \in L^p([0,T])$ for all $p < \infty$ and all $T > 0$, and the LDI condition is satisfied with*

$$\int_0^T \log \frac{1}{d(t)}\,\mathrm{d}t \;\le\; \lambda T + \log(1/C)\cdot T.$$

In particular, STC holds with any exponent $\alpha > 0$, and the LDI integral is bounded linearly in T.

Proof. From $d(t) \le C\,e^{-\lambda t}$ we obtain $\log(1/d(t)) \ge \lambda t - \log C$, and also

$$\log \frac{1}{d(t)} \;\le\; \log \frac{1}{C\,e^{-\lambda t}} \;=\; \lambda t + \log \frac{1}{C}.$$

Hence

$$\int_0^T \log \frac{1}{d(t)}\,\mathrm{d}t \;\le\; \int_0^T \Big(\lambda t + \log(1/C)\Big)\,\mathrm{d}t \;=\; \frac{\lambda T^2}{2} + \log(1/C)\cdot T.$$

For the LDI condition (LDI), with $u' \in L^2$:

$$\int_0^T \|u'(t)\| \log \frac{R}{d(t)}\,\mathrm{d}t \;\le\; \|u'\|_{L^2} \left(\int_0^T \Big(\log(R/C) + \lambda t\Big)^2\,\mathrm{d}t\right)^{1/2} \;<\; \infty$$

by Hölder's inequality, since the right factor is a polynomial in T. The L^p integrability $\log(1/d) \in L^p([0,T])$ for all $p < \infty$ follows since $\log(1/d(t)) \le \lambda t + \log(1/C)$, which is in $L^\infty([0,T])$ for each finite T. □

Remark 7.8 (Exponential tracking as the strongest STC regime)*.* Lemma 7.7 shows that exponential tracking is the "easiest" dynamical regime for LDI: the log-distance is bounded by a linear function of time, making LDI automatic for any trajectory with finite energy ($u' \in L^2$). This is the regime expected for strong solutions of the 3D Navier–Stokes equations conditional on global regularity (Example 8.1). The non-trivial case is when exponential tracking fails—e.g., near potential blow-up—where the weaker STC condition (polynomial sublevel decay) or Variant A (BV log-distance) become the relevant tools.

8 Applications to Dissipative PDEs

We illustrate how the abstract framework applies to concrete PDE settings. In each case, the semiflow possesses a compact global attractor, and the question is whether TLL and LDI hold.

8.1 3D Navier–Stokes on $\mathbb{T}^3$

Example 8.1 (Navier–Stokes equations)*.* Let $H = L^2(\mathbb{T}^3;\mathbb{R}^3)$ (divergence-free), $\mathcal{A}$ the global attractor (in the sense of Temam [19]) with smooth forcing f. By Temam [19]:

- $\mathcal{A}$ is compact in H^1 (hence in H).
- Every $a \in \mathcal{A}$ is C^∞.

- Exponential tracking holds *conditional on global regularity*: for strong solutions existing on $[0, \infty)$, $d(t) \le C_0\, e^{-\lambda t}$ for $t \ge t_0$. (For Leray–Hopf weak solutions, only algebraic attraction in L^2 is known unconditionally.)

The attractor has finite fractal dimension $d_f(\mathcal{A}) \le c_1\, G^{2/3}(1 + \log G)^{1/3}$ (where G is the Grashof number; cf. [7]), but generically has reach 0—ruling out Federer's theorem.

Status of TLL: Open. The squeezing property of the Navier–Stokes semiflow (cf. [10], Chapter 14) provides exponential contraction of high-frequency differences on the attractor. Combined with Mané projector estimates, this suggests that the nearest-point projection has at most log-Lipschitz modulus along trajectories—but a rigorous proof requires quantitative control of the projection stability in the high-mode complement $Q_N H$.

Status of LDI: Follows from exponential tracking on $[t_0, T]$ for any fixed $T < \infty$ (Remark 7.6). The critical question is the blow-up regime $T \nearrow T^*$, where $\|u'\| \to \infty$. The BV bound (2) yields Assumption G′ of [11], which implies global regularity via the attractor-distance argument.

Connection to the log-Dirichlet quotient: The companion paper [11] introduces the *log-Dirichlet quotient* $Q_{\mathrm{LD}}(t) = \|\nabla(u - v)\|^2/\|u - v\|^2$ and shows that its integrability transforms the Gronwall estimate from exponential to algebraic growth. The LDI condition of the present paper provides precisely the mechanism that ensures Q_{LD} integrability: the BV chain rule controls the composition $\log d \circ u$, which bounds the log-Dirichlet quotient from above.

8.2 Reaction-diffusion systems

Example 8.2 (Reaction-diffusion on bounded domains). Consider $\partial_t u = D\Delta u + f(u)$ on a bounded domain $\Omega \subset \mathbb{R}^n$ with Dirichlet or Neumann boundary conditions, where $D > 0$ is a diffusion matrix and f satisfies a dissipativity condition $\langle f(u), u\rangle \le C - c|u|^2$. The semiflow possesses a compact global attractor $\mathcal{A} \subset L^2(\Omega)$ with finite fractal dimension [19].

For scalar equations ($u \in \mathbb{R}$) in one space dimension with a gradient nonlinearity $f(u) = -V'(u)$, the attractor consists of equilibria and heteroclinic orbits, and the nearest-point projection is Lipschitz (in fact, the attractor has a Lipschitz-graph structure as an inertial manifold; cf. [14]). TLL and LDI hold trivially.

For systems in higher dimensions, the attractor may have non-integer fractal dimension and reach zero. The log-Lipschitz framework then provides a path to BV selections that Federer's theorem does not. The spectral gap condition for inertial manifolds is satisfied for many reaction-diffusion systems [8], making the attractor geometry more regular than in the Navier–Stokes case.

8.3 Damped wave equations

Example 8.3 (Damped semilinear wave equation). Consider $\partial_{tt} u + \gamma\, \partial_t u = \Delta u + f(u)$ on $\Omega \subset \mathbb{R}^3$ with $\gamma > 0$ and subcritical nonlinearity. The semiflow on $H_0^1 \times L^2$ possesses a compact global attractor with finite fractal dimension (cf. [3]). The exponential tracking property holds with rate $\lambda = \gamma/2$ (spectral gap of the damped wave operator; cf. [3], Chapter V).

The attractor geometry is intermediate between reaction-diffusion (where inertial manifolds often exist) and Navier–Stokes (where they do not). The log-Lipschitz framework

provides a *uniform* approach applicable across all three settings, with TLL as the single geometric input.

9 Discussion

9.1 The new mathematical interface

The central contribution of this note is the identification of the *critical regularity interface* for BV selections on fractal attractors. The interface is characterised by a single question:

> **Core Question.** For a dissipative semiflow with compact attractor $\mathcal{A} \subset H$, does the nearest-point projection satisfy trajectory log-Lipschitz control (TLL)?

This is a question about the *regularity of the attractor as seen from the trajectory*, not about the global geometry of $\mathcal{A}$. The distinction is crucial: global geometric conditions (positive reach, inertial manifold) impose structure on the *entire* attractor, while TLL requires regularity only in the direction of the dynamical approach.

9.2 Relation to existing regularity theories

Condition	Projection	Composition	Fractal?	Source
Positive reach > 0	Lipschitz-1	AC $\Rightarrow$ AC	No	Federer [9]
Prox-regular	locally Lipschitz	AC $\Rightarrow$ AC (local)	No	Poliquin et al. [17]
Hausdorff-BV set map	BV selection exists	—	Yes	Chistyakov [5]
Mané projector	Hölder^{-1}	AC $\Rightarrow$ Hölder	Yes	Mané [15]
TLL + LDI	**log-Lipschitz**	**AC $\Rightarrow$ BV**	**Yes**	**This note**

The TLL+LDI framework is the only entry that simultaneously (i) accommodates fractal geometry (non-integer dimension, reach 0), (ii) yields BV regularity (sufficient for Gronwall-type arguments), and (iii) is formulated as a trajectory-level condition (independent of global attractor structure).

9.3 Limitations

- **Selection dependence.** The TLL condition is formulated for a *given* measurable nearest-point selection P. On non-convex compact sets, the nearest-point map is generically multi-valued, and different selections may have different regularity. The framework requires the existence of *at least one* selection satisfying TLL; it does not claim that every measurable selection does so. In practice, the "best" selection (e.g., the one minimising the metric derivative) is the natural candidate.

- **TLL is open for 3D NS.** The squeezing property and Mané projector estimates suggest log-Lipschitz stability of the projection, but a rigorous proof requires quantitative bounds on the high-mode projection complement that are not currently available.

- **LDI is non-trivial only under blow-up.** For smooth solutions on compact time intervals, LDI holds automatically. The condition becomes critical only in the limit $T \nearrow T^*$ (blow-up time), where $\|u'\| \to \infty$.
- **BV, not AC.** The framework yields BV regularity, which suffices for Stieltjes-integral-based Gronwall arguments but not for pointwise differential inequalities. Upgrading BV to AC would require Lipschitz projection (positive reach)—exactly the condition we are trying to avoid.

10 Conclusion

We have introduced a regularity framework for nearest-point projections onto compact attractors that occupies the critical gap between Hölder (insufficient for BV) and Lipschitz (incompatible with fractal geometry). The trajectory log-Lipschitz condition (TLL) controls the projection modulus at the attractor distance scale, and the log-distance integrability condition (LDI) ensures that the resulting BV sums converge.

The framework reduces conditional regularity criteria for dissipative PDEs—including the 3D Navier–Stokes equations—to a single geometric-dynamical question: does the nearest-point projection have at most logarithmic blowup along dissipative trajectories? This question is independent of the specific PDE and depends only on the interplay between attractor geometry and the semiflow's contraction properties.

11 Time-averaged selection and BV regularity

The central obstruction to closing the regularity gap is the BV control of the nearest-point selection $v(t) \in \arg\min_{a\in\mathcal{A}} \|u(t) - a\|$. We propose replacing the pointwise selection by a *time-averaged* selection that is automatically BV-regular.

Definition 11.1 (Moreau–Yosida regularised selection)**.** For $\varepsilon > 0$, define the *time-averaged nearest-point selection*

$$v_\varepsilon(t) \;:=\; \arg\min_{a\in\mathcal{A}} \int_{t-\varepsilon}^{t+\varepsilon} \|u(s) - a\|^2 \, ds.$$

Proposition 11.2 (Automatic BV regularity)**.** *Under the standing assumptions (finite-dimensional attractor, energy dissipation bound), the regularised selection v_ε satisfies:*

(i) Uniqueness: *$v_\varepsilon(t)$ is unique for a.e. t (strict convexity of the time-integrated cost).*

(ii) BV bound: $\mathrm{Var}(v_\varepsilon; [0,T]) \;\le\; C_\varepsilon \int_0^T \|\partial_t u(s)\| \, ds \;<\; \infty$.

(iii) TLL inheritance: *If u satisfies TLL with constant C_{TLL}, then v_ε satisfies TLL with constant $C_{\mathrm{TLL}} + O(\varepsilon)$.*

(iv) Gronwall compatibility: *The Gronwall estimates of the main theorem (Theorem 6.1) remain valid for v_ε with an additional error term $O(\varepsilon \, \|\partial_t u\|_{L^2})$.*

Proof sketch. (i) The map $a \mapsto \int_{t-\varepsilon}^{t+\varepsilon} \|u(s) - a\|^2 \, ds$ is strictly convex: the integrand is strictly convex for a.e. s (since $u(s) \neq a$ for a.e. s and $a \in \mathcal{A}$), and integration preserves strict convexity.

(ii) The first-variation condition for the minimiser yields

$$\|v_\varepsilon(t+h) - v_\varepsilon(t)\| \;\le\; \frac{1}{2\varepsilon}\int_{t-\varepsilon}^{t+\varepsilon} \|u(s+h) - u(s)\|\, ds \;\le\; \frac{h}{2\varepsilon}\int_{t-\varepsilon}^{t+\varepsilon} \|\partial_t u(s)\|\, ds.$$

Integrating over $t \in [0,T]$ and applying Fubini gives the BV bound with $C_\varepsilon = T/(2\varepsilon)$.

(iii) The TLL constant degrades by the time-averaging window:

$$|\log d_\varepsilon(t+h) - \log d_\varepsilon(t)| \;\le\; C_{\mathrm{TLL}}\, |\log |h|| \;+\; C\,\varepsilon/|h|,$$

where the second term arises from the mismatch between pointwise and averaged distance.

(iv) The Gronwall estimate picks up an additional forcing term from the time-averaging:

$$\frac{d}{dt} H(u \mid v_\varepsilon) \;\le\; -\alpha\, H + \beta + \gamma\,\varepsilon,$$

where γ depends on $\|\partial_t u\|$ but not on H. The standard Gronwall lemma applies; the additional $\gamma\varepsilon$ term vanishes in the limit $\varepsilon \to 0$. □

Remark 11.3 (Passage to the limit $\varepsilon \to 0$). The key question is whether the BV bound survives the limit $\varepsilon \to 0$:

- If $\sup_\varepsilon \mathrm{Var}(v_\varepsilon; [0,T]) < \infty$, then a subsequence v_{ε_k} converges in L^1 to a BV selection v_0, which is the *canonical* nearest-point selection.
- The uniform BV bound holds if and only if the attractor $\mathcal{A}$ has a "time-averaged reach" $\mathrm{reach}_{\mathrm{avg}}(\mathcal{A}) > 0$, which is strictly weaker than positive reach: it allows isolated cusps of measure zero.

This is the precise formulation of the "BV regularity gap" and connects to:

- *Moreau–Yosida regularisation* in convex analysis (the time-averaged cost is the Moreau envelope),
- *Viscous selection* in gradient systems (the time average acts as artificial viscosity),
- *Thermodynamic selection* in the Dark Energy companion paper (the IR averaging of $\Phi[\rho]$).

12 Open Directions

The TLL+LDI framework opens several concrete research directions grouped by theme.

12.1 Pilot verification systems for TLL and LDI

Remark 12.1 (Pilot verification systems for TLL and LDI). Three candidate systems for pilot verification of the TLL and LDI conditions:

1. *Reaction-diffusion (2D):* Smooth nonlinearity, well-understood attractors. TLL and LDI are expected to hold due to the parabolic smoothing.
2. *Hyperviscous NSE ($(-\Delta)^\beta$, $\beta \ge 5/4$):* Inertial manifolds exist (Mallet-Paret–Sell, 1988), guaranteeing TLL. Serves as a bridge system between the tractable reaction-diffusion case and full NS ($\beta = 1$).

3. *Kuramoto–Sivashinsky (1D):* Deterministic chaos without singularities. The attractor is finite-dimensional and smooth, making TLL+LDI verifiable by direct computation.

Remark 12.2 (Ginzburg–Landau as TLL/LDI bridge). The complex Ginzburg–Landau equation $\partial_t u = (1 + i\alpha)\Delta u + u - (1 + i\beta)|u|^2 u$ possesses a finite-dimensional inertial manifold for certain parameter ranges. Complete verification of TLL+LDI on this system, followed by a lifting argument to 2D NS (via parameter continuation in the nonlinearity), would provide a concrete path to the full NS regularity problem.

12.2 Mathematical tools for TLL and LDI

Remark 12.3 (Bridge lemma: squeezing to TLL). The squeezing property (Foias–Temam) states that high modes contract faster than low modes diverge. Combined with the determining modes theorem, this implies: $d(u(t), \mathcal{A}) \leq Ce^{-\gamma_N t} d(u(0), \mathcal{A}) + C' \sup_{s \leq t} |P_N u(s) - P_N v(s)|$ for some $v \in \mathcal{A}$. Taking logarithms, $\log(1/d(t)) \geq \gamma_N t - C''$, which is the log-modulus bound underlying TLL.

Remark 12.4 (Doubling and Assouad structure). TLL can be reformulated in terms of the Assouad dimension of the attractor trajectory: if the trajectory $\{u(t) : t \in [0, T]\}$ has Assouad dimension < 1, then the "approximate tangent cone" has opening angle bounded by $C/|\log r|$, which is equivalent to TLL with rate $1/|\log r|$. This connects the regularity theory to geometric measure theory. Recent structural results on sets of positive reach [13] provide additional tools for understanding the geometry of attractor neighbourhoods.

Remark 12.5 (LDI as tail-integral equivalence). The LDI condition $\int_0^T \log(1/d(t))\,\mathrm{d}t < \infty$ is equivalent to the tail-integral condition $\int_0^R \frac{1}{s} \left(\int_{\{0<d\leq s\}} \|u'(t)\|\,\mathrm{d}t \right) \mathrm{d}s < \infty$, relating the integrability of $\log(1/d)$ to the "speed" of the trajectory near the attractor. This formulation is more amenable to energy estimates.

Remark 12.6 (Proposition 7.5 clarification). The BV $\to L^\infty$ embedding in Proposition 7.5 requires the additional condition $d(t) > 0$ a.e.: the trajectory must not touch the attractor on a set of positive measure. This is automatically satisfied for Leray–Hopf solutions (which are analytic in space and hence cannot coincide with an attractor element on an interval unless they are that element).

12.3 Stochastic extensions and advanced tools

Remark 12.7 (Stochastic stabilisation). Adding Feller-type noise $\sigma\,\mathrm{d}W$ to the Navier–Stokes equations can enforce TLL "almost surely": the noise prevents the trajectory from approaching the attractor tangentially (which would violate log-Lipschitz), and the Feller property ensures the transition semigroup maps bounded functions to continuous ones, providing the regularity needed for the selection principle.

Remark 12.8 (High-mode stability conjecture). A sufficient condition for TLL in the NS context: if the high Fourier modes $Q_N u = (I - P_N)u$ satisfy $\|Q_N u(t)\| \leq CN^{-\alpha}$ uniformly on the attractor for some $\alpha > 1/2$, then $d(t) \leq C'N^{-\alpha}$ and TLL follows from the determining modes estimate. This "high-mode stability conjecture" is weaker than inertial manifold existence (which requires $\alpha > 1$) and may be verifiable via Galerkin energy estimates. Recent sharp dimension estimates for NS-type attractors [12] provide quantitative input for such bounds.

Remark 12.9 (Variational analysis perspective). The theory of metric regularity and variational analysis (Rockafellar–Wets [16], Dontchev–Rockafellar [6]) provides a natural home for the TLL and LDI conditions: TLL is related to the Aubin property of the attractor projection, and LDI is a calmness condition on the distance function. The full toolkit of variational analysis—including Ekeland's variational principle, the Mordukhovich criterion, and the Attouch–Théra duality—is applicable and underexplored in this context.

12.4 Further NS-specific research directions

Remark 12.10 (Further research directions).

1. *Selection regularity via trajectory attractors:* Formulate the minimal metric derivative selection in the Chepyzhov–Vishik trajectory attractor framework, where non-uniqueness is handled by considering entire trajectories rather than initial data.

2. *Quantitative tracking lemmas:* Establish measure estimates $|\{t : d(t) \leq \varepsilon\}| \leq C\varepsilon^{\alpha}$ via energy/absorbing-ball arguments and "no sticky contact" conditions.

3. *Ekeland variational principle:* Use Ekeland's principle to construct "almost minimisers" with controlled variation, bypassing the need for positive reach of $\mathcal{A}$.

4. *Set-valued BV selection:* Determine which dynamical regularity conditions on the flow force bounded Hausdorff variation of the minimiser set $\{u \in \mathcal{A} : d(u, u(t)) = d_{\mathcal{A}}(u(t))\}$.

5. *Approximate prox-regularity:* Establish a modified reach after removing a meager set (Lebesgue-null): the attractor may fail to have positive reach on a negligible set but retain it generically. Results of Lytchak [13] on the structure of sets of positive reach suggest that such a decomposition may be feasible.

6. *Weak selection functions:* Replace strong Lipschitz projections with selection functions of bounded Hausdorff variation, which exist under weaker regularity assumptions.

7. *Geometric Euler system:* Construct a coherent hierarchy of finite Galerkin approximations $\mathcal{A}_N \to \mathcal{A}$ as a "reach surrogate": even if $\mathrm{reach}(\mathcal{A}) = 0$, the Galerkin approximations $\mathcal{A}_N$ have positive reach and converge in Hausdorff distance.

12.5 Cross-paper connections

Remark 12.11 (DFC from turbulence as topological funnel for TLL). The Detailed Flux Constraint (DFC) from the turbulence companion paper [11] implies that energy flows "downhill" through the inertial range. Translated to the attractor geometry, this means trajectories approach $\mathcal{A}$ through a topological funnel: the DFC constrains the approach direction, enforcing a log-Lipschitz rate. This provides a physical mechanism for TLL that is independent of the abstract functional-analytic arguments.

AI Disclosure

In the preparation of this manuscript, AI-assisted tools (Claude, Anthropic) were used in a supporting capacity, particularly for literature research, text structuring, and formulation

assistance. The conceptual design, scientific argumentation, and responsibility for the entire content rest solely with the author.

References

[1] L. Ambrosio, N. Gigli, and G. Savaré, *Gradient Flows in Metric Spaces and in the Space of Probability Measures*, 2nd ed., Lectures in Mathematics ETH Zürich, Birkhäuser, Basel, 2008.

[2] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Mathematical Monographs, Clarendon Press, Oxford, 2000.

[3] A. V. Babin and M. I. Vishik, *Attractors of Evolution Equations*, Studies in Mathematics and its Applications, vol. 25, North-Holland, Amsterdam, 1992.

[4] V. Barbu and M. Cannone, *Locally Lipschitz composition operators in spaces of functions of bounded variation*, arXiv preprint arXiv:1602.04490, 2016.

[5] V. V. Chistyakov, *Selections of bounded variation*, J. Appl. Anal. **10** (2004), no. 1, 1–82.

[6] A. L. Dontchev and R. T. Rockafellar, *Implicit Functions and Solution Mappings: A View from Variational Analysis*, Springer Monographs in Mathematics, Springer, New York, 2009.

[7] P. Constantin, C. Foias, and R. Temam, *Attractors representing turbulent flows*, Mem. Amer. Math. Soc. **53** (1985), no. 314.

[8] A. Eden, C. Foias, B. Nicolaenko, and R. Temam, *Exponential Attractors for Dissipative Evolution Equations*, Research in Applied Mathematics, vol. 37, Wiley–Masson, Paris, 1994.

[9] H. Federer, *Curvature measures*, Trans. Amer. Math. Soc. **93** (1959), 418–491.

[10] C. Foias, O. Manley, R. Rosa, and R. Temam, *Navier–Stokes Equations and Turbulence*, Encyclopedia of Mathematics and its Applications, vol. 83, Cambridge University Press, 2001.

[11] L. Geiger, *A Thermodynamic Obstruction to Finite-Time Blow-Up in the 3D Navier–Stokes Equations (Conditional Framework)*, Zenodo, 2026. DOI: to be assigned.

[12] A. Ilyin, V. Kalantarov, and S. Zelik, *Attractors and their dimensions for the 3D Fractional Navier–Stokes–Voigt Equations*, arXiv preprint arXiv:2511.04783, 2025.

[13] A. Lytchak, *A note on subsets of positive reach*, Math. Nachr. **297** (2024), no. 3, 932–942.

[14] J. Mallet-Paret and G. R. Sell, *Inertial manifolds for reaction diffusion equations in higher space dimensions*, J. Amer. Math. Soc. **1** (1988), no. 4, 805–866.

[15] R. Mané, *On the dimension of the compact invariant sets of certain non-linear maps*, Lecture Notes in Mathematics, vol. 898, Springer, Berlin, 1981, pp. 230–242.

[16] R. T. Rockafellar and R. J-B Wets, *Variational Analysis*, Grundlehren der mathematischen Wissenschaften, vol. 317, Springer, Berlin, 1998.

[17] R. A. Poliquin, R. T. Rockafellar, and L. Thibault, *Local differentiability of distance functions*, Trans. Amer. Math. Soc. **352** (2000), no. 11, 5231–5249.

[18] S. Saks, *Theory of the Integral*, 2nd revised ed., Monografie Matematyczne, vol. 7, G. E. Stechert & Co., New York, 1937.

[19] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, 2nd ed., Applied Mathematical Sciences, vol. 68, Springer, New York, 1997.