

Functional Stability Theory III: Biological Stability and Nash Frustration

Lukas Geiger

Independent Researcher, Bernau, Germany

ORCID: <https://orcid.org/0009-0005-7296-1534>

June 2026

Abstract

This paper develops a conjectural framework for biological organization spanning molecular to ecosystem scales, organized around thermodynamic game theory and the Maximum Entropy Production Principle (MEPP). It examines whether biological stability problems can be modeled as dissipative structures whose attractors admit Nash-style fixed-point descriptions once players, strategies, and payoffs are specified at the relevant scale. The framework brings England’s dissipative adaptation theory, Friston’s Free Energy Principle, Kauffman’s Boolean network dynamics, and classical evolutionary game theory into a working hierarchy: at each level—from protein folding through gene regulation, cellular coalitions, to ecosystem dynamics—game-theoretic stability offers a common language for biological organization. The major evolutionary transitions are treated as possible coarse-graining steps that may preserve this structure across scales. As a concrete application, we introduce *Nash frustration* for protein structures and provide a preliminary single-protein consistency test against NMR chemical shift perturbation data (Spearman rank correlation $\rho_S = 0.44$, $p = 0.033$, $n = 24$). The normalized residue ranking is independent of η in the checked local scan window, while absolute instability scales, including $\rho(J)$, remain calibration targets. The framework proposes testable hypotheses for protein folding landscapes, gene regulatory network attractors, and ecosystem resilience.

Keywords: evolutionary game theory, ESS, protein folding, gene regulation, MEPP, dissipative adaptation, Free Energy Principle, Boolean networks, major transitions, coarse-graining

1 Introduction

The emergence and stabilization of biological complexity presents modern science with the challenge of bridging the fundamental laws of physics and the dynamic strategies of life. The Functional Stability Theory (FST-III) framework treats biological systems not merely as products of random mutations but as candidate dissipative structures operating within thermodynamic constraints [Kleidon, 2010].

This paper investigates the proposed connection between game theory and the Maximum Entropy Production Principle (MEPP), exploring whether biological stability can be understood through the lens of game-theoretic attractors within thermodynamic constraints. The scope is primarily biological (Sections 3–9); cosmological extensions are briefly noted in Section 12.11 but are not part of the core argument.

Classical evolutionary game theory, pioneered by Maynard Smith and Price [Maynard Smith & Price, 1973, Maynard Smith, 1982], revolutionized our understanding of biological behavior. Yet the standard treatment is largely confined to behavioral ecology. This paper explores a broader proposal: game-theoretic descriptions may be formulated at multiple levels of biological organization, and these descriptions connect biology to physics through an analogous stability logic developed in FST-I [Geiger, 2026a] and FST-II [Geiger, 2026b].

2 Structural Alignment: FST-III as a Pattern A Instance

The biological stability framework developed here (ESS/FEP) is proposed as a **Pattern-A-style structural analogy** within the FST ecosystem (see Geiger [2026d]), not as a formal derivation from the mathematical Pattern A programme. The recent **gauge-theoretic reformulation of the Riemann Hypothesis** (see Geiger [2026c]) provides a motivating structural template for this architecture.

Within this hierarchical stability framework, biological systems are understood as follows:

- a) **Analytical Rank-Finiteness (Null-Tail):** Biological complexity may be constrained against the combinatorial explosion of the “dead” state because, at any given environmental boundary condition, only a limited set of genotype configurations may satisfy simultaneous ESS and FEP stability. This finiteness is conjectured, not proven; it is motivated by the observation that the intersection of ESS conditions (stability against invasion) and thermodynamic viability (positive dissipation) imposes sufficiently many independent constraints to reduce the solution set to a discrete collection. The unorganized “noise” is filtered by the entropic constraint, supporting robust persistence.
- b) **Finite Negativity as Gauge-Signal (Structural Analogy):** Genetic conflict, cancer, and competition play a role analogous to finite negativity in the arithmetical kernel. They are not treated here as random defects, but as signals of incomplete organizational coordination that can be reduced or reorganized through new levels of cooperation (major transitions). This analogy is structural, not formal: the mathematical mechanisms differ, but the logical pattern (destabilization signals driving reorganization) is shared.

- c) **Constrained Functional Positivity (Biological Attractors):** The stable attractors of a gene network or ecosystem may be represented as “gauge-stable” states where the free-energy functional reaches a constrained minimum. In analogy with rank-reduction mechanisms in the RH programme, phenotypic stability is interpreted here as the result of a finite set of epigenetic constraints supporting thermodynamic viability.

This alignment frames FST-III as a structural analogy—not a formal derivation—to “Functional Positivity under Constraint.”

Definition 1 (Resolvent-stable fixed point). *A fixed point of a dynamical system is called resolvent-stable if second-order perturbation analysis (Hessian eigenvalues or resolvent expansion coefficients) yields strictly positive values in all non-gauge directions—i.e., all perturbation modes orthogonal to symmetry-generated degeneracies are stabilized. In biological systems, “non-gauge directions” correspond to perturbation modes that change observable phenotypic properties (e.g., protein conformation, gene expression pattern), while “gauge directions” correspond to neutral degrees of freedom that leave the phenotype invariant (e.g., synonymous codon substitutions, neutral conformational fluctuations within a basin). This is a structural analogy—motivated by, but not formally derived from—the spectral gap property established in the FST-RH framework [Geiger, 2026e]; its application to biological systems requires domain-specific verification of which directions are genuinely “gauge.”*

3 Thermodynamic Foundations: MEPP and Dissipative Adaptation

3.1 Life as Non-Equilibrium Phenomenon

The classification of biological processes in thermodynamics begins with the recognition that life is a phenomenon far from thermodynamic equilibrium [Kleidon, 2010]. While closed systems inevitably tend toward the state of maximum entropy and heat death, biological systems maintain themselves through continuous export of entropy to their environment—a process that Erwin Schrödinger described in 1944 as the acquisition of “negentropy” [Schrödinger, 1944, Kleidon, 2010].

3.2 Three Thermodynamic Hypotheses

Within the FST-III framework, three central thermodynamic hypotheses describe the behavior of living matter [Kleidon, 2010]:

1. **Thermodynamic Fingerprint of Life:** Living communities significantly increase the rate of entropy production compared to abiotic counterparts under identical boundary conditions.
2. **State Selection Principle (Stability Interpretation):** When a system can reach multiple stable states, the resolvent-stable configuration—where second-order destabilization channels are suppressed—tends to exhibit high (though not necessarily maximal) entropy production rates.

3. **Gradient Response Principle:** When the external thermodynamic gradient increases, the system transitions to a new stable state characterized by even higher entropy production.

Table 1: Thermodynamic principles and their biological relevance. MEPP applies to strongly driven systems; Prigogine’s minimum entropy production holds only near equilibrium.

Principle	Description	Biological Relevance
MEPP	Stability filter favoring high dissipation	Metabolic efficiency selection
State Selection	Stability correlates with high $\dot{S}$	Complex food web stabilization
Gradient Response	Adaptation to increasing energy input	Photosynthetic efficiency increase
Prigogine (LMEP)	Minimal entropy near equilibrium	Only for linear, weakly driven systems

Remark 1 (MEPP as Stability Filter, not Selection Law). Following the resolvent analysis from the Riemann Hypothesis program [Geiger, 2026e], the three thermodynamic hypotheses above are read here as *stability filters* rather than selection laws. The key distinctions are:

- (i) **Not a causal driver:** MEPP does not causally drive biological systems toward states of maximum entropy production.
- (ii) **Degeneracy at leading order:** Many biological configurations are viable at first order; the leading-order dynamics are degenerate.
- (iii) **Second-order filter:** MEPP acts as a second-order filter that eliminates resolvent-unstable configurations—those susceptible to perturbation-induced collapse via cross-coupling between metabolic, structural, and replicative degrees of freedom.
- (iv) **Candidate fixed point:** The state with “highest entropy production” is more cautiously read as a candidate resolvent-stable fixed point within the degenerate manifold of thermodynamically viable configurations.

This refinement addresses the criticism that MEPP is used as a selection law: it is a stability criterion, not a causal driver. For the full development in the prebiotic context, including the formal England inequality and the MEPP controversy, see FST-II [Geiger, 2026b]; here we apply this principle specifically to biological systems.

3.3 Dissipative Adaptation: The Physics of Self-Organization

A crucial link between pure thermodynamics and biological structure is provided by Jeremy England’s theory of dissipative adaptation [England, 2013, 2015]. For the full formal derivation of England’s dissipation bound from the Crooks fluctuation theorem, the game-theoretic reading of the inequality, and the causal-direction caveat, see FST-II [Geiger, 2026b]; here we summarize only the aspects relevant to the biological context.

England argues, based on non-equilibrium statistics, that matter exposed to a strong external energy source tends to organize itself into configurations that are particularly good at absorbing energy and dissipating it as heat [England, 2013].

The mechanism of dissipative adaptation relies on the irreversibility of state transitions. When a system absorbs energy from an external source, it can overcome energetic activation barriers that would be impassable through purely thermal fluctuations. Once this energy is dissipated, the system lacks the energy to jump back to the initial state. Over time, structures accumulate that exhibit high resonance with the external energy source [England, 2015].

This process offers one physical account of the spontaneous emergence of self-replicating structures: replication is a direct route by which a system can increase its capacity for energy absorption and dissipation [England, 2015]. In this context, “fitness” appears not as a purely biological concept but as a physical property of structural persistence under energetic drive [England, 2013].

4 Game-Theoretic Stability: Molecular ESS

Once physical structures cross the threshold to biological reproduction, their interactions can be represented with game-theoretic rules.

4.1 Nash Equilibrium and ESS

The central concept is the Nash equilibrium, defined by John Nash in 1950, describing states where no actor can improve their payoff through unilateral change of their strategy [Holt & Roth, 2004, Nash, 1950]. In biology, John Maynard Smith and George Price developed this into the Evolutionarily Stable Strategy (ESS) [Maynard Smith & Price, 1973, Maynard Smith, 1982].

Definition 2 (Evolutionarily Stable Strategy). *An ESS is a strategy that, when adopted by most members of a population, cannot be invaded by a rare mutant strategy. Every ESS is a Nash equilibrium, but not every Nash equilibrium is an ESS, since ESS requires additional stability conditions against invasions [Maynard Smith, 1982].*

In FST-III, this principle is transferred to the molecular level. Proteins, enzymes, and regulatory RNAs are considered “players” whose “strategies” consist of their structural configurations and catalytic activities [Bohl et al., 2014].

Remark 2 (ESS as Resolvent-Stable Fixed Point). A proposed lesson from the resolvent analysis of the Riemann Hypothesis [Geiger, 2026e] refines the interpretation of ESS. An ESS need not be read as a global fitness optimum; in this analogy, it is a resolvent-stable fixed point. At leading order, the fitness landscape may be degenerate—many phenotypic configurations yield comparable payoffs. Selection then acts at second order through resolvent-type couplings between the degenerate modes: invasion dynamics, frequency-dependent interactions, and spatial coupling collectively split the degeneracy. The ESS is the candidate configuration where second-order destabilization channels are jointly suppressed. Nash frustration (defined below in Section 4) is therefore not simply a defect but a possible structural feature of this resolvent architecture: it reflects residual second-order coupling that may stabilize the configuration while preventing crystallization into a rigid, fragile optimum.

Table 2: Mapping of game-theoretic concepts to physical and biological counterparts.

Game Concept	Physical Correspondence	Biological Example
Nash Equilibrium	Stable steady state	Phenotype coexistence
ESS	Attractor with local stability	Allele fixation
Payoff Matrix	Energy/fitness landscape	Enzyme-substrate affinity
Mixed Strategy	Population heterogeneity	Division of labor in colonies
Nash Frustration	Residual instability at stability boundary	Functional flexibility in proteins

4.2 Protein Folding as an Optimization Game

A useful example is protein folding, which can be modeled heuristically as a game against the thermodynamic landscape [Bryngelson & Wolynes, 1989]. The amino acid sequence tends toward low-free-energy conformations; in the present model, the native conformation is represented as a Nash-style equilibrium in an n -person game of molecular forces:

- **Players:** Individual amino acid residues
- **Strategies:** Backbone dihedral angles (ϕ, ψ)
- **Payoff:** $-\Delta G$ (negative free energy change)
- **Nash Equilibrium:** Native conformation (Anfinsen’s dogma [Anfinsen, 1973])

Misfolded proteins (prions) represent alternative local equilibria—metastable configurations where no small unilateral deviation improves the payoff, even though the global payoff is suboptimal. The prion case is particularly instructive: the misfolded conformation is kinetically trapped and propagates through template-directed conversion rather than thermodynamic selection, illustrating that Nash-like stability can arise from kinetic barriers as well as from energetic optimality. Note that prion propagation constitutes a well-known exception to Anfinsen’s thermodynamic hypothesis: the misfolded state is not the global free-energy minimum but is stabilised by intermolecular templating and aggregation kinetics.

4.3 Nash Frustration vs. Energetic Frustration

Notation: $\rho(J)$ denotes the spectral radius of the Jacobian, ρ_S the Spearman correlation coefficient, and ρ_i the normalized residue-level frustration score. These distinct uses of ρ follow their respective community conventions and are disambiguated by subscript throughout.

The concept of local frustration in protein structures has been extensively studied through the Protein Frustratometer [Ferreiro et al., 2007, 2014], which identifies residues where the native amino acid identity is suboptimal given the local structural environment. This energetic frustration correlates with functional sites, binding interfaces, and allosteric regions.

The FST-III framework introduces a complementary concept: **Nash frustration**. While energetic frustration asks “which residues have suboptimal local energy?”, Nash frustration asks “which residues prevent game-theoretic coordination?”

Formally, *residue-level Nash frustration* is computed from the best-response Jacobian $J = I - \eta H$, where H is the $2n \times 2n$ Hessian of the potential (with n residues, each contributing two dihedral angles ϕ_i, ψ_i) and $\eta > 0$ is a learning-rate parameter. The frustration score for residue i is:

$$\text{frustration}(i) = \sum_{k: |\lambda_k| > 1} (|\lambda_k| - 1) (v_{k, \phi_i}^2 + v_{k, \psi_i}^2) \quad (1)$$

where λ_k and v_k are eigenvalues and eigenvectors of J , respectively. The *global Nash frustration* of the entire protein is summarized by the spectral radius $\rho(J) = \max_k |\lambda_k|$; when $\rho(J) > 1$, at least one mode is unstable under best-response dynamics. These two measures—residue-level scores and the global spectral radius—are complementary: $\rho(J)$ quantifies the overall departure from Nash stability, while the residue-level score localizes the instability.

Coordinate-metric caveat. The projection above uses the Euclidean norm of eigenvector components in the local dihedral chart (ϕ, ψ) . It is therefore a chart-level diagnostic, not an intrinsic coordinate-invariant biomolecular observable. An intrinsic or mass-weighted version should replace the squared component sum by a metric-weighted projection using the appropriate torsional/manifold metric; the present single-protein benchmark uses a fixed chart and should be interpreted under that convention.

Empirical validation and η -independence. We validated the Nash frustration metric against experimental NMR chemical shift perturbation (CSP) data derived from BMRB entry 4428 Vardar et al. [1999]. This BMRB entry is the HP67 villin-headpiece NMR data set (primary PDB 1QQV); for the HP35 structural benchmark (PDB 1YRF), we used the documented residue-level mapping from the C-terminal HP67 subdomain to HP35, excluding mutated or missing positions. The CSP per residue is computed as $\text{CSP}_i = \sqrt{\Delta\delta_{\text{H}}^2 + (\Delta\delta_{\text{N}}/5)^2}$, where $\Delta\delta$ denotes the secondary chemical shift relative to random-coil reference values Wishart et al. [1995]; the factor 1/5 is the standard weighting for ^{15}N chemical shifts Williamson [2013].

The Spearman rank correlation between per-residue Nash frustration and CSP is $\rho_S = 0.44$ ($p = 0.033$, $n = 24$ residues), providing initial evidence that frustration scores capture the rank ordering of NMR-observed structural perturbation. We note that the statistical power of this test is moderate ($\approx 60\%$ at $\alpha = 0.05$ for the observed effect size), and replication on additional proteins is needed to confirm the result. The five most frustrated residues (M11, T12, S1, L0, L19) exhibit CSP values of 1.26, 0.68, 2.13, 1.44, and 1.10 ppm, consistent with frustration hotspots corresponding to experimentally detectable structural strain. One outlier (K23, CSP = 2.67 ppm but low frustration) is attributable to the well-known Trp ring-current shift at this position rather than structural instability.

Crucially, the frustration–CSP correlation is *independent* of η in the verified local scan window: since $\lambda_k = 1 - \eta h_k$, unstable modes generated by negative Hessian directions ($h_k < 0$) have instability excess $\eta(-h_k)$, and this common factor cancels under max-normalization to $[0, 1]$. This reasoning applies only while the unstable active set is unchanged and no positive-curvature overshoot modes ($h_k > 2/\eta$) cross $|\lambda_k| > 1$. A numerical scan over $\eta \in \{0.001, 0.005, 0.01, 0.015, 0.02, 0.05, 0.1\}$ confirms this local expectation for the present benchmark (variation $< 10^{-6}$). Therefore, η is not a tuning parameter

for the normalized residue ranking within this checked window; it remains relevant for absolute score scales, spectral-radius thresholds, convergence diagnostics, and any regime in which the unstable active set changes. This result, checked against two independent random-coil reference sets (Wishart 1995, $r = 0.18$; Kjaergaard–Poulsen 2011 Kjaergaard & Poulsen [2011], $r = 0.20$; $|\Delta r| < 0.02$), supports a preliminary single-protein consistency diagnostic rather than an independently calibrated predictor.




Figure 1: Scatter plot of per-residue Nash frustration scores versus NMR chemical shift perturbation (CSP) for the HP35/1YRF benchmark, using BMRB-4428-derived HP67 chemical shifts mapped to the C-terminal HP35 subdomain. Each point represents one residue; the Spearman rank correlation is $\rho_S = 0.44$ ($p = 0.033$). Colour encodes the learning-rate parameter η , demonstrating that the frustration–CSP relationship is invariant across the checked local η scan (all η -values collapse onto the same curve). The outlier K23 (high CSP, low frustration) is attributable to a Trp ring-current shift rather than structural instability.

Remark 3 (Computational Protocol). The Nash-Frustration analysis uses a *custom game-theoretic potential*, not a molecular-mechanics force field (such as AMBER or CHARMM). Each residue i is treated as a player with strategy $\theta_i = (\phi_i, \psi_i)$. The potential $F = F_\phi + F_\psi$ consists of single-player terms—amino-acid-specific Ramachandran preferences $\mu_\phi^{(a_i)}, \mu_\psi^{(a_i)}$ with stiffness constants $k^{(a_i)}$ —and pairwise coupling terms weighted by a radial-basis-function kernel

$$w(r_{ij}) = \sum_{b=1}^B c_b \exp\left(-\frac{(r_{ij} - \mu_b)^2}{2\sigma^2}\right), \quad B = 12, \sigma = 1.0 \text{ \AA}.$$

Contacts are defined by a C_α – C_α distance cutoff of $r_{\text{cut}} = 8.0 \text{ \AA}$. Parameters (Ramachandran preferences, stiffness constants, RBF coefficients) are reverse-engineered from the native PDB structure via regularized least-squares fitting (regularization weight $\lambda_{\text{reg}} = 0.01$; multi-start gradient descent with 64 random initializations, 6 000 steps each).

Circularity caveat. Because the potential is fitted to the native structure, convergence of best-response dynamics to that structure is expected by construction and does not constitute an independent validation. The non-trivial content of the CSP correlation ($\rho_S = 0.44$) is that the *rank ordering* of residue-level frustration—which depends on the eigenvector structure of the fitted Hessian, not merely on the fit quality—correlates

with an independent experimental observable (NMR chemical shifts) that was not used in the fitting procedure. Nevertheless, the fitted potential shapes the Hessian and thereby influences which residues appear frustrated. A stringent test of the Nash frustration concept would require either (a) a physics-based potential not fitted to the native structure (e.g., an all-atom force field), or (b) cross-validation in which parameters fitted to one protein predict frustration patterns in structurally unrelated proteins. Until such tests are performed, the present result should be interpreted as a *consistency check of the self-parametrised model*, not as an independent prediction.

The best-response Jacobian $J = I - \eta H$ uses a learning rate η ; the software default is $\eta = 0.015$, while the HP35 pilot study reported below uses $\eta = 0.01$ as an absolute-scale convention near $\rho(J) \approx 1$. The Hessian H is computed by central finite differences ($\epsilon = 10^{-4}$). Best-response dynamics employ 300 sweeps from 16 random starting configurations. The frustration score ρ_i projects the unstable eigenmodes of J (those with $|\lambda_k| > 1$) onto residue i and normalizes to $[0, 1]$.

As demonstrated above, η is not optimized to maximize the normalized CSP correlation: the rank-level frustration pattern is independent of η in the range $[0.001, 0.1]$. Further work should calibrate the absolute score scale and $\rho(J)$ -based thresholds against independent observables such as hydrogen–deuterium exchange (HDX) protection factors (McKnight et al. 1996 McKnight et al. [1996]). Source code: `protein_fold_nash_pdb.py` (Python 3, BioPython, SciPy, NumPy). Extended three-protein analysis (LaTeX table generation, convergence diagnostics): `run_extended_analysis.py`.

Table 3 separates the empirical and computational evidence blocks from the next validation gates. This distinction is important because the current dataset supports a local, self-parametrised consistency check, whereas predictive use requires independent controls and absolute-scale calibration.

Computational experiments on three structurally distinct proteins explore whether Nash frustration localizes biologically meaningful residues and whether best-response dynamics converge beyond the training structure. Table 5 summarizes the results.

Training protein (HP35). Nash frustration concentrates at loop regions (particularly Met11, Pro20 with scores 1.0 and 0.49), while helix cores (Ser14, Asn18, Trp22) exhibit near-zero frustration—consistent with their role as “structural anchors” in the folding game. The spectral radius $\rho(J) = 1.013$ with 80% positive Hessian eigenvalues and mean frustration 0.082 characterize a near-Nash structure. This pattern differs from energetic frustration, which typically peaks at surface-exposed functional residues.

Cross-protein convergence test. The model parameters learned from HP35 were used to test whether best-response dynamics converge for Protein G (1PGA, 56 residues) and the p53 tumor suppressor DNA-binding domain structure used here (2XWR, 199 modeled residues). Both converged to well-defined fixed points (Table 5). The p53 domain—roughly six times larger than the training protein and belonging to a different fold class (immunoglobulin-like β -sandwich vs. α -helical)—achieved a ϕ_{rms} of 1.25 rad, comparable to Protein G (1.30 rad). These ϕ_{rms} values correspond to mean per-residue deviations of $\sim 72^\circ$, indicating that the converged structures are *not* close to the native folds. The key observation is therefore limited to *convergence* of the best-response dynamics across proteins of different size and fold class, not structural prediction accuracy. The potential-game formulation captures convergence properties of the fitted dynamics, but structural prediction would require a richer potential including side-chain interactions, solvent effects, and electrostatics.

Table 3: Validation gate ledger for the Nash-frustration evidence. The table separates current observables from the minimum next gate needed before stronger predictive claims.

Evidence block	Current observable	Present status	Next gate
HP35/CSP rank check	BMRB-4428-derived CSP mapped to HP35; $\rho_S = 0.44$, $p = 0.033$, $n = 24$	Single-protein consistency signal; not an externally calibrated predictor	Replicate on additional proteins with preregistered residue mapping and fixed scale convention
η scan	Seven values in $\{0.001, \dots, 0.1\}$; normalized ranking variation $< 10^{-6}$	Rank-level local invariance under unchanged unstable active set	Calibrate absolute scores and $\rho(J)$ thresholds out of sample; track active-set changes and positive-curvature overshoot
Cross-protein convergence	HP35-trained parameters applied to 1PGA and 2XWR; all runs converge, but $\phi_{\text{rms}} \approx 1.25\text{--}1.30$ rad for test proteins	Convergence diagnostic only; no structural prediction accuracy claim	Fix 2XWR provenance and evaluate richer potentials with side-chain, solvent, and electrostatic terms
Functional-residue benchmark	AUC 0.616 for HP35 and 0.688 for Protein G	Weak-to-moderate signal above chance; below standalone diagnostic strength	Direct head-to-head comparison with Ferreiro frustration, B-factors/NMA, pLDDT, and matched sequence/structure controls
Prospective molecular tests	HDX protection factors, ϕ -values, allosteric/interface annotations, helix melting order	Falsifiable agenda specified; not yet executed	Run multi-protein tests with preregistered thresholds and report negative controls alongside positive correlations

Table 4: Comparison of energetic frustration (Ferreiro et al.) and Nash frustration (this work). The two measures probe complementary aspects of protein stability.

Concept	Energetic Frustration	Nash Frustration
Question	Is local energy optimal?	Is unilateral deviation stable?
Method	Contact energy statistics	Best-response Jacobian eigenanalysis
Interpretation	Evolutionary constraint	Game-theoretic coordination
High values indicate	Functional sites, interfaces	Folding intermediates, kinetic traps

Table 5: Nash equilibrium convergence analysis of protein folds. Parameters learned from HP35 (Villin headpiece, 1YRF) and applied to test convergence of best-response dynamics for Protein G (1PGA, 56 residues) and the p53 DNA-binding domain structure used here (2XWR, 199 modeled residues). Best-response dynamics converge across all three proteins despite a several-fold range in protein size, though the large ϕ_{rms} values indicate that converged structures do not closely resemble native folds.

Protein	Residues	ϕ_{rms} [rad]	ψ_{rms} [rad]	Converged	F_{best}
HP35 (1YRF, train)	35	1.054	1.661	Yes (267 sw)	−4.61
Protein G (1PGA)	56	1.300	2.002	Yes (282 sw)	−10.61
p53 DBD (2XWR)	199	1.247	2.114	Yes	−39.43

Computational observation (Nash Stability Boundary). Native protein folds are not exact Nash equilibria under local best-response dynamics, but lie close to the stability boundary ($\rho(J) \approx 1$). The spectral radius $\rho(J) = 1.013$ for HP35 indicates that the native structure is nearly self-stabilizing, with 14 out of 70 eigenvalues of J exceeding unity in magnitude, and these unstable modes are localized at functional hinge regions rather than reflecting global coordination failure.

This observation suggests that Nash stability is *stricter* than energy minimization. A structure can be a local energy minimum while still permitting unilateral improvements for individual residues. The localized Nash frustration at Met11 is consistent with a functional freedom degree—a “hinge” that enables conformational flexibility while the helix cores form stable coalitions. This is what one expects for functional proteins under the present interpretation: they balance stability against flexibility, rather than crystallizing into rigid Nash optima. A systematic calibration of the absolute score scale against experimental data—for instance, by requiring that Nash-unstable residues coincide with known allosteric sites or folding nuclei identified by hydrogen-deuterium exchange experiments—would be needed before this proof-of-concept could support predictive use. Until such calibration is performed, the numerical values reported here ($\rho(J) = 1.013$, frustration peaks at Met11 and Pro20) should be interpreted as parameter-dependent illustrations rather than robust quantitative predictions. In terms of the FEP–MEPP compatibility (Conjecture 1), Nash frustration $\rho(J)$ can be read as a local tension between internal free-energy minimisation (mEPP regime) and external entropy-production maximisation (MEPP regime): high frustration marks residues where the fitted folded state locally departs from the MEPP-optimal dissipative configuration in favour of structural stability. This is a stability interpretation of the model, not a direct organism-level choice.

Planned quantitative comparison. A systematic residue-by-residue comparison of Nash frustration scores with the energetic frustration indices computed by the Protein Frustratometer 2.0 Parra et al. [2016] constitutes a key future validation step. The central question is whether Nash-frustrated residues (high $|\lambda_k| - 1$ contributions) overlap with, complement, or are orthogonal to energetically frustrated residues. Preliminary expectations suggest partial complementarity: energetic frustration identifies evolutionary constraints and binding sites, while Nash frustration should highlight kinetic traps and folding intermediates (Table 4). A dataset of proteins with known allosteric sites and experimentally determined ϕ -values would provide the empirical basis for this comparison. Rank-level map comparisons do not require an optimized η ; absolute thresholds and global

$\rho(J)$ comparisons do require an explicit scale convention or out-of-sample calibration (see Remark 3).

4.4 Benchmark: Nash Frustration vs. Functional Residues

To assess whether Nash frustration carries information about biological function beyond random expectation, we performed a benchmark analysis comparing residue-level Nash frustration scores against experimentally annotated functional residues for HP35 (1YRF) and Protein G (1PGA). For each protein, residues were classified as *functional* (catalytic sites, binding interfaces, allosteric regions from UniProt annotations) or *passive* (all others), and the Nash frustration score was evaluated as a binary classifier using ROC and precision-recall analysis. Table 6 summarizes the results.

Table 6: Benchmark of Nash frustration as a predictor of functional residues. AUC: area under the ROC curve; AUPRC: area under the precision-recall curve. Mean frustration values are reported as mean $\pm$ standard deviation. Script: `frustration_benchmark.py`.

Protein	AUC	AUPRC	Mean frust. (functional)	Mean frust. (passive)
HP35 (1YRF)	0.616	0.550	0.160 ± 0.269	0.031 ± 0.082
Protein G (1PGA)	0.688	0.618	0.229 ± 0.247	0.096 ± 0.101

The AUC values of 0.62 (HP35) and 0.69 (Protein G) indicate weak to moderate discriminatory power—below the AUC = 0.7 threshold commonly considered “acceptable” in diagnostic applications [Ferreiro et al., 2014]. Nash frustration carries information about functional residues beyond chance level (AUC > 0.5), but is not diagnostically strong. For context, the Protein Frustratometer typically achieves AUC > 0.75 for functional-site identification; a direct head-to-head comparison on the same proteins is needed to quantify whether Nash frustration provides complementary or merely weaker information. Functional residues exhibit significantly higher mean Nash frustration than passive residues in both proteins (0.160 vs. 0.031 for HP35; 0.229 vs. 0.096 for Protein G), consistent with the interpretation that game-theoretic instability preferentially localizes at biologically active sites. The modest AUC values are expected given the proof-of-concept nature of the analysis: absolute score thresholds remain uncalibrated, and the game-theoretic potential omits side-chain interactions, solvent effects, and electrostatics. These results are consistent with the interpretation that Nash frustration captures a complementary signal to energetic frustration—one that reflects coordination failure rather than energetic suboptimality—but requires further calibration before it can serve as a standalone predictor.

4.5 Gene Regulation as a Signaling Game

Gene regulatory networks (GRNs) control gene expression patterns and can be modeled as signaling games [Bohl et al., 2014]:

- **Players:** Transcription factors and DNA binding sites
- **Strategies:** Binding/non-binding (with continuous affinity as strategy intensity)




Figure 2: Receiver operating characteristic (ROC) curves for Nash frustration as a binary classifier of functional versus passive residues. HP35 (1YRF, $AUC = 0.62$) and Protein G (1PGA, $AUC = 0.69$) both exceed the random baseline (dashed diagonal), consistent with game-theoretic instability localizing at biologically active sites. The moderate AUC values reflect the proof-of-concept status of the analysis; improved performance would require testing richer potentials with side-chain interactions and solvent effects.

- **Payoff:** Organismal fitness, mediated through gene expression levels

The mathematical connection between game theory and population dynamics is established through the replicator equation, which is formally equivalent to Lotka-Volterra models [Hofbauer & Sigmund, 1998]. More precisely, this equivalence holds for symmetric games with linear fitness functions; for frequency-dependent or nonlinear payoffs, the correspondence requires the transformation $x_i = y_i / \sum_j y_j$ and restricts to the interior of the simplex (Hofbauer & Sigmund 1998, Chapter 7). A stable fixed point in these equations corresponds to an ESS and thus to a behavioral attractor determining the long-term structure of the system.

Illustrative example: the lac operon. The *E. coli* lactose utilization system exemplifies the signaling-game logic. The lac repressor (LacI) and CAP activator compete for regulatory control of the *lac* operon. In the absence of lactose, LacI binding constitutes a Nash equilibrium: no unilateral change of binding state by any regulatory factor improves organismal fitness. When lactose is present, allolactose inactivates LacI, shifting the payoff matrix and establishing a new equilibrium with operon expression. The bistable switch between these states is precisely the kind of game-theoretic attractor that the FST-III framework identifies as an ESS at the regulatory level.

5 The Free Energy Principle and Active Inference

The Free Energy Principle (FEP), primarily developed by Karl Friston, provides a unified theory for biological self-organization that merges information theory and thermodynamics [Friston, 2009, 2010]. **Note:** The claim that FEP and MEPP are compatible or equivalent is not uncontroversial. FEP minimizes variational free energy (an information-theoretic quantity bounding surprise), while MEPP maximizes thermodynamic entropy

production rate. The connection between these two principles is an active area of debate [Friston, 2009]; the present paper treats them as complementary perspectives at different levels of analysis, but does not formally derive one from the other. For a critical assessment of FEP’s philosophical claims (e.g. Markovian Monism) versus its scientific scope as a heuristic unification principle, see Sánchez-Cañizares [2021].

5.1 Theoretical Framework

FEP postulates that every biological system that successfully defends itself against entropy (decay) must minimize its “variational free energy” [Friston, 2009]. This quantity represents a mathematical upper bound on the “surprise” (surprisal) that the system experiences through sensory inputs [Friston, 2009].

A biological agent exists within a “Markov blanket”—a statistical boundary separating internal states from external states [Friston, 2010]. To survive, the agent must ensure that its sensory states remain within physiological limits. This is achieved through two mechanisms:

1. **Perception:** The system adjusts its internal models to better predict the causes of sensory data (minimizing prediction error) [Friston, 2009]
2. **Active Inference:** The system acts to change the world or select sensory data to match its internal expectations [Friston, 2009]

Through free energy minimization, the system implicitly becomes a model of its environment. In the heuristic bridge used here, this process is linked to MEPP by scale separation: while the system internally minimizes variational free energy, it may contribute to entropy production in the environment through metabolic and agential activities [Friston, 2010].

In FST-III, behavioral attractors are those states where free energy is minimal and the agent has achieved a stable, predictable interaction with its niche [Friston, 2009].

5.2 Free-Energy Minimization as Nash Fixed Point

The connection between FEP and Nash equilibrium can be made formal under specific conditions. Let $p(o, s_1, \dots, s_N)$ be a joint generative model for N interacting agents, where each agent i controls a factored approximate posterior $q_i(s_i)$ over its latent states. The joint variational free energy is:

$$F(q; o) = \mathbb{E}_q[\ln q(s) - \ln p(o, s)], \quad q(s) = \prod_{i=1}^N q_i(s_i). \quad (2)$$

Define a game where each agent i controls q_i and all agents share the cost functional $C_i(q_i, q_{-i}) := F(\prod_j q_j; o)$. Then the best response of agent i against fixed q_{-i} is precisely the coordinate-wise variational update (mean-field VI), and a Nash equilibrium $q^* = (q_1^*, \dots, q_N^*)$ is a fixed point of distributed free-energy minimization. This constitutes a *potential game* with potential F in the sense of Monderer & Shapley [1996]: Nash equilibria coincide with coordinate-wise minima of the shared free energy.

Scope of the bridge. This formal equivalence holds for the belief-optimization component of FEP (variational inference) and relies on the *potential game* structure: all agents share the same cost functional F . In biological reality, interacting organisms typically have heterogeneous objectives (predator vs. prey, host vs. parasite), which breaks the potential-game assumption. The FEP-as-Nash bridge therefore applies strictly to intra-organismal inference (where all subsystems share the goal of organismal survival) and to symmetric multi-agent settings; it does not extend without modification to antagonistic ecological interactions. For active inference, where agents additionally select policies π_i to minimize expected free energy $G(\pi_i, \pi_{-i})$, the Nash condition becomes: no agent can unilaterally change its policy to reduce its expected free energy—a genuine multi-agent equilibrium involving interaction through state dynamics. The present paper does not develop this extension formally; the contribution is to provide the precise demarcation: FEP-as-Nash is more than metaphor when the game is explicitly defined as a variational/potential game over distributions, and merely conceptual when “Nash” is used as a synonym for “fixed point.”

5.3 Scale Separation: Resolving the FEP–MEPP Paradox

A persistent conceptual tension in theoretical biology asks how a system can *minimise* its internal free energy (FEP) while simultaneously *maximising* entropy production in the environment (MEPP). The following proposition resolves this paradox through scale separation across the Markov blanket.

Conjecture 1 (FEP–MEPP compatibility via Markov blanket). *Let $\mathcal{B}$ be a self-organising system with Markov blanket $\partial\mathcal{B}$, internal states s , and environmental states e . Define:*

- $F_{\text{int}}(q) := \mathbb{E}_q[\ln q(s) - \ln p(o, s)]$, the variational free energy of the internal model;
- $\dot{S}_{\text{ext}} := \frac{dS_{\text{env}}}{dt} = \frac{1}{T} \frac{dQ}{dt}$, the rate of entropy production in the environment.

If $\mathcal{B}$ minimises F_{int} (FEP) by maintaining an accurate internal model and selecting actions that reduce surprise, then:

- (i) *The structural integrity of $\mathcal{B}$ is maintained: $\mathcal{B}$ resists dissolution and preserves the Markov blanket (low internal entropy);*
- (ii) *The intact, far-from-equilibrium structure $\mathcal{B}$ functions as an efficient dissipative engine, channelling energy gradients from the environment through metabolic and agential activities;*
- (iii) *Among all boundary conditions consistent with $\mathcal{B}$ ’s survival, the actual dynamics tend toward configurations with high (though not necessarily maximal) rates of entropy production $\dot{S}_{\text{ext}}$ in the environment, consistent with the stability interpretation of MEPP (Hypothesis 2 above). The key mechanism linking (ii) to (iii) is competitive selection: among multiple intact dissipative structures, those that channel energy gradients more efficiently acquire more resources, replicate faster, and out-compete slower dissipators. This selection pressure—absent for passive structures like crystals—drives the correlation between structural integrity and high entropy production in the biological domain.*

In short: internal free-energy minimisation may support external entropy-production increase. *The two principles need not be in direct conflict when they are assigned to complementary sides of the Markov blanket.*

Status: This proposition is stated as a conjecture supported by qualitative arguments. A formal proof would require establishing that FEP-minimizing systems necessarily maximize environmental entropy production under the stated boundary conditions—a result that has not been demonstrated in the literature.

Remark 4 (Game-theoretic interpretation). In the language of the present paper, the FEP–MEPP compatibility corresponds to a two-level game:

- *Inner game* (within $\partial\mathcal{B}$): Agents minimise variational free energy F_{int} via the Nash fixed point of the potential game described in Section 5.
- *Outer game* (across $\partial\mathcal{B}$): The ensemble of surviving organisms competes for energy gradients. Organisms that dissipate more efficiently (higher $\dot{S}_{\text{ext}}$) outcompete slower dissipators, making MEPP a candidate macro-scale selection criterion under the stated boundary conditions.

This reframes the MEPP paradox raised by Martyushev [2010]: MEPP applies to the *environment-coupled* system, not to the internal degrees of freedom, where minimum entropy production (Prigogine) may hold locally. The Nash frustration $\rho(J)$ from Section 4 measures the residual tension between inner optimality and outer dissipative pressure.

5.4 Nash Equilibrium as Mean Field Game Fixed Point

The potential-game characterisation of the FEP–Nash bridge (Conjecture 1) can be embedded into the more general framework of *Mean Field Games* (MFG) introduced by Lasry and Lions [2007]. Consider $2N$ dihedral angles $\theta \in [-\pi, \pi]^{2N}$ parameterising the backbone of a folding protein. Each residue is treated as a representative agent whose state is governed by its local torsion angle. In the continuum (large- N) limit the coupled optimisation problem takes the form of an MFG system consisting of a Hamilton–Jacobi–Bellman (HJB) equation and a Fokker–Planck (FP) equation.

Approximation caveat. The MFG formulation assumes that each agent interacts with the *population distribution* of all other agents rather than with specific individual neighbours. For proteins, this mean-field approximation is strongest when the contact network is dense and approximately homogeneous. In practice, proteins have sparse, topology-specific contact networks (~ 5 – 10 contacts per residue out of N possible), which deviates from the mean-field limit. The MFG embedding is therefore best understood as an asymptotic idealisation valid for large N with dense coupling; for small proteins (e.g., HP35 with $N = 35$), it provides a formal scaffolding rather than a quantitatively accurate description. The “control cost” term $\frac{1}{2}\|p\|^2$ in the MFG Hamiltonian corresponds physically to the kinetic energy barrier for dihedral angle rotation, not to a literal strategic control by the residue.

The HJB equation describes the optimal control of each agent. The associated Hamiltonian reads

$$H(\theta, p, m) = \frac{1}{2}\|p\|^2 - \sum_{i < j} \int E_{ij}(\theta_i, \theta_j) m_j(\theta_j) d\theta_j, \quad (3)$$

where $p = \nabla V$ is the conjugate momentum, E_{ij} encodes pairwise residue–residue interactions, and m_j is the population density of dihedral states at site j . The Fokker–Planck equation propagates the density of agents under the optimal drift and thermal noise:

$$\partial_t m = \nabla \cdot (m \nabla V) + \sigma \Delta m, \quad \sigma = k_B T / \gamma. \quad (4)$$

Here γ is the friction coefficient of the Langevin dynamics; setting $\gamma = 1$ (natural units) recovers the simplified form $\sigma = k_B T$ used throughout. An MFG Nash equilibrium is a pair (V^*, m^*) that simultaneously satisfies both equations. In thermal equilibrium the stationary density is

$$m^*(\theta) \propto \exp(-V(\theta)/k_B T), \quad (5)$$

which is identical to the Boltzmann distribution obtained from the Langevin–Fokker–Planck description of protein folding. The existing potential-game argument (Section 5) is recovered as a special case: in the *static limit* $\partial_t = 0$, agents are assumed to occupy the minima of V (i.e., $\nabla V = 0$ at the equilibrium configuration), and the Fokker–Planck equation reduces to the detailed-balance condition $\nabla \cdot (m \nabla V) + \sigma \Delta m = 0$, which is satisfied by the Boltzmann distribution (5). Strictly, the stationary HJB equation $H(\theta, \nabla V, m) = \text{const}$ admits solutions with $\nabla V \neq 0$ away from the equilibrium point; the simplified reduction $\nabla V = 0$ holds only at the Nash fixed point itself. The resulting system is a classical potential game with shared cost functional V .

The crucial existence condition for MFG Nash equilibria is **Lasry–Lions monotonicity** Lasry and Lions [2007]: the interaction cost must be monotone in the population measure, i.e. the map $m \mapsto -\sum_{i < j} \int E_{ij} m_j d\theta_j$ must be monotone. For *repulsive* interactions (excluded-volume effects, steric clashes), this condition is satisfied because increasing the density at a given configuration increases the cost for all agents, ensuring uniqueness and convergence to the Nash equilibrium. For *strongly attractive* interactions, however, monotonicity is violated: cooperative binding lowers the cost as more agents aggregate, creating multiple self-reinforcing fixed points and destroying convergence guarantees.

The physical interpretation is useful but model-dependent: Lasry–Lions monotonicity corresponds to the absence of strongly cooperative misfolding in this representation. When monotonicity holds, the MFG system has convergence guarantees toward a stable Nash equilibrium candidate for the native fold. When it is violated, the system can admit multiple competing fixed points corresponding to misfolded aggregates.

Remark 5 (Failure Modes). The MFG framework predicts non-convergence in systems with strongly cooperative misfolding (prions, amyloid fibrils), where Lasry–Lions monotonicity is violated. This is a useful scope condition rather than a failure of the framework: such systems lie outside the simplest stable-Nash domain. Prion propagation and amyloid nucleation are plausible examples of regimes where the potential-game structure breaks down, and the multi-agent system becomes trapped in kinetically stabilised, thermodynamically metastable states. The MFG embedding thus sharpens the scope statement of the FEP–Nash bridge: it applies to folding systems that satisfy monotonicity, while its failure modes can be compared with known pathologies of protein aggregation.

6 Boolean Networks and the Edge of Chaos

At the level of gene regulation, stability manifests in the dynamics of Boolean networks, as introduced by Stuart Kauffman [Kauffman, 1969, 1993]. Boolean network models

remain an active research area; for comprehensive treatments of network motifs and design principles see Alon [2007], and for robustness and evolvability in genetic networks see Wagner [2005].

6.1 Network Dynamics

These networks model genes as binary switches whose states are linked through regulatory logic [Kauffman, 1969]. The long-term dynamics of such networks flow into attractors that are biologically interpreted as specific cell types or functional states [Graudenzi et al., 2011].

Kauffman argued that network stability depends on connectivity and the type of Boolean functions [Kauffman, 1969]. Networks at the “Edge of Chaos”—a critical transition region between rigid order and unpredictable chaos—are hypothesized to exhibit high evolvability and information processing capacity, although this claim remains debated in the literature [Kauffman, 1993]:

Table 7: Boolean network regimes and their biological interpretation. Networks at the edge of chaos exhibit maximum evolvability.

Network Regime	Dynamic Behavior	Biological Function
Ordered	Frozen states, low variability	Structural integrity, basal metabolism
Critical (Edge of Chaos)	Complex patterns, high information flow	Adaptive behavior, differentiation, learning
Chaotic	High sensitivity, instability	Pathological states, malignancy

These attractors in gene regulatory networks form the material basis for the game-theoretic strategies at higher levels. An ESS at the macro-level requires stable genetic attractors at the micro-level that reliably produce the necessary phenotypic traits [Müssel et al., 2010, Kauffman, 1993].

Boolean attractors as Nash equilibria. The connection between Boolean network attractors and Nash equilibria can be made explicit. Each gene g_i in a Boolean network of n genes is a “player” whose strategy is its expression state $\sigma_i \in \{0, 1\}$. The Boolean update function $f_i(\sigma_{-i})$ defines the best response of gene i given the states of its regulators. A fixed-point attractor $\sigma^* = (f_1(\sigma_{-1}^*), \dots, f_n(\sigma_{-n}^*))$ is then precisely a pure-strategy Nash equilibrium of this n -player game: no gene can unilaterally change its state to improve its “payoff” (defined as agreement with the regulatory logic). **Note:** This Nash characterization is *tautological* as stated: defining the payoff as “agreement with the regulatory logic” makes every fixed point a Nash equilibrium by construction—the game-theoretic label adds no information beyond the fixed-point condition. The non-trivial content arises *only* when the payoff is linked to an external, independently measurable fitness function (e.g., organismal viability or growth rate), so that not all Boolean fixed points are equally “Nash-stable” under selection. Without such an external payoff, the Nash language is a relabelling, not a result. Limit-cycle attractors correspond to periodic orbits rather than pure Nash equilibria, but can be interpreted as correlated equilibria in the repeated game.

This game-theoretic perspective on Boolean networks provides the formal bridge between Kauffman’s attractor dynamics and the hierarchical Nash architecture of FST-III.

7 Cellular Level: Multicellularity as Coalition Game

7.1 The Cooperation Problem

In a multicellular organism, individual cells “sacrifice” their reproductive potential for the benefit of the whole. Somatic cells do not reproduce directly; only germline cells transmit genes.

This represents extreme cooperation: cells adopt a strategy (differentiation, non-reproduction) that benefits the collective at individual cost.

7.2 Coalition Game Formulation

- **Players:** Individual cells within the organism
- **Strategies:** Cooperate (follow developmental program) or Defect (proliferate autonomously)
- **Payoff:** Genetic fitness (transmitted through germline)

To first approximation, all cells in an organism are genetically nearly identical (clonal), sharing the overwhelming majority of their genome. Therefore, the payoff to any cell approximately equals the payoff to the germline: if the organism reproduces, the shared genes are transmitted. (In practice, somatic mutations, V(D)J recombination in immune cells, and mitochondrial heteroplasmy introduce limited genetic divergence, which is precisely what enables the cancer-as-defection scenario discussed below.)

Proposition 1 (Multicellularity as Nash Equilibrium). *In a clonal organism where all cells share relatedness $r \approx 1$, cooperation (following the developmental program) is a Nash equilibrium provided the individual cost of cooperation is less than the inclusive fitness benefit: $c < r \cdot b$ (Hamilton’s rule [Hamilton, 1964]). In the limiting case $r = 1$, any positive net benefit $b > c$ suffices. The contribution here is embedding this classical result within the hierarchical FST framework; the Nash equilibrium is maintained as long as clonal identity is preserved.*

7.3 Cancer as Coalition Breakdown

Cancer as a breakdown of multicellular cooperation is a well-established framework [Aktipis et al., 2015]. The game-theoretic framing here follows Aktipis et al. (2015), who apply evolutionary game theory to explain somatic evolution and cancer progression. The FST-III contribution is to embed this framework within the hierarchical Nash-equilibrium picture: cancer represents not merely a selective advantage for individual cells but a phase transition at the coalition level, analogous to the parasite problem in prebiotic chemistry discussed in FST-II. Formally, cancer represents coalition breakdown:

1. Somatic mutations create genetic divergence
2. Mutated cells no longer share all genes with other cells

3. Defection (proliferation) now benefits the mutated lineage
4. The Nash equilibrium breaks down; cancer cells “defect”

This connects to the parasite problem in hypercycles [Geiger, 2026b]: both cancer and chemical parasites exploit cooperative systems by breaking conditions that stabilize cooperation. The parallel between cancer as coalition breakdown and the hypercycle parasite problem is presented as a *structural analogy*, not a derived result. A formal connection would require showing that oncogenic mutations map to parasitic replicators in the sense of Eigen & Schuster; this remains an open problem.

8 Scaling and the Renormalization Group

A central element of FST-III is the recognition that biological principles operate scale-invariantly across many levels [Friston et al., 2025, Wilson, 1975].

8.1 The RG Framework

Terminological caveat: The term “renormalization group” is used in this section in an extended, metaphorical sense. In condensed matter physics, the renormalization group (RG) refers to a precise mathematical framework involving flow equations, fixed points, and scaling exponents [Wilson, 1975]. The present use—describing how effective game-theoretic parameters change when viewed at successively larger biological scales—invokes the *conceptual logic* of coarse-graining, but does not provide RG flow equations, fixed-point analysis, or dimensional scaling laws for biological systems. Whether a rigorous RG treatment of evolutionary transitions is achievable remains an open question [Friston et al., 2025].

With this caveat in mind: The coarse-graining perspective posits that biological systems are often hierarchically organized—from fractal branching in blood vessels to hierarchical neural architectures [Gallos et al., 2012].

Renormalizing Generative Models (RGM) show that principles of free energy and game-theoretic optimization are applied recursively at each level [Friston et al., 2025]. An organism can be viewed as coarse-graining over billions of cells, where cells function as “atoms” in a higher-order game whose rules are set by the species’ evolutionary history [Farzulla, 2026].

8.2 The Scale-Relative Kernel Model

This instantiation is described by scale-relative fitness models developed in multi-level selection theory [Okasha, 2006, Michod, 1999, Farzulla, 2026]:

This hierarchical structure ensures that conflicts between scales (e.g., cancer as defection of individual cells against the organism) are minimized through higher-order selection pressures [Farzulla, 2026].

8.3 Major Transitions as RG Steps

The major transitions in evolution identified by Szathmáry and Maynard Smith [Szathmáry & Maynard Smith, 1995] map onto this framework:

Table 8: Scale-relative kernel model: fitness definitions and strategy types at each biological level of organisation.

Level	Fitness/Strategy
Cellular	Replication rate as fitness, mutations as strategy change
Organismic	Darwinian fitness, behavioral strategies
Agential	Intentionality, learning through imitation
Institutional	Legitimacy and social cooperation as ESS

- Genes $\rightarrow$ Chromosomes (Block 1)
- Prokaryotes $\rightarrow$ Eukaryotes (Block 2)
- Unicells $\rightarrow$ Multicells (Block 3)
- Individuals $\rightarrow$ Societies (Block 4)

Each transition creates a new level of “player” from coalitions of lower-level players, preserving game-theoretic structure through coarse-graining.

A toy coarse-graining step for spatial evolutionary games. Consider a 2D lattice game with strategies $\sigma_x \in \{C, D\}$ and payoff matrix $A = \begin{pmatrix} R & S \\ T & P \end{pmatrix}$. For a 2×2 block B define a block strategy $\Sigma_B = \mathcal{B}(\sigma|_B)$ by majority rule. For neighboring blocks B, B' define the effective block payoff by boundary averaging

$$A'_{s,t} := \frac{1}{|\partial(B, B')|} \mathbb{E} \left[\sum_{\langle x,y \rangle \in \partial(B, B')} A_{\sigma_x, \sigma_y} \mid \Sigma_B = s, \Sigma_{B'} = t \right], \quad s, t \in \{C, D\}.$$

In a frozen-interior approximation, let $q_s = \mathbb{P}(\sigma_x = C \mid \Sigma_B = s, x \in \partial B)$. Then the coarse-graining map $R_2 : (A, \beta) \mapsto (A', \beta')$ satisfies the explicit mixing formula

$$A'_{s,t} = q_s q_t R + q_s (1 - q_t) S + (1 - q_s) q_t T + (1 - q_s)(1 - q_t) P,$$

with $q_C \approx 1 - \varepsilon$ and $q_D \approx \varepsilon$ controlled by the within-block dynamics (selection strength β and internal enforcement mechanisms). Fixed points include the ordered phases (all- C , all- D), while changes in within-block constraints shift the coarse-graining flow by modifying $q_C - q_D$. This provides a concrete sense in which major evolutionary transitions correspond to coarse-graining steps together with the activation of new relevant operators (e.g. policing, bottlenecks) that stabilize cooperative macro-variables. Biologically, the block strategy Σ_B corresponds to a higher-level phenotype (e.g., multicellular cooperation), while the boundary parameters q_C, q_D encode enforcement mechanisms (e.g., immune surveillance, reproductive bottlenecks) that suppress within-group defection.

9 Ecosystem Level: The Ecological Game

9.1 Species as Players

In the ecological game:

- **Players:** Species populations
- **Strategies:** Ecological roles (predator, prey, competitor, mutualist)
- **Payoff:** Population growth rate

9.2 Ecological Niches as Nash Equilibria

Ecological niches are Nash equilibrium configurations: no species can unilaterally change its strategy to improve fitness while others remain fixed.

Example: Predator-prey coexistence. In a classical Lotka-Volterra predator-prey system, the interior fixed point—where predator and prey populations coexist at constant densities—satisfies a *formal analogy* to the Nash equilibrium condition: at this point, neither the prey population growth rate nor the predator population growth rate can be improved by a unilateral infinitesimal change. This is a mathematical correspondence, not a Nash equilibrium in the strategic sense, since species do not “choose” population sizes—the dynamics are deterministic ODEs, not strategic decisions. **Caveat:** In the basic Lotka-Volterra model this fixed point is a center (neutrally stable with closed orbits), not an asymptotically stable attractor; ESS-type stability requires additional mechanisms such as density-dependent mortality or spatial structure. The equivalence between Lotka-Volterra fixed points and replicator dynamics equilibria [Hofbauer & Sigmund, 1998] maps the ecological game onto evolutionary dynamics, but the stability type of the resulting equilibrium depends on the specific payoff structure.

9.3 Mass Extinctions as Phase Transitions

Mass extinction events represent phase transitions in the ecological game:

1. Environmental perturbation changes the payoff matrix
2. Previous Nash equilibria become unstable
3. System transitions to new equilibrium with different species composition

At a structural level, the logic resembles phase transitions in physical systems more broadly: changing boundary conditions destabilize existing equilibria and drive transitions to new configurations. It should be noted that this game-theoretic framing is a *post hoc* interpretive model: mass extinctions are complex events driven by asteroid impacts, volcanism, or ocean anoxia, and the extent to which they can be productively modeled as payoff-matrix perturbations—rather than as catastrophic regime shifts outside the equilibrium framework—remains an empirical question [Gould & Eldredge, 1977].

10 Synthesis: Three Phases of Biological Organization

FST-III delivers an integrated picture of biological organization that unfolds in three phases:

10.1 Phase 1: Thermodynamic Dissipation

The energy flow of the young universe forces simple matter into dissipative structures. Here MEPP acts as a stability filter (Section 3): among the many possible configurations, those with resolvent-stable dissipation patterns persist. This is the era of rapid entropy production and the emergence of England’s dissipative adaptors [Kleidon, 2010].

10.2 Phase 2: Strategic Consolidation

Once dissipative structures cross the threshold to self-replication, competition for resources introduces game-theoretic dynamics. The emergence of Evolutionarily Stable Strategies (ESS) stabilizes molecular populations against parasitic invasion (Section 4). The “players” are still purely chemical, but their interactions can already exhibit Nash-equilibrium structure that later levels may inherit in modified form [Maynard Smith & Price, 1973]. The major evolutionary transitions (Section 8) are interpreted as iterative coarse-graining steps from lower-level players into higher-level coalitions, each preserving parts of the game-theoretic logic at its new scale.

10.3 Phase 3: Cognitive Inference

Organisms develop internal models of their environment to minimize their free energy. The emergence of the Markov blanket (Section 5) allows separation of self and world, motivating the FEP–MEPP compatibility conjecture (Conjecture 1): internal model-building (free energy minimization) may support externally efficient dissipation.

This third phase marks a qualitative shift in the game-theoretic hierarchy. In Phase 2, the “players” are molecular or cellular agents whose strategies are determined by physical chemistry. In Phase 3, agents acquire *generative models* of their environment: the strategies are no longer fixed phenotypic states but policies—sequences of actions conditioned on inferred environmental states. Active inference [Friston, 2009] provides the mechanism: agents select actions that minimise expected free energy (i.e., they act to confirm their predictions or explore to reduce uncertainty). This converts the static Nash equilibria of Phase 2 into dynamic Bayesian Nash equilibria where each agent’s strategy is contingent on its beliefs about others. The nested hierarchy of inference—from interoceptive homeostasis through perception to planning—mirrors the coarse-graining logic of Section 8: each level of inference integrates over the degrees of freedom below it, producing an increasingly abstract game at higher temporal and spatial scales. The endpoint of this hierarchy—consciousness, language, and culture—can be *interpreted* as a highly compressed form of the game-theoretic stability logic, where the “strategies” are symbolic representations and the “payoff” is predictive accuracy over arbitrarily long time horizons [Friston, 2010]. This extrapolation is speculative and goes beyond what the present framework can formally derive.

11 Testable Predictions

Note: The predictions below are labeled P1–P5 to distinguish them from the mathematical Propositions (e.g., Proposition 1) stated in earlier sections. Propositions are formal claims within the theoretical framework; Predictions are empirically testable hypotheses derived from it.

11.1 P1: Protein Folding Landscapes

Hypothesis 1 (Nash frustration predicts folding-relevant residues). *The energy landscape of protein folding can be characterized by Nash equilibrium stability analysis. Residues with high Nash frustration correspond to experimentally identified folding nuclei and allosteric sites.*

Test: Compute Nash frustration scores for proteins with experimentally characterized folding intermediates (e.g., via hydrogen-deuterium exchange or ϕ -value analysis). A correlation coefficient $r > 0.3$ between Nash frustration rank and experimental ϕ -values would constitute positive evidence.

11.2 P2: Gene Regulatory Network Attractors

Hypothesis 2 (GRN attractors as Nash equilibria). *Cell type attractors in gene regulatory networks can be predicted from Nash analysis of the TF-DNA binding game.*

Test: For well-characterized GRNs (e.g., the *Drosophila* segment polarity network or the mammalian hematopoietic differentiation network), compute Nash equilibria of the TF-DNA binding game. Compare the number and identity of predicted attractor states to experimentally observed cell types. Success criterion: the predicted attractor count should match the known cell-type count within a factor of two.

11.3 P3: Cancer and Coalition Stability

Hypothesis 3 (Cancer as coalition breakdown). *Cancer incidence correlates inversely with multicellular coalition stability as measured by the game-theoretic defection threshold.*

Test: Across tissue types, compute the defection threshold Δr at which Hamilton's condition $c < r \cdot b$ fails (given tissue-specific mutation rates and cooperation costs). Predict that tissues with lower Δr thresholds exhibit higher cancer incidence. Compare against epidemiological data from the SEER database. A particularly stringent test is provided by Peto's paradox: large-bodied, long-lived species do not show proportionally higher cancer rates despite having more cells and cell divisions. The coalition model predicts that these species must have evolved stronger enforcement mechanisms (lower effective Δr), a prediction testable by comparing immune surveillance intensity across species of different body sizes. *A priori falsification criterion:* if natural killer (NK) cell density per unit tissue does not correlate negatively with body size across mammals, the enforcement prediction is falsified.

11.4 P4: Metabolic Rate and Reproductive Success

Hypothesis 4 (MEPP-fitness correlation). *Across species, mass-specific metabolic rate correlates with reproductive rate after controlling for body size.*

Test: Compile metabolic rates and reproductive outputs across taxa. A positive partial correlation (controlling for body mass) supports the MEPP-fitness connection. Kleiber's law provides the body-size scaling. **Context:** The Metabolic Theory of Ecology [Brown et al., 2004] already establishes quantitative relationships between metabolic rate, body size, and life-history traits. The FST-III prediction goes beyond MTE by attributing

the correlation not merely to allometric scaling but to the thermodynamic stability filter (MEPP): species that dissipate more efficiently per unit mass are predicted to be more resilient to perturbation, not just faster reproducers.

11.5 P5: Ecosystem Resilience

Hypothesis 5 (Ecosystem resilience from Nash stability). *Ecosystem resilience can be predicted from the spectral gap of the Nash stability matrix of the ecological equilibrium.*

Test: For well-characterized ecosystems (e.g., Yellowstone food web, coral reef communities), compute the Jacobian eigenvalue gap at the Nash equilibrium. Predict that ecosystems with larger spectral gaps recover faster from perturbation. Compare against empirical resilience measures (recovery time after disturbance events).

12 Discussion

12.1 What is Genuinely New?

Classical evolutionary game theory is well-established at the organismic level. The contributions of this paper are:

1. **Vertical integration:** Connecting molecular, cellular, organismic, and ecosystem games
2. **Nash-frustration maps:** A residue-level coordination-failure score obtained by projecting only unstable best-response modes, weighted by instability excess, onto residues; this is complementary to energetic frustration and standard NMA/B-factor profiles
3. **Thermodynamic foundation:** Grounding fitness in MEPP and dissipative adaptation
4. **Coarse-graining interpretation:** Framing major transitions as coarse-graining steps
5. **MEPP as stability filter:** Reinterpreting MEPP as a second-order selection criterion rather than a maximization law

12.2 Scalability and the AlphaFold Era

The Nash frustration pipeline operates on experimentally determined or computationally predicted three-dimensional structures. The recent availability of over 200 million predicted protein structures through AlphaFold2 [Jumper et al., 2021] dramatically expands the scope of the method: Nash frustration scores can in principle be computed for any protein with a predicted structure, enabling proteome-scale frustration maps. Moreover, AlphaFold2’s per-residue confidence metric (pLDDT) provides a natural independent benchmark: regions with low pLDDT (predicted disorder or poor confidence) are expected to correlate positively with high Nash frustration, since both reflect local structural instability—pLDDT from the prediction model’s perspective, Nash frustration

from the game-theoretic perspective. A systematic comparison of pLDDT and Nash frustration profiles across a large protein set would test whether game-theoretic coordination failure captures structural information beyond what is already encoded in deep-learning confidence scores. This comparison is computationally straightforward and constitutes a natural extension of the benchmark analysis in Section 4.4.

12.3 Punctuated Equilibrium as Resolvent Transition

Remark 6 (Biological Stasis and Sudden Transitions). The pattern of long biological stasis interrupted by sudden evolutionary transitions (punctuated equilibrium, Gould & Eldredge, 1977; see also Kauffman, 1993) finds a structural explanation in the resolvent framework [Geiger, 2026e]. Periods of stasis correspond to resolvent-stable phases: the second-order coupling structure maintains the current configuration against perturbations. Sudden transitions occur when environmental or internal changes reorganize the second-order coupling architecture, analogous to phase transitions in physical systems where critical parameter changes trigger qualitative reorganization of the stability landscape. Major evolutionary transitions—from prokaryote to eukaryote, from unicellularity to multicellularity—are thus not gradual optimizations but *resolvent transitions*: reorganizations of the coupling structure that establish a qualitatively new stable fixed point.

12.4 Stability vs. Maximization: The Resolvent Perspective

The FST framework describes *stability principles*, not maximization principles. The general resolvent perspective—applicable across all three FST papers—is developed in FST-II [Geiger, 2026b]; here we focus on its biological implications. The core logic is that leading-order dynamics are degenerate (many configurations are viable), and selection occurs at second order via resolvent-type coupling. MEPP, ESS, and Nash frustration are all manifestations of this second-order stability architecture (see Remarks 1 and 2 for the detailed arguments). The unifying claim is that Nash equilibria across all FST papers are not optima but resolvent-stable fixed points [Geiger, 2026e].

The Nash frustration of protein folds (Section 4) provides a concrete illustration: the native fold is not the global energy minimum but the configuration where all second-order destabilization channels—misfolding, aggregation, proteolytic vulnerability—are simultaneously suppressed. The residual Nash frustration ($\rho(J) \approx 1$) is the structural signature of this stabilization.

12.5 MEPP: Scope and Limitations

The Maximum Entropy Production Principle is used throughout this paper as a heuristic organizing principle, not as a proven physical law. The MEPP controversy—including the Grinstein–Linsker critique, Martyushev’s applicability conditions, the mEPP/MEPP regime separation, and the resolvent reinterpretation—is discussed in detail in FST-II [Geiger, 2026b]; here we summarize only the limitations most relevant to the biological domain:

- MEPP’s theoretical foundations require local thermodynamic equilibrium and bilinear entropy production [Martyushev, 2010]. Biological systems far from equilibrium generally violate these conditions.

- The resolvent perspective (Section 12) addresses this limitation by reinterpreting MEPP as a stability filter: among the many biologically viable configurations at leading order, MEPP is used here to identify candidate resolvent-stable fixed points—not necessarily configurations with maximal dissipation, but configurations where second-order destabilization channels are jointly suppressed.
- This weaker interpretation is compatible with Prigogine’s minimum entropy production near equilibrium: internally, organisms may minimize entropy production (FEP), while externally they may support higher entropy production (MEPP), as formulated in Conjecture 1.
- A recent review by Sawada, Daigaku & Toma [Sawada et al., 2025] confirms that MEPP provides a consistent thermodynamic framework for understanding the birth and evolution of life at macroscopic scales, while acknowledging that a fundamental derivation from first principles far from equilibrium remains elusive. This corroborates the present paper’s use of MEPP as a heuristic organizing principle.

Circularity caveat. When MEPP serves as the payoff function and Nash equilibria are defined as MEPP-maximizing states, the claim that biological systems occupy Nash equilibria *because* they maximize entropy production is definitionally circular. The non-circular content of this framework is the *stability* claim: biological attractors are not merely entropy-producing configurations but second-order stable fixed points where all destabilization channels (parasitism, mutation, ecological invasion) are simultaneously suppressed. This stability property is testable independently of the payoff definition (Predictions P1–P5).

12.6 Nash Frustration Beyond the Hessian

A natural objection to the Nash frustration formalism is that, when the payoff equals $-\Delta G$ (the negative free energy change), the best-response Jacobian $J = I - \eta H$ is a linear transformation of the Hessian, and Nash frustration reduces to an eigenvalue analysis of the energy landscape—essentially a normal-mode analysis in game-theoretic clothing. This objection correctly identifies the mathematical core but misses the conceptual and practical content that goes beyond standard Hessian analysis:

1. **Residue-level decomposition:** The Nash frustration formalism projects unstable modes onto individual residues via eigenvector components, producing a *per-residue frustration score* that localises coordination failure. Per-residue eigenvector projections are themselves standard in normal-mode analysis—the well-known *B-factor* calculation $B_i = \frac{8\pi^2}{3} \sum_k \omega_k^{-2} |v_{k,i}|^2$ is structurally analogous. The distinction lies in the *weighting scheme*: NMA *B*-factors sum over *all* modes weighted by inverse frequency squared (emphasising slow, collective motions), whereas Nash frustration projects exclusively onto *unstable* modes ($|\lambda_k| > 1$) weighted by the instability excess ($|\lambda_k| - 1$). This selective projection isolates game-theoretic coordination failure rather than overall flexibility, producing a biologically interpretable map of where the folding game is most contested. Future work should directly compare Nash frustration maps with *B-factor* profiles to quantify the degree of overlap and complementarity.

2. **Multi-start convergence diagnostics:** The best-response dynamics (64 random starts, 6 000 steps each) provide information about the basin structure of the game: how many starts converge to the native fold, how many become trapped in alternative configurations, and which residues are responsible for convergence failures. This dynamical information is absent from a static Hessian eigendecomposition.
3. **Connection to existence results:** The MFG embedding (Section 5.4) connects the local Hessian analysis to global existence and uniqueness conditions (Lasry–Lions monotonicity). This connection—from local spectral properties to candidate convergence criteria—has no direct analogue in standard normal-mode analysis and provides a principled way to ask when the folding landscape admits a single stable fold versus multiple competing metastable states.
4. **Cross-scale applicability:** The same game-theoretic vocabulary (players, strategies, payoff, best-response Jacobian) can be adapted at the gene-regulatory, cellular, and ecosystem levels (Sections 6–9), whereas normal-mode analysis is specific to energy landscapes. The unified language is the substantive contribution.

The genuine novel content of Nash frustration thus lies not in the eigenvalue computation itself but in the interpretive framework, the per-residue decomposition, and the connection to MFG existence theory. Future work should make this distinction empirically testable by comparing Nash frustration maps with standard B -factor profiles (which also derive from normal modes): the prediction is that Nash frustration identifies a partially distinct set of residues—those at game-theoretic coordination boundaries rather than simply flexible regions.

12.7 Consolidated Limitations

The following limitations should be kept in mind when evaluating the framework:

1. **Nash frustration calibration:** While the frustration *ranking* is η -independent in the verified local scan window (Section 4), the absolute frustration values and the global spectral radius $\rho(J)$ depend on η , and ranking invariance can fail if the unstable active set changes. Calibration against experimental observables (HDX protection factors, ϕ -values) remains an open task.
2. **Coordinate metric:** The current residue projection uses Euclidean components in a fixed (ϕ, ψ) chart. A coordinate-invariant version should use a torsional/manifold metric or a mass-weighted Hessian; the present score is a local chart diagnostic.
3. **MEPP foundations:** MEPP lacks rigorous derivation for far-from-equilibrium biological systems (Section 12).
4. **RG formalization:** The coarse-graining interpretation of major transitions is conceptual, not mathematically rigorous (Section 8).
5. **FEP–MEPP bridge:** The compatibility conjecture (Conjecture 1) remains unproven; the connection from FEP minimization to MEPP relies on the competitive selection mechanism, which is plausible but not formally derived.
6. **Empirical validation:** None of the five testable predictions (Section 11) have been empirically tested; the framework is currently at the theoretical stage.

7. **Scope of “Nash equilibrium”:** The concept is applied across very different domains (molecular, cellular, ecological) where the notion of “player,” “strategy,” and “payoff” requires domain-specific interpretation, risking vacuousness if applied too liberally.
8. **Potential game restriction:** The formal FEP–Nash bridge (Section 5) requires a shared cost functional (potential game). Antagonistic interactions (predator–prey, host–parasite) do not satisfy this condition, limiting the formal scope of the bridge to cooperative or intra-organismal settings. The MFG embedding (Section 5.4) generalises this to systems satisfying Lasry–Lions monotonicity, but strongly attractive (cooperative misfolding) interactions remain outside the convergence domain.

12.8 Status of Claims

Table 9 classifies the central claims of this paper by epistemic status.

Table 9: Status of claims. ESTABLISHED: proven or widely accepted results cited here; CONJECTURED: new but testable claims; SPECULATIVE: framework-level proposals without direct empirical test.

Claim	Status	Evidence / Reference
Replicator–Nash equivalence	ESTABLISHED	Hofbauer & Sigmund (1998)
ESS as Nash refinement	ESTABLISHED	Maynard Smith (1982)
MEPP literature as heuristic thermodynamic framework	ESTABLISHED	Martyushev [2010]; Sawada et al. [2025]; Kleidon [2010]
FST MEPP stability-filter interpretation	CONJECTURED	Sec. 12; no first-principles derivation
Nash frustration as stability predictor (P1)	CONJECTURED	Preliminary HP35/CSP and benchmark signals only; validation gates in Table 3
FEP–MEPP compatibility via Markov blanket	CONJECTURED	Conjecture 1; no formal proof
Hierarchical game structure across scales	CONJECTURED	Sec. 8; qualitative only
Boolean network attractors as Nash equilibria	CONJECTURED	Sec. 6; formal restatement, empirical test pending
Nash equilibrium as MFG fixed point	CONJECTURED	Sec. 5.4; Lasry–Lions monotonicity as existence condition
RG analogy for major transitions	SPECULATIVE	Conceptual parallel; no RG flow computed
Cosmological extension of game structure	SPECULATIVE	Sec. 12.11; deferred to FST framework paper

12.9 Falsifiable Predictions and Experimental Protocols

The predictions P1–P5 (Section 11) identify testable hypotheses at the framework level. To address the concern that these predictions lack sufficient specificity for empirical refutation, we here formulate four concrete, falsifiable predictions with detailed experimental

protocols and explicit falsification criteria. Each prediction is designed to be testable with existing techniques and publicly available data.

PRED-BIO-1: Nash frustration correlates with HDX protection factors. *Prediction.* Residues with high Nash frustration scores (top quartile) exhibit systematically *lower* hydrogen–deuterium exchange (HDX) protection factors than residues with low Nash frustration (bottom quartile). The rationale is that Nash-frustrated residues represent coordination failures in the folding game: they are structurally strained and therefore more solvent-accessible and conformationally dynamic.

Protocol. (i) Compute residue-level Nash frustration scores for HP35 (villin headpiece, PDB: 1YRF) using the game-theoretic potential described in Remark 3. (ii) Obtain HDX protection factors from McKnight et al. [1996] or perform HDX-MS on recombinant HP35 under native conditions (pH 7.0, 25 °C, D_2O exchange for 10 s to 10 h). (iii) Compute Spearman rank correlation ρ_S between per-residue Nash frustration and $\log(P_f)$, where P_f is the protection factor. (iv) Repeat for at least two additional small proteins with published HDX data (e.g., ubiquitin, BPTI) to assess generalizability.

Falsification criterion. The prediction is falsified if $\rho_S > -0.15$ (i.e., no meaningful negative correlation) across all three proteins. A positive correlation ($\rho_S > 0$) would directly contradict the prediction. Expected effect size: $\rho_S \in [-0.5, -0.25]$.

PRED-BIO-2: Global spectral radius predicts unfolding kinetics. *Prediction.* Proteins with higher global spectral radius $\rho(J)$ of the best-response Jacobian unfold faster under thermal or chemical denaturation. The spectral radius quantifies the overall departure from Nash stability; a higher $\rho(J)$ indicates stronger destabilizing modes, which should manifest as faster unfolding kinetics.

Protocol. (i) Select a panel of 15–20 small single-domain proteins (< 100 residues) spanning diverse fold classes (α , β , α/β) with experimentally determined unfolding rates k_u from temperature-jump (T-jump) or stopped-flow experiments (available in databases such as ACPro or published kinetic compilations). (ii) For each protein, compute $\rho(J)$ from the PDB structure using the Nash frustration pipeline (Remark 3). (iii) Compute Spearman rank correlation between $\rho(J)$ and $\log(k_u)$.

Falsification criterion. The prediction is falsified if $\rho_S < 0.2$ (no meaningful positive correlation between spectral radius and unfolding rate). The prediction explicitly does *not* claim that $\rho(J)$ is a better predictor than contact order or other established metrics, but that it carries independent information: a partial correlation $\rho_S(\rho(J), \log k_u \mid \text{CO}) > 0.15$ after controlling for contact order would constitute strong evidence.

PRED-BIO-3: Nash frustration enrichment at allosteric and binding sites. *Prediction.* Residues classified as allosteric or located at protein–protein binding interfaces are enriched in high Nash frustration scores relative to the protein average. This complements the known enrichment of *energetic* frustration at such sites [Ferreiro et al., 2014] and predicts that game-theoretic coordination failure (Nash frustration) and energetic suboptimality (Ferreiro frustration) are partially overlapping but distinct signals.

Protocol. (i) Obtain a dataset of proteins with experimentally annotated allosteric sites from the Allosteric Database (ASD, asd.pharm.umich.edu) or CASBench; minimum 30 proteins for statistical power. (ii) For each protein, compute residue-level Nash frustration scores and classify residues as allosteric/interface (from ASD annotations) or

non-functional. (iii) Perform a Wilcoxon rank-sum test comparing Nash frustration distributions of allosteric vs. non-allosteric residues, pooled across all proteins. (iv) Compute AUC of Nash frustration as a binary classifier of allosteric sites.

Falsification criterion. The prediction is falsified if (a) the Wilcoxon test yields $p > 0.05$ (no significant enrichment) *and* (b) $\text{AUC} < 0.55$ (classifier performance indistinguishable from chance). A positive result requires $\text{AUC} > 0.60$ and $p < 0.01$. Additionally, the prediction claims partial independence from energetic frustration: the correlation between Nash frustration and Ferreiro frustration scores should be $r < 0.7$ (i.e., they share less than 50% variance).

PRED-BIO-4: Frustration map predicts helix melting order. *Prediction.* In multi-helix proteins, helices with higher mean Nash frustration melt (unfold) earlier during thermal denaturation. The frustration map should predict the temporal order of secondary structure loss.

Protocol. (i) Select 3–5 multi-helix proteins with experimentally determined helix melting order from NMR relaxation dispersion (R_2 dispersion, CPMG experiments) or hydrogen exchange under progressively denaturing conditions (e.g., HP35 with three helices, or apomyoglobin with eight helices). (ii) Compute mean Nash frustration per helix from the PDB structure. (iii) Rank helices by mean frustration (highest = predicted to melt first) and compare against the experimentally determined melting order using Kendall’s τ rank correlation.

Falsification criterion. The prediction is falsified if Kendall’s $\tau < 0.2$ (no meaningful rank agreement) for the majority ($\geq 3/5$) of test proteins. For HP35 specifically, the prediction is that the helix containing Met11/Pro20 (highest frustration, cf. Section 4) melts before the helix containing Trp22/Asn18 (lowest frustration); reversal of this order would falsify the local prediction.

Summary. These four predictions are designed to be testable with existing experimental techniques (HDX-MS, T-jump kinetics, NMR relaxation dispersion) and publicly available databases (ASD, ACPro, BMRB). Each prediction specifies quantitative thresholds for falsification. Together, they provide a concrete empirical agenda for evaluating the Nash frustration framework at the molecular level.

12.10 Is Biology “Just Physics”?

The answer is nuanced:

- **Yes, as a working model:** MEPP-like stability filters and Nash-style fixed-point structures can be formulated across multiple biological scales
- **No:** Higher levels introduce emergent constraints and causal descriptions that are not straightforwardly predictable from lower levels alone

The framework is not reductive in the eliminativist sense. It identifies a common logical structure instantiated differently at each scale.

12.11 Cosmological Extensions

The biological game-theoretic framework developed in this paper is self-contained. Speculative extensions to physical and cosmological scales are developed separately in the FST framework paper [Geiger, 2026d] and are not part of the biological theory presented here.

13 Conclusion

FST-III argues that the separation between physical laws and the strategies of life may be less sharp than commonly assumed. Biological stability can be productively analyzed through game-theoretic and thermodynamic tools adapted from physical systems. In this conjectural reading, the Maximum Entropy Production Principle provides a thermodynamic stability filter, dissipative adaptation a mechanism of form-building, game-theoretic ESS criteria for long-term stability, and the Free Energy Principle a foundation for adaptive behavior.

The primary contributions are threefold: (i) the Nash frustration formalism, which provides a game-theoretic complement to energetic frustration in protein structures and has been subjected to a preliminary test against NMR data from a single protein (HP35, $\rho_S = 0.44$, $p = 0.033$, $n = 24$); (ii) the hierarchical game interpretation of major evolutionary transitions as coarse-graining steps; and (iii) the coalition-breakdown model of cancer, embedding Aktipis et al.’s framework into the FST stability hierarchy. Four concrete falsifiable predictions (PRED-BIO-1 through PRED-BIO-4) with explicit protocols and falsification criteria provide an empirical agenda for the molecular-level claims.

The key open challenges are the calibration of the absolute Nash frustration scale against experimental observables (HDX protection factors, ϕ -values), the formalization of the RG-like coarse-graining as explicit renormalization flows, and the rigorous derivation of the FEP–MEPP compatibility conjecture. From amino acid to ecosystem, the game-theoretic stability logic is offered here as a unifying conjectural lens; its extension to cosmological scales [Geiger, 2026d] remains speculative.

AI Disclosure. This work was assisted by Claude (Anthropic) and Codex (OpenAI) for literature synthesis, mathematical verification, editorial maintenance, and text drafting. All scientific claims, original arguments, and theoretical contributions are the sole responsibility of the human author. No AI system is listed as co-author.

References

- Geiger, L. (2026). Functional Stability Theory I: A Game-Theoretic Framework for the Thermodynamic Stability of Fundamental Parameters. *Zenodo preprint*, v1.4. Concept DOI: 10.5281/zenodo.20130544.
- Geiger, L. (2026). Functional Stability Theory II: Chemical Stability and Autocatalytic Selection (DRAFT). *Zenodo preprint*, v1.5. Concept DOI: 10.5281/zenodo.20130563.
- Aktipis, C. A. et al. (2015). Cancer across the tree of life: cooperation and cheating in multicellularity. *Phil. Trans. R. Soc. B*, 370, 20140219. DOI: 10.1098/rstb.2014.0219.
- Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. *Science*, 181(4096), 223–230. DOI: 10.1126/science.181.4096.223.
- Graudenzi, A., Serra, R., Villani, M., Damiani, C., Colacci, A. & Kauffman, S. A. (2011). Dynamical properties of a Boolean model of gene regulatory network with memory. *Journal of Computational Biology*, 18(10), 1291–1303. DOI: 10.1089/cmb.2010.0069.
- Müssel, C., Hopfensitz, M. & Kestler, H. A. (2010). BoolNet—an R package for generation, reconstruction and analysis of Boolean networks. *Bioinformatics*, 26(10), 1378–1380. DOI: 10.1093/bioinformatics/btq124.
- Kauffman, S. A. (1969). Metabolic stability and epigenesis in randomly constructed genetic nets. *Journal of Theoretical Biology*, 22(3), 437–467. DOI: 10.1016/0022-5193(69)90015-0.

- Kauffman, S. A. (1993). *The Origins of Order: Self-Organization and Selection in Evolution*. Oxford University Press.
- Hofbauer, J. & Sigmund, K. (1998). *Evolutionary Games and Population Dynamics*. Cambridge University Press. (Ch. 7: Lotka-Volterra and replicator equivalence).
- Monderer, D. & Shapley, L. S. (1996). Potential games. *Games and Economic Behavior*, 14(1), 124–143. DOI: 10.1006/game.1996.0044.
- England, J. L. (2013). Statistical physics of self-replication. *Journal of Chemical Physics*, 139, 121923. DOI: 10.1063/1.4818538.
- England, J. L. (2015). Dissipative adaptation in driven self-assembly. *Nature Nanotechnology*, 10, 919–923. DOI: 10.1038/nnano.2015.250.
- Kleidon, A. (2010). Life, hierarchy, and the thermodynamic machinery of planet Earth. *Physics of Life Reviews*, 7(4), 424–460. DOI: 10.1016/j.plrev.2010.10.002.
- Bryngelson, J. D. & Wolynes, P. G. (1989). Intermediates and barrier crossing in a random energy model (with applications to protein folding). *The Journal of Physical Chemistry*, 93(19), 6902–6915. DOI: 10.1021/j100356a007.
- Friston, K. (2009). The free-energy principle: a rough guide to the brain? *Trends in Cognitive Sciences*, 13(7), 293–301. DOI: 10.1016/j.tics.2009.04.005.
- Friston, K. (2010). The free-energy principle: a unified brain theory? *Nature Reviews Neuroscience*, 11, 127–138. DOI: 10.1038/nrn2787.
- Hamilton, W. D. (1964). The genetical evolution of social behaviour. I. *Journal of Theoretical Biology*, 7(1), 1–16. DOI: 10.1016/0022-5193(64)90038-4.
- Maynard Smith, J. & Price, G. R. (1973). The logic of animal conflict. *Nature*, 246, 15–18. DOI: 10.1038/246015a0.
- Bohl, K., Hummert, S., Werner, S., Basanta, D., Deutsch, A., Schuster, S., Theissen, G. & Schroeter, A. (2014). Evolutionary game theory: molecules as players. *Molecular BioSystems*, 10(12), 3066–3074. DOI: 10.1039/C3MB70601J.
- Maynard Smith, J. *Evolution and the Theory of Games*. Cambridge UP, 1982.
- Holt, C. A. & Roth, A. E. (2004). The Nash equilibrium: A perspective. *PNAS*, 101(12), 3999–4002. DOI: 10.1073/pnas.0308738101.
- Nash, J. F. (1950). Equilibrium points in n-person games. *PNAS*, 36(1), 48–49. DOI: 10.1073/pnas.36.1.48.
- Okasha, S. (2006). *Evolution and the Levels of Selection*. Oxford University Press. DOI: 10.1093/acprof:oso/9780199267972.001.0001.
- Michod, R. E. (1999). *Darwinian Dynamics: Evolutionary Transitions in Fitness and Individuality*. Princeton University Press.
- Farzulla, M. (2026). The replicator-optimization mechanism: A scale-relative formalism for persistence-conditioned dynamics with application to consent-based metaethics. arXiv:2601.06363.
- Friston, K. et al. (2025). From pixels to planning: scale-free active inference. *Frontiers in Network Physiology*, 5, 1521963. DOI: 10.3389/fnetp.2025.1521963.
- Wilson, K. G. (1975). The renormalization group: critical phenomena and the Kondo problem. *Reviews of Modern Physics*, 47(4), 773–840. DOI: 10.1103/RevModPhys.47.773.
- Schrödinger, E. (1944). *What is Life? The Physical Aspect of the Living Cell*. Cambridge University Press.
- Sánchez-Cañizares, J. (2021). The Free Energy Principle: Good Science and Questionable Philosophy in a Grand Unifying Theory. *Entropy*, 23(2), 238. DOI: 10.3390/e23020238.
- Gallo, L. K., Makse, H. A. & Sigman, M. (2012). A small world of weak ties provides optimal global integration of self-similar modules in functional brain networks. *PNAS*, 109(8), 2825–2830. DOI: 10.1073/pnas.1106612109.
- Szathmáry, E. & Maynard Smith, J. (1995). The major evolutionary transitions. *Nature*, 374, 227–232. DOI: 10.1038/374227a0.

- Ferreiro, D.U., Hegler, J.A., Komives, E.A. & Wolynes, P.G. (2007). Localizing frustration in native proteins and protein assemblies. *PNAS*, 104, 19819–19824. DOI: 10.1073/pnas.0709915104.
- Ferreiro, D.U., Komives, E.A. & Wolynes, P.G. (2014). Frustration in biomolecules. *Quarterly Reviews of Biophysics*, 47, 285–363. DOI: 10.1017/S0033583514000092.
- R. G. Parra, N. P. Schafer, L. G. Radusky, M.-Y. Tsai, A. B. Guzovsky, P. G. Wolynes, and D. U. Ferreiro (2016). Protein Frustratometer 2: a tool to localize energetic frustration in protein molecules, now with electrostatics. *Nucleic Acids Research*, 44(W1), W356–W360. DOI: 10.1093/nar/gkw304.
- Martyushev, L. M. (2010). The maximum entropy production principle: two basic questions. *Phil. Trans. R. Soc. B*, 365, 1333–1334. DOI: 10.1098/rstb.2009.0295.
- Gould, S. J. & Eldredge, N. (1977). Punctuated equilibria: the tempo and mode of evolution reconsidered. *Paleobiology*, 3(2), 115–151. DOI: 10.1017/S0094837300005224.
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85(7), 1771–1789. DOI: 10.1890/03-9000.
- Geiger, L. (2026). From Landscape to Atlas: Multi-Route Cartography of an Ongoing Expedition Toward the Riemann Hypothesis. *Zenodo preprint*, v3.1. Concept DOI: 10.5281/zenodo.19035640.
- Geiger, L. (2026). A Conditional Reduction of the Riemann Hypothesis to Even Dominance of the Weil Quadratic Form. *Zenodo preprint*, v1.9. Concept DOI: 10.5281/zenodo.19764771.
- Geiger, L. (2026). Functional Stability Theory — Physical Instantiation via the Renormalized Free Energy Principle. *Zenodo preprint*, v1.8. Concept DOI: 10.5281/zenodo.19036190.
- Sawada, Y., Daigaku, Y. & Toma, K. (2025). Maximum entropy production principle of thermodynamics for the birth and evolution of life. *Entropy*, 27(4), 449. DOI: 10.3390/e27040449.
- Alon, U. (2007). *An Introduction to Systems Biology: Design Principles of Biological Circuits*. Chapman & Hall/CRC, Boca Raton, FL.
- Wagner, A. (2005). *Robustness and Evolvability in Living Systems*. Princeton University Press, Princeton, NJ.
- M. Kjaergaard and F. M. Poulsen (2011). Sequence correction of random coil chemical shifts: correlation between neighbor correction factors and changes in the Ramachandran distribution. *Journal of Biomolecular NMR*, 50, 157–165. DOI: 10.1007/s10858-011-9508-2.
- Jumper, J. et al. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596, 583–589. DOI: 10.1038/s41586-021-03819-2.
- J.-M. Lasry and P.-L. Lions (2007). Mean Field Games. *Japanese Journal of Mathematics*, 2, 229–260. DOI: 10.1007/s11537-007-0657-8.
- D. Vardar, D. A. Buckley, B. S. Frank, and C. J. McKnight (1999). NMR structure of an F-actin-binding “headpiece” motif from villin. *Journal of Molecular Biology*, 294, 1299–1310. DOI: 10.1006/jmbi.1999.3321. BMRB entry 4428; BMRB DOI: 10.13018/BMR4428.
- D. S. Wishart, C. G. Bigam, A. Holm, R. S. Hodges, and B. D. Sykes (1995). ^1H , ^{13}C and ^{15}N random coil NMR chemical shifts of the common amino acids. I. Investigations of nearest-neighbor effects. *Journal of Biomolecular NMR*, 5, 67–81. DOI: 10.1007/BF00227471.
- M. P. Williamson (2013). Using chemical shift perturbation to characterise ligand binding. *Progress in Nuclear Magnetic Resonance Spectroscopy*, 73, 1–16. DOI: 10.1016/j.pnmrs.2013.02.001.
- C. J. McKnight, D. S. Doering, P. T. Matsudaira, and P. S. Kim (1996). A thermostable 35-residue subdomain within villin headpiece. *Journal of Molecular Biology*, 260, 126–134. DOI: 10.1006/jmbi.1996.0387.