

The BSD Conjecture as a Positivity Normal-Form: A Structural Survey

Lukas Geiger¹

¹*Independent Researcher, Bernau im Schwarzwald, Germany*

ORCID: [0009-0005-7296-1534](https://orcid.org/0009-0005-7296-1534)*

(Dated: May 1, 2026)

We reformulate the Birch and Swinnerton-Dyer conjecture as a *positivity normal-form*: a structural characterisation, not a proof. Four axiomatic constraints – finite arithmetic capacity, co-operative height positivity, controlled Iwasawa growth, and Euler-system composition coherence – jointly characterise the BSD leading-term formula as the unique “saturation profile”: no alternative leading-term structure can simultaneously satisfy all four axioms (conditional on these axioms being satisfied; for rank ≥ 2 , some axioms remain conjectural). The Néron-Tate height pairing provides the canonical positivity structure, playing the role for BSD that Frobenius positivity plays for the Weil conjectures. We show that the Gross-Zagier theorem is the rank-1 instantiation of this framework, coupling an analytic derivative to a non-negative height. For general rank r , we formulate the *Height Saturation Conjecture*: $L^{(r)}(E, 1)$ is proportional to the regulator $R_E = \det(\langle P_i, P_j \rangle) > 0$, forcing $\text{ord}_{s=1} L(E, s) = \text{rank } E(\mathbb{Q})$. The precise obstruction to a full proof is identified: the absence of a universal higher Gross-Zagier formula and a complete Kolyagin system for arbitrary rank.

I. INTRODUCTION

A. The BSD conjecture

Let $E/\mathbb{Q}$ be an elliptic curve with conductor N and L -function $L(E, s) = \sum_{n \geq 1} a_n n^{-s}$ (which exists for all $E/\mathbb{Q}$ by the modularity theorem [16]). The *Birch and Swinnerton-Dyer conjecture* (BSD) [1] asserts:

Conjecture I.1 (BSD – Weak form). $\text{ord}_{s=1} L(E, s) = \text{rank } E(\mathbb{Q})$.

Conjecture I.2 (BSD – Strong form). *The leading coefficient of $L(E, s)$ at $s = 1$ satisfies:*

$$\frac{L^{(r)}(E, 1)}{r!} = \frac{\Omega_E \cdot R_E \cdot \prod_p c_p \cdot |\text{Sha}(E/\mathbb{Q})|}{|E(\mathbb{Q})_{\text{tors}}|^2}, \quad (1)$$

where $r = \text{rank } E(\mathbb{Q})$, Ω_E is the real period, R_E is the regulator (determinant of the Néron-Tate height pairing), c_p are the Tamagawa numbers, and $\text{Sha}(E/\mathbb{Q})$ is the Tate-Shafarevich group. We write $E_{\text{tors}} := E(\mathbb{Q})_{\text{tors}}$ throughout for brevity.

The conjecture is known for:

- **Rank 0:** If $L(E, 1) \neq 0$, then $\text{rank } E(\mathbb{Q}) = 0$ and $|\text{Sha}| < \infty$ (Kolyagin [3], using Gross-Zagier [2]).
- **Rank 1:** If $\text{ord}_{s=1} L(E, s) = 1$, then $\text{rank } E(\mathbb{Q}) = 1$ and $|\text{Sha}| < \infty$ (Gross-Zagier + Kolyagin).
- **Rank ≥ 2 :** Open in general.

B. The positivity perspective

This paper proposes that the BSD conjecture, like the Hodge Conjecture and the Weil conjectures, is fundamentally a *positivity statement*. The key observation:

The Néron-Tate height pairing is a canonical positive-definite quadratic form on the Mordell-Weil group. It detects arithmetic “directions” – rational points modulo torsion – via their positive energy cost. The BSD conjecture asserts that the L -function “sees” exactly these directions, no more and no fewer.

The parallel to existing positivity proofs is:

Conjecture	Positive form	Result
Weil	Frobenius eigenvalues	Deligne 1974
Hodge	Hodge-Riemann form	Open
BSD	Néron-Tate height	Rank ≤ 1

The Gross-Zagier theorem [2] (extended to all Shimura curves by Yuan-Zhang-Zhang [25]) is the canonical example: it couples the first derivative $L'(E, 1)$ to the height $\hat{h}(P_K)$ of a Heegner point, establishing:

$$L'(E, 1) \neq 0 \iff \hat{h}(P_K) > 0 \iff P_K \text{ non-torsion}. \quad (2)$$

This is a *positivity identity*: an analytic object (derivative of L) equals a non-negative geometric object (height), and non-vanishing on one side forces non-triviality on the other.

C. Normal-form theorems

By a *positivity normal-form theorem* we mean a result of the following type: given a set of structural axioms

* AI tools (Claude by Anthropic) were used for mathematical formalization, literature verification, and editorial refinement. All mathematical content was developed, verified, and approved by the author.

(positivity, finiteness, composition coherence), there exists a *unique* functional relationship between the analytic and geometric invariants of the object in question, and this relationship is forced by the conjunction of the axioms. The Weil conjectures for the zeta function of a variety over $\mathbb{F}_q$ provide the archetype: Frobenius positivity (the Riemann hypothesis for varieties) forces the unique factorization of the zeta function via eigenvalues on étale cohomology.

D. Structure of this paper

We develop four axioms (Section III) whose conjunction forces the BSD leading-term formula as the unique “normal form.” We show how the rank ≤ 1 case is already proven within this framework (Section V), identify the precise obstructions for higher rank (Section VI), and discuss the relationship to Iwasawa theory and Euler systems (Section VII).

This paper is self-contained; all main results depend only on definitions and proofs within this document. Connections to the broader FST programme are cited for context but are not logically required.

II. THE POSITIVITY STRUCTURE OF BSD

A. The Néron-Tate height

Let $E/\mathbb{Q}$ be an elliptic curve. The *Néron-Tate (canonical) height* $\hat{h} : E(\mathbb{Q}) \rightarrow \mathbb{R}_{\geq 0}$ is the unique function satisfying:

1. $\hat{h}(P) \geq 0$ for all $P \in E(\mathbb{Q})$, with equality iff P is torsion.
2. $\hat{h}(nP) = n^2 \hat{h}(P)$ (quadratic scaling).
3. $\hat{h}$ differs from the naive (Weil) height by a bounded function.

The associated *Néron-Tate pairing* is:

$$\langle P, Q \rangle = \frac{1}{2} \left(\hat{h}(P + Q) - \hat{h}(P) - \hat{h}(Q) \right). \quad (3)$$

In particular, $\langle P, P \rangle = \hat{h}(P)$ for all $P \in E(\mathbb{Q})$.

Theorem II.1 (Néron-Tate positivity). *The pairing $\langle \cdot, \cdot \rangle$ is positive semi-definite on $E(\mathbb{Q})$ and positive definite on $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$.*

This is the BSD analog of the Hodge-Riemann bilinear form: a canonical, non-degenerate, positive-definite quadratic form on the “interesting” part of the arithmetic structure.

B. The regulator as positivity detector

Let $P_1, \dots, P_r$ be generators of $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$, where $r = \text{rank } E(\mathbb{Q})$. The *regulator* is:

$$R_E = \det(\langle P_i, P_j \rangle)_{1 \leq i, j \leq r}. \quad (4)$$

Proposition II.2 (Regulator positivity). *$R_E > 0$ whenever $r \geq 1$, and $R_E = 1$ by convention when $r = 0$.*

Proof. The matrix $(\langle P_i, P_j \rangle)$ is the Gram matrix of a positive-definite form. Its determinant is strictly positive. $\square$

The regulator is a *positivity certificate* for the existence of r independent rational points. It cannot vanish or be negative: positive-definiteness of $\hat{h}$ makes $R_E > 0$ an automatic consequence of $\text{rank } E(\mathbb{Q}) = r$.

C. The BSD formula as a positivity identity

The strong BSD formula (1) can be read as:

$$\underbrace{\frac{L^{(r)}(E, 1)}{r!}}_{\text{analytic}} = \underbrace{\Omega_E}_{>0} \cdot \underbrace{R_E}_{>0} \cdot \prod_{p \geq 1} c_p \cdot \underbrace{\frac{|\text{Sha}|}{|E_{\text{tors}}|^2}}_{>0}. \quad (5)$$

Every factor on the right is non-negative, and $R_E > 0$ whenever $r \geq 1$. The formula asserts that the analytic side – which has no a priori reason to be non-negative or to have the “right” order of vanishing – is *forced* to match this positive structure.

This is precisely the pattern of a *positivity normal-form theorem*: the analytic object is constrained by the positivity of the geometric/arithmetic object to which it is equal.

III. THE FOUR AXIOMS

We formulate four structural conditions that any coherent arithmetic theory of L -functions and Mordell-Weil groups must satisfy.

A. Axiom I: Finite Arithmetic Capacity

Axiom III.1 (Finite Arithmetic Capacity). *The Tate-Shafarevich group $\text{Sha}(E/\mathbb{Q})$ is finite:*

$$|\text{Sha}(E/\mathbb{Q})| < \infty. \quad (6)$$

Equivalently, the p^∞ -Selmer group has $\mathbb{Z}_p$ -corank equal to $\text{rank } E(\mathbb{Q})$ for every prime p . (The n -Selmer groups $\text{Sel}_n(E/\mathbb{Q})$ are always finite by a standard argument using finiteness of the class group and the weak Mordell-Weil theorem; the nontrivial content of this axiom is the finiteness of Sha .)

Status. Finiteness of Sha is known for rank ≤ 1 (Kolyagin [3]) and for certain higher-rank cases. It is expected to hold in general but remains unproven.

Role in the No-Go argument. If Sha were infinite, one could imagine a curve with $\text{rank } E(\mathbb{Q}) = 0$ but $\text{ord}_{s=1} L(E, s) = 2$: the “missing rank” would be hidden in Sha. Axiom I excludes this loophole.

B. Axiom II: Cooperative Height Positivity

Axiom III.2 (Cooperative Height Positivity). *The Néron-Tate height pairing is positive-definite on $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$:*

$$\langle P, P \rangle > 0 \quad \forall P \in E(\mathbb{Q}) \setminus E(\mathbb{Q})_{\text{tors}}. \quad (7)$$

Moreover, the height pairing is “cooperative”: heights of independent points combine quadratically via the Gram determinant $R_E = \det(\langle P_i, P_j \rangle) > 0$.

Status. Unconditionally true (a theorem, not a conjecture).

Role in the No-Go argument. Positivity ensures that the right-hand side of BSD has a definite sign structure: $R_E > 0$ is forced. Any analytic formula for $L^{(r)}(E, 1)$ must be compatible with this positivity.

C. Axiom III: Controlled Iwasawa Growth

Axiom III.3 (Controlled Iwasawa Growth). *In the cyclotomic $\mathbb{Z}_p$ -extension $\mathbb{Q}_\infty/\mathbb{Q}$, the p -primary Selmer groups grow in a controlled, asymptotically predictable fashion. Writing $X_n = \text{Sel}_{p^\infty}(E/\mathbb{Q}_n)^\vee$ for the Pontryagin dual (a finitely generated $\mathbb{Z}_p$ -module whose $\mathbb{Z}_p$ -rank equals $\text{rank } E(\mathbb{Q}_n)$):*

$$\text{ord}_p(|X_n^{\text{tors}}|) = \mu p^n + \lambda n + \nu + O(1), \quad (8)$$

where X_n^{tors} denotes the $\mathbb{Z}_p$ -torsion submodule of X_n , and $\mu, \lambda \geq 0$ are the Iwasawa invariants. The Iwasawa main conjecture provides a p -adic L -function $L_p(E, s)$ whose characteristic ideal matches the Selmer module.

Status. The Iwasawa main conjecture for elliptic curves has been established in important cases, with different results covering complementary aspects:

- **Kato [4]:** Proves one divisibility direction ($\text{char}_\Lambda(\text{Sel}^\vee) \subseteq (\mathcal{L}_p)$, i.e., the characteristic ideal of the Selmer dual is contained in the ideal generated by the p -adic L -function) for all primes p at which the modular form is ordinary and the residual representation satisfies mild conditions. This holds without restriction on the rank and provides an upper bound on the Selmer module.
- **Skinner-Urban [6]:** Prove the full main conjecture (both divisibilities, hence equality) for *ordinary* primes p (i.e., $p \nmid a_p$), under technical conditions including that $\bar{\rho}_{E,p}$ is irreducible and that

the residual representation satisfies certain local hypotheses. This covers a large class of elliptic curves at ordinary primes, but does not apply at supersingular primes.

For supersingular primes, the main conjecture requires a different formulation (Pollack, Kobayashi, Sprung) and is the subject of ongoing work. The full main conjecture for all primes and all elliptic curves remains open.

Role in the No-Go argument. The controlled growth formula (8) (note: “controlled” refers to the asymptotically predictable growth governed by μ and λ , not strict monotonicity, since the $O(1)$ error allows bounded fluctuations) prevents the analytic rank from exhibiting unbounded “oscillations” between layers of the $\mathbb{Z}_p$ -tower. Together with the main conjecture, it couples the p -adic L -function to the Selmer module, providing a p -adic positivity constraint.

D. Axiom IV: Euler-System Composition Coherence

Axiom III.4 (Euler-System Composition Coherence). *There exists an Euler system $\{c_m\}_{m \in \mathcal{M}}$ – a coherent family of cohomology classes indexed by moduli m – satisfying norm-compatibility relations:*

$$\text{cores}_{m\ell/m}(c_{m\ell}) = P_\ell(\text{Frob}_\ell^{-1}) c_m, \quad (9)$$

where P_ℓ is the Euler factor at ℓ and cores is the corestriction. The Euler system provides upper bounds on Selmer groups through the Kolyagin derivative machinery:

$$|\text{Sel}(E/\mathbb{Q})| \leq C \cdot |(\text{image of Euler system})|, \quad (10)$$

for a computable constant C .

Status. Euler systems exist for many elliptic curves:

- **Heegner points** (for rank 1): the classical Euler system [3].
- **Kato’s Euler system:** from K -theory of modular curves [4].
- **Beilinson-Flach elements:** for Rankin-Selberg convolutions [13].

Role in the No-Go argument. The Euler system is the “exclusion machine”: it bounds the Selmer group from above, excluding the possibility of hidden rank in Sha or in unexpected Selmer classes. Combined with positivity (Axiom II), it forces the analytic and algebraic ranks to agree.

Remark III.5 (Root number and parity). The functional equation of $L(E, s)$ determines the *root number* $w(E) = \pm 1$, which constrains the parity of $\text{ord}_{s=1} L(E, s)$: if $w(E) = +1$, then $\text{ord}_{s=1} L(E, s)$ is even; if $w(E) = -1$, it is odd. The *parity conjecture* – that $(-1)^{\text{rank } E(\mathbb{Q})} = w(E)$ – is proven in many cases (Nekovář [23]; T. and

V. Dokchitser [24]). While this parity constraint is not one of our four axioms, it provides an additional consistency check: any “alternative profile” violating the parity conjecture would be excluded by the functional equation alone.

Remark III.6 (Independence of the axioms). For rank ≤ 1 , Axioms I–IV are independently established theorems (Kolyvagin, Néron-Tate, Kato/Skinner-Urban, Heegner points). For rank ≥ 2 , Axioms I and IV are *not* independent of BSD: finiteness of Sha (Axiom I) is itself a consequence of the full BSD conjecture, and the existence of a complete Kolyvagin system of sufficient depth (Axiom IV) is not established. The conditional result (Proposition IV.1) should therefore be read as a *structural characterisation*: it identifies the BSD formula as the unique normal form compatible with the axioms, rather than providing an independent proof of BSD.

E. Heuristic parallels

The four axioms exhibit structural parallels to the variational framework of [18]. These parallels are *purely heuristic and motivational* – they suggest why the axiom structure is natural, but do not constitute rigorous mathematical connections and are not used in any proof.

- **Axiom I $\leftrightarrow$ Finite geometric capacity (Axiom A).** The finiteness of Sha is the arithmetic analog of finite capacity: the “hidden mass” is bounded, preventing infinite discrepancies from being absorbed into the right-hand side of the BSD formula.
- **Axiom II $\leftrightarrow$ Cooperative reinforcement (Axiom B).** Each independent rational point costs positive energy (height). The quadratic scaling $\hat{h}(nP) = n^2 \hat{h}(P)$ mirrors cooperative growth in the saturation ODE.
- **Axiom III $\leftrightarrow$ Controlled growth (Axiom C).** As one ascends the $\mathbb{Z}_p$ -tower, arithmetic complexity grows in a controlled, predictable fashion – no oscillations or jumps in the Selmer rank occur.
- **Axiom IV $\leftrightarrow$ Composition law (Axiom D).** Norm-compatible steps in the Euler system compose coherently. The norm-compatibility (9) ensures that the Euler system is a globally consistent family, not a collection of ad hoc local constructions.

IV. THE NORMAL-FORM THEOREM

A. Statement

Proposition IV.1 (BSD Normal-Form Characterisation – Conditional Structural Result). (*Conditional on*

Axioms I–IV.) *The following conclusions hold only under the assumption that Axioms I–IV are satisfied. For rank ≤ 1 , this assumption is fulfilled by established theorems; for rank ≥ 2 , Axioms I and IV themselves depend on unproven conjectures (see Remark III.6). This is a structural characterisation, not a new proof of BSD.*

Let $E/\mathbb{Q}$ be an elliptic curve satisfying Axioms I–IV. Then:

1. The analytic rank equals the algebraic rank:

$$\text{ord}_{s=1} L(E, s) = \text{rank } E(\mathbb{Q}). \quad (11)$$

2. The leading term has the normal form:

$$\frac{L^{(r)}(E, 1)}{r!} = \Omega_E \cdot R_E \cdot \prod_p c_p \cdot \frac{|\text{Sha}(E/\mathbb{Q})|}{|E(\mathbb{Q})_{\text{tors}}|^2}, \quad (12)$$

with all factors on the right determined by the arithmetic of E .

3. No alternative leading-term profile is compatible with Axioms I–IV.

Remark IV.2 (Conditional status and logical circularity at rank ≥ 2). Proposition IV.1 is a *conditional structural characterisation*: it shows that the BSD formula is the unique normal form *given* Axioms I–IV. **It does not provide an unconditional proof of BSD, and for rank ≥ 2 the argument is logically circular.** For rank ≤ 1 , all four axioms are independently established theorems (Kolyvagin, Néron-Tate, Kato/Skinner-Urban, Heegner points), so the conclusion holds unconditionally and constitutes a genuine structural reformulation of known results. For rank ≥ 2 , however, Axiom I (finiteness of Sha) is itself a consequence of the full BSD conjecture, and Axiom IV (existence of a complete Kolyvagin system of sufficient depth) is not established independently of BSD. The proposition therefore has the logical structure “BSD + known facts $\Rightarrow$ BSD is the unique normal form” at rank ≥ 2 , which is logically valid but mathematically trivial as a proof strategy. Its value lies in *structural analysis*: identifying the BSD formula as the unique output of the four axioms clarifies *which* components are needed for a proof, even though it does not provide one.

The analogy to the Weil conjectures requires qualification: Grothendieck’s reformulation via étale cohomology was *not* circular – étale cohomology was an independently constructed mathematical object, and the reformulation proved rationality and the functional equation outright, reducing the problem to the Riemann hypothesis for varieties. By contrast, our axioms are not independently established for rank ≥ 2 . The analogy holds *in spirit* (identifying the correct structural framework) but not *in logical force*.

B. Proof structure

The proof strategy has three components, mirroring the saturation theorem:

Step 1: Positivity forces the sign. By Axiom II, $R_E > 0$ whenever $r \geq 1$. By Axiom I, $|\text{Sha}| < \infty$, so the right-hand side of the BSD formula is a positive real number (for $r \geq 1$) or equals $L(E, 1)/\Omega_E$ times arithmetic factors (for $r = 0$). The analytic side must match this sign.

Step 2: Composition forces the order. [Status: open for rank ≥ 2 ; proven for rank ≤ 1 .] For rank 0, Kato's Euler system [4] shows that $L(E, 1) \neq 0$ implies $\text{rank } E(\mathbb{Q}) = 0$. For rank 1, Kolyvagin's Euler system [3] shows that if $\text{ord}_{s=1} L(E, s) = 1$ (and the Heegner point is non-torsion), then $\text{rank } E(\mathbb{Q}) = 1$ and $|\text{Sha}| < \infty$. **For rank ≥ 2 , the general inequality $\text{rank } E(\mathbb{Q}) \leq \text{ord}_{s=1} L(E, s)$ is not established** – it is equivalent to one direction of the BSD conjecture and requires the unproven hypothesis (H2). The reverse inequality $\text{ord}_{s=1} L(E, s) \leq \text{rank } E(\mathbb{Q})$ requires the height coupling (H1): if $L^{(r)}(E, 1)$ is proportional to $R_E > 0$ (Axiom II), then L cannot vanish to order higher than $r = \text{rank } E(\mathbb{Q})$ without contradicting the positivity of the regulator. *Note: both directions of this step are conditional on unproven hypotheses for rank ≥ 2 (see Remark below).*

Remark IV.3 (Heuristic nature of the reverse inequality in Step 2). The direction $\text{ord}_{s=1} L(E, s) \leq \text{rank } E(\mathbb{Q})$ relies on the full BSD leading-term formula as an assumption: without the BSD formula, the argument that “the leading term vanishes faster than the regulator, violating positivity” remains heuristic. Step 2 should therefore be understood as conditional on the BSD formula, not as a self-contained deduction from Axioms I–IV alone (cf. Remark IV.2 for the full discussion of logical dependencies).

Step 3: Uniqueness of the normal form. Given the rank equality, the BSD formula is the unique expression compatible with:

- The functional equation of $L(E, s)$ (determines Ω_E).
- The Birch lemma and its generalizations (determines the local factors c_p).
- The regulator R_E (determined by the height pairing).
- Finiteness of Sha (Axiom I).

C. Uniqueness of the leading-term profile

Proposition IV.4 (Uniqueness of the leading-term formula – Proof sketch). *Let $E/\mathbb{Q}$ be an elliptic curve with $\text{rank } E(\mathbb{Q}) = r$, satisfying Axioms I–IV and the height*

coupling hypothesis (H1). Then the BSD leading-term formula (1) is the unique expression

$$\frac{L^{(r)}(E, 1)}{r!} = \Phi(E)$$

satisfying the following five constraints simultaneously:

- (U1) **Analytic-algebraic rank equality:** $\text{ord}_{s=1} L(E, s) = \text{rank } E(\mathbb{Q})$.
- (U2) **Positivity of leading coefficient:** $\Phi(E) > 0$ whenever $r \geq 1$ (forced by Axiom II, since any coupling to $R_E > 0$ must preserve the sign).
- (U3) **Integrality of the Sha contribution:** the ratio $\Phi(E)/(\Omega_E \cdot R_E \cdot \prod_p c_p)$ is of the form $|\text{Sha}|/|E(\mathbb{Q})_{\text{tors}}|^2$ with $|\text{Sha}| \in \mathbb{Z}_{>0}$ (forced by Axiom I).
- (U4) **Compatibility with functional equation:** $\Phi(E)$ is compatible with the p -adic interpolation formula $L_p(E, 0) \sim L(E, 1)/\Omega_E^+$ and its higher derivatives (forced by Axiom III, the Iwasawa main conjecture specialization).
- (U5) **Isogeny invariance:** $\Phi(E)$ depends only on the isogeny class of E , not on the choice of Weierstrass model or basis of $E(\mathbb{Q})$.

Heuristic argument (not a rigorous proof). We argue heuristically that (U1)–(U5) jointly determine $\Phi(E)$. **Caveat:** This argument is a plausibility sketch, not a rigorous proof. The rigorous framework for the uniqueness question is the *Bloch-Kato conjecture* (Tamagawa number conjecture) [33, 34], which defines Tamagawa numbers for motives and derives the BSD formula as the special case for the motive $h^1(E)(1)$, eliminating the rational ambiguity of the Beilinson conjectures. Our heuristic argument should be read as motivation for *why* the BSD formula is the expected answer; the Bloch-Kato framework provides the correct mathematical setting for making this precise.

Step 1: The positivity factor must be R_E . By (U2), $\Phi(E) > 0$ for $r \geq 1$, so $\Phi(E)$ must involve a positive-definite invariant of $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$. The only canonical such invariant is the Gram determinant $R_E = \det(\langle P_i, P_j \rangle)$ of the Néron-Tate height, which is the *unique* positive-definite bilinear form on $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$ satisfying quadratic scaling $\hat{h}(nP) = n^2 \hat{h}(P)$, agreement with the Weil height up to $O(1)$, and positive-definiteness modulo torsion (cf. [15], Theorem VIII.9.3). *Note:* This step only shows that *if* $\Phi(E)$ contains a positive-definite factor, it must be R_E ; it does not show that $\Phi(E)$ must contain such a factor at all. The latter requires the height coupling hypothesis (H1).

Step 2: The archimedean and local factors are determined. (U4) (compatibility with the p -adic interpolation formula and functional equation) determines the archimedean factor Ω_E and the local factors c_p through

the Birch lemma and the specialization of the Iwasawa main conjecture.

Step 3: The remaining factor is constrained by integrality. After extracting $\Omega_E \cdot R_E \cdot \prod_p c_p$, (U3) requires the remaining ratio $\Phi(E)/(\Omega_E \cdot R_E \cdot \prod_p c_p)$ to equal $|\text{Sha}(E/\mathbb{Q})|/|E(\mathbb{Q})_{\text{tors}}|^2$ with $|\text{Sha}| \in \mathbb{Z}_{>0}$. *Note:* This step presupposes that the decomposition $\Phi(E) = \Omega_E \cdot R_E \cdot (\text{rest})$ is valid, which is precisely the content of the BSD formula. The “integrality constraint” does not independently derive this decomposition; rather, it is *consistent with* the BSD formula once assumed.

Step 4: Isogeny invariance excludes ad hoc constructions. (U5) ensures that $\Phi(E)$ depends only on the isogeny class, not on auxiliary choices (basis of $E(\mathbb{Q})$, Weierstrass model). The individual quantities Ω_E , R_E , c_p , $|\text{Sha}|$, $|E(\mathbb{Q})_{\text{tors}}|$ are *not* separately isogeny-invariant, but their combination in the BSD formula is, since $L(E, s)$ depends only on the isogeny class (Faltings). This forces $\Phi(E)$ to be the unique isogeny-invariant product of canonical arithmetic data.

In summary, the five constraints make the BSD formula the unique *natural candidate* of the form “product of canonical arithmetic invariants.” This is a weaker claim than a rigorous uniqueness proof: it shows that the BSD formula is the unique expression *within the class of products of known canonical invariants*, not that it is the unique formula satisfying (U1)–(U5) in the space of all possible functions of E . The Bloch-Kato conjecture [33] provides the rigorous motivic framework in which this uniqueness can be made precise. $\square$

D. Exclusion of alternatives

We show that “alternative profiles” – hypothetical scenarios where the BSD formula fails – violate the axioms:

Proposition IV.5 (Consistency check: scenarios excluded by the axioms). *The following scenarios are excluded by the axioms. Note: For $\text{rank} \leq 1$, the exclusions restate known results (Kato, Kolyagin) in the language of our axiom system. For $\text{rank} \geq 2$, only (E3) and (E4) are excluded, and both follow trivially from the axiom assumptions. The non-trivial cases (E1) and (E2) remain conditional for $\text{rank} \geq 2$.*

(E1) Analytic zero without geometric direction: $\text{ord}_{s=1} L(E, s) = r > \text{rank } E(\mathbb{Q})$. *Note that Euler-system bounds (Axiom IV) establish the opposite inequality $\text{rank } E(\mathbb{Q}) \leq \text{ord}_{s=1} L(E, s)$, so they do not directly exclude this scenario. Rather, exclusion of (E1) requires Axiom I ($|\text{Sha}| < \infty$) combined with the height coupling hypothesis (H1): if the coupling $L^{(r)}(E, 1) = C_r \cdot R_E$ holds (where $r = \text{rank } E(\mathbb{Q})$), then $\text{ord} > r$ would force $L^{(r)}(E, 1) = 0$ (since L vanishes to order $> r$), while $R_E > 0$ (Axiom II) and $C_r > 0$, yielding a contradiction. For $\text{rank} \leq 1$, Gross-Zagier provides this coupling*

explicitly. Note: For $\text{rank} \geq 2$, this exclusion is conditional on (H1) and (H2).

(E2) Geometric rank without analytic derivative: $\text{rank } E(\mathbb{Q}) = r > \text{ord}_{s=1} L(E, s) = s$. *Under the assumption of the leading-term formula and the Euler-system bound (H2), this is excluded: (H2) gives $r = \text{rank } E(\mathbb{Q}) \leq \text{ord}_{s=1} L(E, s) = s$, directly contradicting $r > s$. Thus for $\text{rank} \leq 1$ (where Axiom IV is proven unconditionally via Kato’s Euler system), (E2) is excluded outright. **For $\text{rank} \geq 2$, exclusion is conditional:** it depends on the full Euler-system bound (H2), which itself requires the existence of a Kolyagin system of sufficient depth – a major open problem. Without (H2), the inequality $\text{rank} \leq \text{ord}$ is not established, and no contradiction arises; thus (E2) cannot currently be excluded for $\text{rank} \geq 2$. Supplementarily, under the assumption that the leading-term formula holds, the height coupling (H1) provides a compatible perspective: it asserts that $L^{(r)}(E, 1)/r!$ equals $C_r \cdot R_E$ as a leading-term identity, coupling the r -th derivative to the regulator precisely when $\text{ord}_{s=1} L(E, s) = r$; if $\text{ord} = s < r$, the hypothesis of the coupling formula is not met.*

(E3) Infinite Sha absorbing discrepancy: $\text{ord}_{s=1} L(E, s) = r$ but $|\text{Sha}| = \infty$, allowing a different right-hand side. *Excluded by Axiom I.*

(E4) Oscillating rank in $\mathbb{Z}_p$ -towers: Rank jumps non-monotonically through cyclotomic layers. *Excluded by Axiom III.*

V. RANK ≤ 1 : POSITIVITY IN ACTION

A. Rank 0: Kato’s exclusion

Proposition V.1 (Rank 0 saturation). *Let $E/\mathbb{Q}$ with $L(E, 1) \neq 0$. Then all four axioms are verified, $\text{rank } E(\mathbb{Q}) = 0$, $|\text{Sha}| < \infty$, and the BSD formula holds.*

Proof sketch. Axiom II applied: If $\text{rank } E(\mathbb{Q}) \geq 1$, there exists a non-torsion point P with $\hat{h}(P) > 0$. However, the direct implication $L(E, 1) \neq 0 \Rightarrow \text{rank} = 0$ is not established via Axiom II alone; rather, it follows from the Euler-system machinery (Axiom IV) below.

Axiom IV applied: Kato’s Euler system [4], constructed from K -theory classes on modular curves, provides the relevant upper bound for rank 0. Specifically, for any prime p at which the residual representation $\bar{\rho}_{E,p}$ is surjective (which holds for all but finitely many p), Kato shows that if $L(E, 1) \neq 0$, the image of his Euler system class in $H^1(\mathbb{Q}, T_p(E))$ is non-trivial, which forces the p^∞ -Selmer group to have $\mathbb{Z}_p$ -corank zero:

$$\text{corank}_{\mathbb{Z}_p} \text{Sel}_{p^\infty}(E/\mathbb{Q}) = 0, \quad (13)$$

hence $\text{rank } E(\mathbb{Q}) = 0$. Combined with the Selmer bound, this also yields $|\text{Sha}(E/\mathbb{Q})| < \infty$. (Note: for rank 0, the Heegner-point Euler system is not directly applicable since the Heegner point is torsion when $L(E, 1) \neq 0$; Kato’s Euler system is the appropriate tool.)

The rank-0 case is fully “saturated” in the positivity framework: all four axioms are verified (Axiom III at all primes of good ordinary reduction with surjective residual representation; the remaining primes are handled by complementary results or are finite in number), and the No-Go argument excludes $\text{rank} \geq 1$. $\square$

B. Rank 1: Gross-Zagier as a positivity identity

The Gross-Zagier theorem [2] states: for a suitable imaginary quadratic field K satisfying the Heegner hypothesis (every prime dividing the conductor N splits in K – such a K always exists by the Chinese Remainder Theorem and quadratic reciprocity),

$$L'(E, 1) = \frac{8\pi^2 \|f\|^2}{\sqrt{|D_K|}} \cdot \hat{h}(P_K), \quad (14)$$

where f is the newform associated to E , $\|f\|^2 = \int_{\Gamma_0(N) \backslash \mathcal{H}} |f(z)|^2 \frac{dx dy}{y^2}$ is the Petersson norm (the L^2 -norm of f with respect to the hyperbolic measure on the modular curve), D_K is the discriminant, and $P_K \in E(K)$ is the Heegner point. (The formula is stated up to an explicit nonzero rational factor depending on K ; for $|D_K| > 4$ this factor equals 1. The relationship between the Petersson norm $\|f\|^2$ and the Néron period Ω_E in the BSD formula involves the Manin constant c_M ; for the optimal curve in each isogeny class, $c_M = 1$ is verified computationally for all conductors up to 500,000 in the Cremona database [22]¹ (unconditionally for $N \leq 130,000$; conditionally on optimality of the first curve for $N > 400,000$). More generally, Česnavičius [32] proved $c_M = 1$ for all semistable elliptic curves over $\mathbb{Q}$.)

Proposition V.2 (Gross-Zagier as positivity identity). *The Gross-Zagier formula (14) is a positivity identity:*

- The left side $L'(E, 1)$ has no a priori sign constraint.
- The right side is a product of $\hat{h}(P_K) \geq 0$ (by Axiom II) and positive constants.
- Non-vanishing: $L'(E, 1) \neq 0 \iff \hat{h}(P_K) > 0 \iff P_K$ is non-torsion.

This is the BSD analog of the Hodge-Riemann bilinear relation: the analytic object is equal to a geometrically

positive quantity, and the positivity forces the correct rank.

Kolyvagin’s Euler system then completes the argument: it shows $\text{rank } E(\mathbb{Q}) = 1$ (the Heegner point generates a finite-index subgroup) and $|\text{Sha}| < \infty$.

Remark V.3. The rank-1 proof follows the exact pattern of a No-Go theorem:

1. Assume $\text{ord}_{s=1} L(E, s) = 1$.
2. Gross-Zagier converts the analytic datum to a positive geometric datum ($\hat{h}(P_K) > 0$).
3. Kolyvagin excludes alternative explanations ($\text{rank} > 1$ or $|\text{Sha}| = \infty$).
4. Conclusion: $\text{rank } E(\mathbb{Q}) = 1$, and the BSD formula holds.

No explicit construction of rational points is needed beyond the Heegner point (which serves as a “seed” for the Euler system, not as a constructive proof of rank).

C. Numerical illustration

Example V.4 (Curve 11a – rank 0). The elliptic curve $E : y^2 + y = x^3 - x^2 - 10x - 20$ (Cremona label 11a1, conductor $N = 11$; [22]) is the curve of smallest conductor. It has $\text{rank } E(\mathbb{Q}) = 0$, $E(\mathbb{Q})_{\text{tors}} \cong \mathbb{Z}/5\mathbb{Z}$, and $|\text{Sha}| = 1$ (proven). With $R_E = 1$ (by convention for rank 0), $\Omega_E = 1.2692\dots$, $c_{11} = 5$ (Kodaira type I_5 , split multiplicative reduction), and $|E(\mathbb{Q})_{\text{tors}}| = 5$:

$$L(E, 1) = \Omega_E \cdot R_E \cdot c_{11} \cdot \frac{|\text{Sha}|}{|E_{\text{tors}}|^2} = 1.2692 \cdot 1 \cdot 5 \cdot \frac{1}{25} = 0.2538\dots$$

This matches the numerical value $L(E, 1) = 0.2538\dots$ to all available precision. All four axioms are verified: $|\text{Sha}| = 1$ (I), height positivity vacuously satisfied for $r = 0$ (II), Iwasawa main conjecture proven at all primes $p \neq 11$ (III), Kato’s Euler system provides the Selmer bound (IV).

The positivity interpretation: rank 0 is the “vacuum state” of the positivity framework – no height direction exists, $R_E = 1$ by convention, and the BSD formula reduces to a ratio of archimedean and local data. The positivity structure is trivially saturated.

Example V.5 (Curve 37a – rank 1). The elliptic curve $E : y^2 + y = x^3 - x$ (Cremona label 37a1, conductor $N = 37$) has $\text{rank } E(\mathbb{Q}) = 1$, generated by $P = (0, 0)$ with $\hat{h}(P) = 0.0511\dots$. The BSD data: $\Omega_E = 5.9869\dots$, $R_E = 0.0511\dots$, $c_{37} = 1$, $|E(\mathbb{Q})_{\text{tors}}| = 1$, $|\text{Sha}| = 1$ (proven). The BSD formula predicts $L'(E, 1) = \Omega_E \cdot R_E \cdot 1 \cdot 1/1 = 0.3059\dots$, which matches the numerical computation $L'(E, 1) = 0.3059\dots$ to all available digits. All four axioms are verified: $|\text{Sha}| = 1 < \infty$ (I), $\hat{h}(P) > 0$ (II), Iwasawa main conjecture proven for $p \neq 37$ (III), Heegner point Euler system (IV).

¹ Cremona’s `ecdata` provides complete tables up to conductor $N = 500,000$; see <https://johncremona.github.io/ecdata/>.

Example V.6 (Curve 389a – rank 2). The elliptic curve $E : y^2 + y = x^3 + x^2 - 2x$ (Cremona label 389a1) has rank $E(\mathbb{Q}) = 2$, generated by $P_1 = (-1, 1)$ and $P_2 = (0, 0)$.² The height matrix is:

$$(\langle P_i, P_j \rangle) = \begin{pmatrix} 0.6868 & 0.2685 \\ 0.2685 & 0.3270 \end{pmatrix}, \quad R_E = \det = 0.1525 \dots$$

The BSD formula predicts $L''(E, 1)/2 = \Omega_E \cdot R_E \cdot \prod c_p \cdot |\text{Sha}|/|E_{\text{tors}}|^2$. With $\Omega_E = 4.9804 \dots$, $c_{389} = 1$, $|E_{\text{tors}}| = 1$, and $|\text{Sha}| = 1$ (numerically verified), this gives $L''(E, 1)/2 = 0.7593 \dots$, matching numerical computation. This is a rank-2 case where Axioms I and IV are not rigorously proven but numerically consistent.

VI. HIGHER RANK: THE OPEN FRONTIER

A. The Height Saturation Conjecture

For general rank r , we formulate:

Conjecture VI.1 (Height Saturation – Reformulation of BSD in positivity language). *Let $E/\mathbb{Q}$ with rank $E(\mathbb{Q}) = r$. Let $P_1, \dots, P_r$ be a basis of $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$. Then:*

$$L^{(r)}(E, 1) \sim R_E = \det(\langle P_i, P_j \rangle) > 0, \quad (15)$$

in the sense that the ratio $L^{(r)}(E, 1)/(r! \cdot \Omega_E \cdot R_E \cdot \prod c_p)$ equals $|\text{Sha}|/|E_{\text{tors}}|^2$.

Remark VI.2 (Relation to the classical BSD conjecture). Conjecture VI.1 is *equivalent* to the strong BSD conjecture (Conjecture I.2): the “height saturation” formulation simply rephrases the BSD leading-term formula in the language of this paper’s positivity framework. The name “Height Saturation” emphasizes the role of the positive-definite regulator $R_E > 0$ as the dominant geometric factor, but the mathematical content is identical to the classical formulation. We retain this reformulation because it highlights the positivity structure, not because it adds new mathematical content.

This is the “higher Gross-Zagier” in its most optimistic form: the r -th derivative of L is proportional to the $r \times r$ regulator – the determinant of a positive-definite matrix.

Remark VI.3 (Disc-averaging formulation). The BSD leading-term formula can be expressed as a contour-integral limit, making the role of the Taylor coefficient explicit. This is a standard complex-analytic identity, not an independent conjecture. For an elliptic curve $E/\mathbb{Q}$ of analytic rank $r = \text{ord}_{s=1} L(E, s)$:

$$\frac{L^{(r)}(E, 1)}{r!} = \lim_{\varepsilon \rightarrow 0} \varepsilon^{-r} \cdot \frac{1}{2\pi i} \oint_{|s-1|=\varepsilon} L(E, s) ds = \Omega_E \cdot \det(\langle P_i, P_j \rangle_{\text{NT}}) \cdot \frac{\# \text{Sha}(E) \cdot \prod c_p}{\# E(\mathbb{Q})_{\text{tors}}^2},$$

² The height matrix depends on the choice of generators; the regulator R_E is invariant under change of $\mathbb{Z}$ -basis of $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$ (since $\det(A^T M A) = (\det A)^2 \det M$ and $\det A = \pm 1$ for $A \in \text{GL}(r, \mathbb{Z})$).

where $P_1, \dots, P_r$ are generators of $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$.

Remark VI.4 (Normalisation of the disc average). The disc-averaging formula in Remark VI.3 should be understood with the standard complex-analytic normalisation: $\frac{1}{2\pi i} \oint_{|s-1|=\varepsilon} L(E, s) ds = \frac{L^{(r)}(E, 1)}{r!} \varepsilon^r + O(\varepsilon^{r+1})$ as $\varepsilon \rightarrow 0$. The height saturation ratio is therefore $\lim_{\varepsilon \rightarrow 0} \varepsilon^{-r} \cdot \frac{1}{2\pi i} \oint_{|s-1|=\varepsilon} L(E, s) ds = \frac{L^{(r)}(E, 1)}{r!}$, which by BSD equals $\Omega_E \cdot R_E \cdot \prod c_p \cdot |\text{Sha}|/|E_{\text{tors}}|^2$.

Remark VI.5 (Height saturation as a selection principle). The conjecture reformulates the BSD leading-term formula as a *limit identity* rather than a pointwise evaluation: the contour average extracts the leading Taylor coefficient, which BSD equates to a product of arithmetic invariants dominated by the positive-definite regulator R_E . This perspective suggests viewing the BSD formula as a *selection principle*: among all leading-term structures of the form “product of canonical arithmetic invariants” (see the heuristic argument of Proposition IV.4), the positivity of the Néron-Tate height pairing makes the BSD formula the unique natural candidate, with Sha providing the remaining arithmetic correction. The rigorous formulation of this uniqueness is provided by the Bloch-Kato conjecture [33].

Remark VI.6 (Numerical test via LMFDB). The Height Saturation Conjecture is amenable to direct numerical verification using the LMFDB database. For elliptic curves $E/\mathbb{Q}$ of rank 2–4, one can compute the discretised Néron-Tate height $\hat{h}_{\text{disc}}(E, D)$ for quadratic twists E_D with $|D| \leq 10^7$ and test whether $\hat{h}_{\text{disc}}(E, D) \cdot R(E)^{-1} \rightarrow 1$ statistically. This replaces the missing algebraic control (Axiom IV for rank ≥ 2) with *typical validity*—a thermodynamic rather than arithmetic statement. Smith’s distribution results for 2^∞ -Selmer groups [29] provide the probabilistic framework for such tests.

Remark VI.7 (Rank-2 height saturation). For 17 rank-2 elliptic curves from the Cremona database (conductors $389 \leq N \leq 5077$), the regulator satisfies $R > 0$ in all cases ($R_{\min} = 0.095$, $R_{\max} = 1.700$). Log-log regression yields $R \sim N^{0.099}$, consistent with sublinear growth. **Caveat:** The sample size ($n = 17$) is too small for statistically significant conclusions; these results should be viewed as preliminary consistency checks, not as statistical confirmation of a scaling law. The positivity $R > 0$ is guaranteed a priori by the positive-definiteness of the Néron-Tate height (Theorem II.1). Script: `compute_rank2_lmfdb.py`.

What would a proof require?

A proof of Conjecture VI.1 via the positivity framework requires three hypotheses, which we label for reference throughout the paper:

(H1) Higher Gross-Zagier formula. An explicit identity:

$$L^{(r)}(E, 1) = C_r \cdot \det(\langle P_i, P_j \rangle), \quad (16)$$

where $P_1, \dots, P_r$ are canonical arithmetic points (“higher Heegner points”) and $C_r > 0$ is an explicit constant.

Status: For $r = 1$, this is Gross-Zagier. For $r = 2$, partial results exist via diagonal restriction of p -adic families [8]. Kim [5] has obtained a higher Gross-Zagier formula refining the connection between Heegner point Kolyvagin systems and Selmer group structure. For general r , this is a major open problem. Recent work on Euler systems for higher-rank motives [14] provides progress but not a complete formula.

(H2) Higher Kolyvagin system. An Euler system machinery that bounds $\text{Sel}(E/\mathbb{Q})$ using r classes simultaneously, establishing:

$$\text{rank } E(\mathbb{Q}) \leq \text{ord}_{s=1} L(E, s). \quad (17)$$

Status: For $r = 1$, this is Kolyvagin’s method. For general r , Mazur-Rubin [7] have developed the theory of Kolyvagin systems abstractly, and Burns-Sano [11] and Burns-Sakamoto-Sano [12] have extended this to higher rank Euler, Kolyvagin, and Stark systems over general coefficient rings. Kim [5] has obtained a Kolyvagin system-theoretic refinement of the Gross-Zagier formula. However, the *existence* of a Kolyvagin system of sufficient depth for arbitrary rank is not established.

(H3) Sha finiteness for general rank. Axiom I must hold unconditionally:

$$|\text{Sha}(E/\mathbb{Q})| < \infty \quad \forall E/\mathbb{Q}. \quad (18)$$

Status: Open for rank ≥ 2 . This is widely expected but no general proof exists.

C. The positivity gap

The situation can be summarized:

Axiom	Rank 0	Rank 1	Rank ≥ 2
I (Finite Sha)	✓	✓	Open
II (Height pos.)	✓	✓	✓
III (Iwasawa)	✓	✓	Partial
IV (Euler sys.)	✓	✓	Open

The “positivity” (Axiom II) is always available – it is a theorem. The obstruction lies in the “composition” (Axioms I, III, IV): the Euler-system/Kolyvagin/Iwasawa machinery is not yet fully developed for higher rank.

In summary: the positivity (Axiom II) is established, but the composition law (Axioms I, III, IV) is incomplete for rank ≥ 2 . The BSD formula is forced once the Euler-system composition is established for arbitrary rank.³

VII. IWASAWA THEORY AS COMPOSITION COHERENCE

A. The p -adic perspective

Iwasawa theory provides the clearest realization of Axiom III (controlled growth) and connects it naturally to Axiom IV (Euler-system coherence). The main conjecture for elliptic curves [4, 6] is established in the following specific cases:

- **Kato’s divisibility (one direction):** For all primes p where E has good ordinary reduction and $\bar{\rho}_{E,p}$ is surjective. Applies to *all ranks*, but only gives the inclusion $\text{char}_\Lambda(\text{Sel}^\vee) \subseteq (\mathcal{L}_p)$, i.e., the characteristic ideal of the Selmer dual is contained in the ideal generated by the p -adic L -function, providing an upper bound on the Selmer module.
- **Skinner-Urban (full equality):** For ordinary primes p with $p \nmid a_p$, provided $\bar{\rho}_{E,p}$ is irreducible and additional local hypotheses hold. Does *not* cover supersingular primes.
- **Supersingular primes:** Require the signed Selmer group formulation (Kobayashi, Pollack, Sprung); the main conjecture in this setting remains largely open.
- **Not covered:** Primes of bad additive reduction, primes where $\bar{\rho}_{E,p}$ is reducible, and the general case without residual irreducibility.

The main conjecture asserts:

$$\text{char}_\Lambda(\text{Sel}_{p^\infty}(E/\mathbb{Q}_\infty)^\vee) = (L_p(E)), \quad (19)$$

where $\Lambda = \mathbb{Z}_p[[T]]$ is the Iwasawa algebra, $\text{Sel}_{p^\infty}(E/\mathbb{Q}_\infty)^\vee$ is the Pontryagin dual of the Selmer group over the cyclotomic $\mathbb{Z}_p$ -extension, and $L_p(E)$ is the p -adic L -function.

Proposition VII.1 (Iwasawa main conjecture as composition law). *The Iwasawa main conjecture provides the composition structure in the following precise sense: let $\text{char}_\Lambda(M)$ denote the characteristic ideal of a finitely generated torsion Λ -module M . The main conjecture identifies $\text{char}_\Lambda(\text{Sel}_{p^\infty}(E/\mathbb{Q}_\infty)^\vee) = (L_p(E))$. The “composition” is the control theorem (Mazur): specialising at layer n of the $\mathbb{Z}_p$ -tower, the Selmer group at level n is related to the Selmer group at level $n-1$ by an exact sequence whose kernel and cokernel are controlled by the Euler factor at p . This layer-by-layer compatibility is the arithmetic analogue of norm-compatibility in Euler systems.*

The algebraic structure is: $\Lambda = \mathbb{Z}_p[[T]]$ is a power series ring, and the characteristic ideal is generated by a single element $f_E(T)$. The specialisation maps $\chi_n : \Lambda \rightarrow$

³ This paper is part of a broader programme [17]; the connections are purely motivational and not used in any proof.

$\mathbb{Z}_p[\text{Gal}(\mathbb{Q}_n/\mathbb{Q})]$, obtained by evaluating at $T = \zeta_{p^n} - 1$, form a projective system:

$$\chi_m = \chi_n \circ \text{pr}_{m,n} \quad \text{for } m \geq n, \quad (20)$$

where $\text{pr}_{m,n}$ is the natural projection. The control theorem ensures that $\text{Sel}_{p^\infty}(E/\mathbb{Q}_n)$ is determined by $\text{Sel}_{p^\infty}(E/\mathbb{Q}_\infty)$ via restriction with bounded kernel and cokernel. This controlled compatible system is the p -adic composition law constraining the arithmetic of E across the tower.

B. The $\mu = 0$ conjecture

The Iwasawa μ -invariant measures the “ p -adic mass” of Sha in the tower. The conjecture $\mu = 0$ (proven by Ferrero-Washington for abelian fields, expected in general) ensures that no infinite p -power contribution hides in Sha. This is precisely Axiom I restricted to the p -adic tower.

C. Specialization to $s = 1$

The key connection between Iwasawa theory and BSD is the *specialization*: evaluating the p -adic L -function at $T = 0$ (or its derivatives) recovers the complex L -function values:

$$L_p(E, 0) \sim L(E, 1)/\Omega_E^+ \quad (\text{up to periods and Euler factors}). \quad (21)$$

For rank r , the relevant specialization involves the r -th derivative of $L_p(E, T)$ at $T = 0$. The Iwasawa main conjecture then relates this to the characteristic ideal of the Selmer module, providing a p -adic version of the BSD formula.

D. The p -adic to archimedean bridge

The Iwasawa main conjecture and Euler-system machinery (Axioms III–IV) live naturally in the p -adic world, while the BSD conjecture concerns the *complex* L -function $L(E, s)$ at the archimedean place $s = 1$. The passage between these two domains is controlled by Fontaine’s comparison isomorphism $H_{\text{dR}}^1(E/\mathbb{Q}_p) \cong H_{\text{ét}}^1(E, \mathbb{Q}_p) \otimes B_{\text{dR}}$ from p -adic Hodge theory, which provides a dictionary between p -adic periods and complex periods (Ω_E), between p -adic heights ($\hat{h}_p$) and archimedean heights ($\hat{h}$), and between p -adic L -values ($L_p(E, 0)$) and complex L -values ($L(E, 1)$).

For rank 0, the bridge is well understood: at a prime p of good ordinary reduction, the interpolation formula gives $L_p(E, 0) = (1 - \alpha_p^{-1})^2 \cdot L(E, 1)/\Omega_E^+$ up to a p -adic unit, where α_p is the unit root of $X^2 - a_p X + p$ (for multiplicative reduction, the Mazur-Tate-Teitelbaum conjecture, proven by Greenberg-Stevens [20], introduces the

$\mathcal{L}$ -invariant). For rank 1, the Gross-Zagier and Perrin-Riou [26] formulas provide compatible archimedean and p -adic height couplings, closing the bridge at the level of first derivatives.

For rank ≥ 2 , the bridge is the deepest obstruction in the positivity framework. Even if a p -adic height coupling $L_p^{(r)}(E, 0) \sim \det(\hat{h}_p(P_i, P_j))$ were established via diagonal cycles [8], the translation to the archimedean formula $L^{(r)}(E, 1) \sim R_E$ requires two additional ingredients: (i) a comparison between p -adic and archimedean regulators, $R_{E,p} \sim R_E$, which is known up to a p -adic unit for CM curves but open in general; and (ii) an interpolation formula for the r -th derivative generalizing Mazur-Tate-Teitelbaum beyond rank 0.

This bridge crystallizes the structural tension in our framework: the *positivity* (Axiom II) is an archimedean phenomenon – $\hat{h}$ is a real-valued positive-definite form – while the *composition* (Axioms III–IV) is a p -adic phenomenon – Euler systems and Iwasawa modules live over $\mathbb{Z}_p$. Unifying these two “positivity domains” via the comparison isomorphism is the key technical challenge for extending the BSD positivity program beyond rank 1.

VIII. THE PRECISE OBSTRUCTION

A. What blocks a full proof

The positivity framework identifies a precise bottleneck: the *absence of a universal higher Gross-Zagier formula*. Specifically:

1. **For rank 1:** The Heegner point P_K provides a canonical “geometric direction” whose height couples to $L'(E, 1)$. The composition law (Kolyvagin’s Euler system) then excludes alternatives.
2. **For rank r :** One would need r independent canonical points $P_1, \dots, P_r$ whose height *matrix* couples to $L^{(r)}(E, 1)$. The candidates are:
 - Higher Heegner points on Shimura varieties of dimension > 1 .
 - Generalized Heegner cycles (Bertolini-Darmon-Prasanna [9]).
 - Diagonal restriction of p -adic families (Darmon-Rotger [8]).

None currently provides a complete “higher Gross-Zagier formula” in the form (16).

B. The obstruction in positivity language

The obstruction can be stated precisely:

The positivity of $R_E = \det(\langle P_i, P_j \rangle) > 0$ is a theorem (from $\hat{h}$ being positive-definite). The

missing piece is a formula equating $L^{(r)}(E, 1)$ to R_E times computable constants – i.e., the explicit coupling of the analytic and geometric positive structures.

Once such a coupling is established, the No-Go argument closes:

1. $L^{(r)}(E, 1) = C_r \cdot R_E > 0$ (positivity of heights), hence $\text{ord}_{s=1} L(E, s) \leq r$.
2. The Euler-system bound (Axiom IV) gives $\text{rank } E(\mathbb{Q}) \leq \text{ord}_{s=1} L(E, s)$, hence $\text{ord}_{s=1} L(E, s) \geq r$.
3. Together: $\text{ord}_{s=1} L(E, s) = r = \text{rank } E(\mathbb{Q})$.

The entire argument rests on coupling two positive structures – one analytic (L -derivatives), one geometric (height pairing) – and showing they must “saturate” at the same order.

IX. CONNECTION TO THE BROADER PROGRAM

A. The positivity pattern across mathematics

The BSD framework fits into the same positivity pattern observed in:

Problem	Positive form	Composition	Status
Weil conj.	Frobenius ev.	Étale coh.	Proven
Hodge conj.	HR bilinear	Galois-HR	Open
BSD	Néron-Tate	Euler sys.	Rank ≤ 1

In each case:

- The *positivity* is the “easy” part (a theorem or a well-motivated condition).
- The *composition/coherence* is the “hard” part (requires deep structural work).
- The *conjunction* of positivity and composition forces the desired result.

B. Link to the Hodge approach

The Néron-Tate height $\hat{h}$ on $E(\mathbb{Q})$ and the Hodge-Riemann form Q_L on $H^{p,p}(X)$ play structurally identical roles:

- Both are canonical positive-definite bilinear forms.
- Both detect “arithmetic directions” (rational points / algebraic cycles).
- Both require a “composition law” (Euler systems / Galois-HR stability) to force the desired conclusion.

The Hodge analog of the Gross-Zagier formula would be: an explicit identity coupling the “analytic” Hodge-Riemann form to the “geometric” cycle class map, showing that positivity of one forces algebraicity of the other.

C. Link to the Szpiro conjecture

The BSD framework connects to the Szpiro conjecture (equivalent to the *abc* conjecture) through the Tamagawa numbers c_p in the leading-term formula (1).

1. Szpiro as a Tamagawa bound

Conjecture IX.1 (Szpiro [35]). *For every $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that for every elliptic curve $E/\mathbb{Q}$ with minimal discriminant $\Delta_{\min}$ and conductor N :*

$$\log |\Delta_{\min}| \leq (6 + \varepsilon) \log N + C_\varepsilon. \quad (22)$$

For a semistable curve $E/\mathbb{Q}$, all bad reduction is multiplicative. The conductor is $N = \prod_{p|\Delta} p$ and for split multiplicative reduction at p , the Tamagawa number equals the valuation of the minimal discriminant: $c_p = v_p(\Delta_{\min})$ ([15], Table IV.9.1). Thus Szpiro is a *prime-weighted* Tamagawa-exponent bound:

$$\sum_{p|N} c_p \log p = \log |\Delta_{\min}| \leq (6 + \varepsilon) \sum_{p|N} \log p + C_\varepsilon. \quad (23)$$

This is stronger and more structured than an unweighted bound on $\prod_p c_p$: a large exponent at a large prime carries more discriminant mass than the same exponent at a small prime.

2. Diagnostic chain: BSD + bounds expose the missing weighted step

The BSD formula (1) yields:

$$\prod_p c_p = \frac{L^{(r)}(E, 1)}{r!} \cdot \frac{|E_{\text{tors}}|^2}{\Omega_E \cdot R_E \cdot |\text{Sha}|}. \quad (24)$$

This identity controls the *unweighted product* of Tamagawa factors. It does not by itself imply Szpiro, because Szpiro controls the prime-weighted sum in (23). A route from BSD to Szpiro therefore requires the following ingredients plus an additional weighted-local upgrade:

(B1) Upper bound on L -values: $L^{(r)}(E, 1)/r! \leq C \cdot N^{r/2+\varepsilon}$. Unconditional for $r = 0$ (convexity bound); conditional on GRH for general r .

(B2) Lower bound on the period: $\Omega_E \geq c \cdot N^{-1/2-\varepsilon}$. Follows from bounds on the modular parametrisation degree.

(B3) Lower bound on the regulator: $R_E \geq c \cdot (\log N)^r$. Requires an effective lower bound on the Néron-Tate height of non-torsion points.

(B4) Torsion bound: $|E_{\text{tors}}|^2 \leq 256$. This is Mazur's torsion theorem ($|E(\mathbb{Q})_{\text{tors}}| \leq 16$).

(B5) Weighted local upgrade: a mechanism converting unweighted control of $\prod_p c_p$ into control of $\sum_{p|N} c_p \log p$, or directly bounding each local discriminant contribution in terms of conductor data.

Combining (B1)–(B4) under the BSD assumption yields only:

$$\prod_p c_p \leq C' \cdot \frac{N^{r/2+1/2+2\varepsilon}}{(\log N)^r}. \quad (25)$$

For semistable curves of rank 0, this gives a strong unweighted Tamagawa-product constraint. It is *not* yet a Szpiro bound unless (B5) supplies the missing prime weights.

Remark IX.2 (Circularity in step (B3)). Even before the weighted upgrade, the chain has a second obstruction: circularity in (B3). The best known effective lower height bound is Silverman's $\hat{h}(P) \geq c \cdot \log N$ for non-torsion P [15], which is *conditional on the abc conjecture*. Since $abc \Leftrightarrow$ Szpiro, the logical structure would be:

$$\text{BSD} + \underbrace{\text{abc}}_{\text{used in (B3)}} + \underbrace{\text{weighted local upgrade}}_{\text{missing (B5)}} \Rightarrow \underbrace{\text{Szpiro}}_{\Leftrightarrow abc}.$$

Breaking this circularity requires a height bound that does not depend on the discriminant or on abc , and a separate mechanism for the prime-weighted local exponents.

3. A theta-product bridge

The following identity, connecting theta values at torsion points to the Dedekind eta function (and hence to the modular discriminant), provides a potential non-circular link between torsion structure and the discriminant.

Proposition IX.3 (Theta-eta product identity at l -torsion). *For any odd prime l and $\tau \in \mathcal{H}$, with $q = e^{2\pi i \tau}$:*

$$\prod_{j=1}^{l-1} \theta_1\left(\frac{j}{l} \middle| \tau\right) = l \cdot \eta(\tau)^{l-3} \cdot \eta(l\tau)^2, \quad (26)$$

where $\theta_1(z|\tau) = -i \sum_{n \in \mathbb{Z}} (-1)^n q^{(n+1/2)^2/2} e^{(2n+1)\pi i z}$ is the Jacobi theta function and $\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$ is the Dedekind eta function.

Proof. The Jacobi triple product gives $\theta_1(z|\tau) = -2q^{1/8} \sin(\pi z) \prod_{n=1}^{\infty} (1 - q^n)(1 - e^{2\pi i z} q^n)(1 - e^{-2\pi i z} q^n)$. Setting $\omega = e^{2\pi i/l}$, the cyclotomic identity $\prod_{j=1}^{l-1} (1 - \omega^j x) = 1 + x + \dots + x^{l-1}$ applied to each factor of the infinite product, together with $\prod_{j=1}^{l-1} \sin(\pi j/l) = l/2^{l-1}$ (the Chebyshev sine identity), yields:

$$\prod_{j=1}^{l-1} \theta_1(j/l|\tau) = l q^{(l-1)/8} \prod_{n=1}^{\infty} (1 - q^n)^{l-3} (1 - q^{ln})^2.$$

Since $\eta(\tau) = q^{1/24} \prod_{n \geq 1} (1 - q^n)$ and $\eta(l\tau) = q^{l/24} \prod_{n \geq 1} (1 - q^{ln})$, the eta-product $l\eta(\tau)^{l-3}\eta(l\tau)^2$ has exactly the prefactor $lq^{(l-3)/24+2l/24} = lq^{(l-1)/8}$ and the same infinite product. This proves the identity. $\square$

Remark IX.4 (Connection to the discriminant). Since $\eta(\tau)^{24} = \Delta(\tau)/(2\pi)^{12}$ (where Δ is the modular discriminant), identity (26) gives:

$$\sum_{j=1}^{l-1} \log |\theta_1(j/l|\tau)| = \log l + \frac{l-3}{24} \log \frac{|\Delta(\tau)|}{(2\pi)^{12}} + \frac{2}{24} \log \frac{|\Delta(l\tau)|}{(2\pi)^{12}}. \quad (27)$$

For large l , the dominant term is $(l/24) \log |\Delta(\tau)|$: the product of theta values at l -torsion points encodes the discriminant.

Remark IX.5 (Tate curve application and the Szpiro connection). For an elliptic curve $E/\mathbb{Q}_p$ with split multiplicative reduction (Tate curve $E_q \cong \mathbb{G}_m/q^{\mathbb{Z}}$), the l -torsion points correspond to $\zeta_l^j \cdot q^{k/l}$ for $0 \leq j, k < l$. The identity (26) then relates the l -torsion structure of E to $v_p(\Delta_{\min}) = c_p$ *without invoking height bounds or the abc conjecture*. This suggests a potential non-circular path to Szpiro: if the l -torsion theta values can be bounded in terms of the conductor N – for instance via the Galois representation $\bar{\rho}_{E,l}$ or the Weil explicit formula for $L(E, s)$ – the resulting bound on c_p would yield the Szpiro conjecture directly.

The missing step is a prime-weighted local conductor estimate of the form:

$$\sum_{p|N} \sum_{j=1}^{l-1} -\log |\theta_1(j/l|\tau_p)|_p \leq C(l, \varepsilon) \cdot \log N, \quad (28)$$

which is not currently established but is the correct weighted target. Bounding the theta values via the conductor connects naturally to the Galois action on $E[l]$, where Serre's modularity results and level-lowering provide the link between the l -torsion representation and the conductor.

Remark IX.6 (Relation to Mochizuki's IUT approach). The theta-eta product identity (26) operates on the same objects – theta values at torsion points of Tate curves – that appear in Mochizuki's Inter-Universal Teichmüller Theory (IUT). However, the IUT approach uses the theta values q^{j^2} (quadratic in j) arising from the multiradial representation, whereas the identity (26) concerns

$\theta_1(j/l|\tau)$ (linear in j). This distinction is analysed in [36], where it is shown that the $O(l)$ -vs- $O(l^2)$ scaling mismatch between the linear and quadratic theta values constitutes a structural obstruction to reproducing the IUT bound via classical methods. The present approach differs fundamentally: it does not attempt to reproduce the IUT mechanism, but instead uses the product identity as a direct bridge from torsion to discriminant within the BSD framework.

X. DISCUSSION

A. What has been achieved

1. **Axiomatization.** Four axioms – finite capacity, height positivity, controlled Iwasawa growth, Euler-system composition – that jointly force the BSD formula as the unique normal form.
2. **Rank ≤ 1 as a completed instance.** The Gross-Zagier + Kolyvagin argument is naturally a positivity No-Go proof: it couples an analytic derivative to a non-negative height and uses composition (Euler systems) to exclude alternatives.
3. **Higher rank identified.** The obstruction to higher rank is precisely the absence of (H1) a universal higher Gross-Zagier formula and (H2) a complete Kolyvagin system – i.e., the composition law is incomplete.
4. **Exclusion of alternatives.** No alternative leading-term formula is compatible with all four axioms, *provided* a coupling between $L^{(r)}(E, 1)$ and the height pairing exists.

B. Limitations

1. **The paper does not prove BSD.** It reformulates it as a positivity normal-form statement and identifies the missing ingredient.
2. **Axiom I (Sha finiteness) is conjectural for rank ≥ 2 .** This is itself a major open problem, though widely expected.
3. **The “higher Gross-Zagier” is speculative.** No complete formula of the form (16) exists for $r \geq 2$. Recent progress (Darmon-Rotger, Loeffler-Skinner-Zerbes) is promising but incomplete.
4. **The axiomatic framework is descriptive, not constructive.** The axioms identify what must be true, not how to prove it.

C. Outlook

1. **p -adic BSD via Iwasawa theory.** The strongest near-term progress likely comes from extending the Iwasawa main conjecture to higher-rank settings, using the Euler systems of Loeffler-Skinner-Zerbes [14] and the higher rank Kolyvagin-Stark system theory of Burns-Sakamoto-Sano [11, 12].
2. **Higher Gross-Zagier.** Darmon’s program of “Stark-Heegner points” and diagonal cycles provides candidate formulas for rank 2. Establishing these rigorously would close the positivity loop for $r = 2$.
3. **Computational verification.** The BSD formula has been verified numerically for thousands of curves of rank ≤ 3 (see [22]). Extending these computations to higher precision and higher rank tests the framework’s predictions.
4. **Statistical BSD.** The work of Bhargava-Shankar [21] shows that a positive proportion of elliptic curves over $\mathbb{Q}$ have rank 0 and satisfy the BSD conjecture. These statistical results provide further evidence that the positivity framework captures the generic behavior.

Remark X.1 (Arakelov bridge to Hodge). The Néron–Tate height pairing encodes positivity at finite (non-archimedean) primes. Together with the Hodge–Riemann form (positivity at the infinite prime, cf. the companion Hodge paper), the Arakelov intersection pairing provides the natural unified framework in which both are special cases of a single positivity principle.

Remark X.2 (Geometric interpretation of the regulator). The regulator $R_E = \det(\langle P_i, P_j \rangle_{\text{NT}})$ is the covolume of the Mordell–Weil lattice $E(\mathbb{Q})/E(\mathbb{Q})_{\text{tors}}$ in the Néron–Tate metric. The BSD formula equates the analytic datum $L^{(r)}(E, 1)$ to this geometric volume (times local correction factors). A connection to random matrix theory exists via the Keating–Snaith conjectures on the distribution of L -function derivatives, but we do not develop this here.

D. Research directions for higher rank

The following remarks survey concrete approaches toward establishing the missing hypotheses (H1)–(H3) for rank ≥ 2 .

1. *Diagonal cycles and higher Gross-Zagier formulas*

Remark X.3 (Diagonal-cycle / Euler-system hybrid for rank-2 CM curves). For an elliptic curve $E/\mathbb{Q}$ with CM by an imaginary quadratic field K , the Selmer group

$\text{Sel}_{p^\infty}(E/K)$ decomposes into $\pm$ -eigenspaces under complex conjugation. If $\text{rank } E(\mathbb{Q}) = 2$ and E has CM, one could attempt to construct two independent cohomology classes – one from a Heegner-type cycle and one from a Beilinson–Flach element – whose heights span the full regulator matrix. The key challenge is establishing the non-degeneracy of the resulting 2×2 height matrix: $\det(\langle c_i, c_j \rangle) > 0$ must be shown to couple to $L''(E, 1)$. Darmon and Rotger [8] provide partial results in this direction via diagonal cycles on triple products of modular curves.

Remark X.4 (Heegner-simplex approach for rank r). For rank r , one seeks r -dimensional cycles on Kuga–Sato varieties or higher-dimensional Shimura varieties whose Abel–Jacobi images generate a full-rank subgroup of the relevant Selmer group. Concretely, one would need:

1. A Shimura variety $\text{Sh}(G, X)$ of dimension $\geq r$ with special cycles $Z_1, \dots, Z_r$ indexed by CM data;
2. Non-trivial Abel–Jacobi images $\text{AJ}(Z_i) \in H_f^1(\mathbb{Q}, V_p(E))$;
3. An explicit formula: $L^{(r)}(E, 1) = C_r \cdot \det(\langle \text{AJ}(Z_i), \text{AJ}(Z_j) \rangle_p)$.

The group-theoretic framework (replacing GL_2 by GSp_{2r} or a product of copies of GL_2) is provided by Loeffler–Skinner–Zerbes [14], but the explicit height formula remains open.

Remark X.5 (Plectic Gross–Zagier formula). Nekovář and Scholl developed a “plectic” extension of Hodge theory, introducing plectic Galois actions and plectic Hodge structures that generalize the classical Hodge decomposition to higher-rank settings. A *plectic Gross–Zagier formula* would express $L^{(r)}(E, 1)$ in terms of the plectic regulator – a higher determinant constructed from the plectic Hodge structure on the motive of E . This program, still in its early stages, has been continued by Li–Huerta and others; a complete formula would directly establish hypothesis (H1) and close the positivity loop for general rank.

Remark X.6 (p -adic deformation of non-diagonal Heegner cycles). Darmon’s program of Stark–Heegner points constructs p -adic points on E from p -adic integration on Shimura curves, without relying on the Heegner hypothesis. For rank ≥ 2 , one could attempt to p -adically deform families of non-diagonal cycles on higher-dimensional Shimura varieties (e.g., unitary Shimura varieties associated to $\text{U}(2, 1)$ or $\text{U}(3)$), obtaining families of cohomology classes whose specializations provide the required r independent elements. The key obstruction is rationality: the p -adic constructions do not automatically yield $\mathbb{Q}$ -rational points, and a descent argument is needed.

2. Euler systems for higher rank

Remark X.7 (Beilinson–Flach elements and Rankin–Selberg convolutions). Kings, Loeffler, and Zerbes [27] con-

struct Euler systems from Beilinson–Flach elements for Rankin–Selberg convolutions $L(f \otimes g, s)$, where f and g are modular forms. When specialized to the case $f = g$ (the symmetric square), these elements provide information about the p^∞ -Selmer group of E beyond what Kato’s Euler system yields. For rank-2 curves, one could attempt to construct a “rank-2 Kolyvagin system” from the combination of Kato’s classes and Beilinson–Flach elements, each providing control over one eigenspace of the Selmer group. The p -adic Eisenstein elements in multi-dimensional Hida families [13] provide the interpolation framework.

Remark X.8 (Higher Kolyvagin systems and Selmer splitting). Mazur and Rubin [7] developed the abstract theory of Kolyvagin systems: a module over the Iwasawa algebra satisfying certain local conditions that yield upper bounds on Selmer groups. Burns and Sano [11] extended this to higher rank systems over general coefficient rings, and Burns, Sakamoto, and Sano [12] proved the existence of a canonical higher Kolyvagin derivative map between Euler and Kolyvagin systems in any given rank, as conjectured by Mazur and Rubin. For rank r , one needs a “rank- r Kolyvagin system” – a collection of r classes satisfying simultaneous norm-compatibility relations. For CM curves, the endomorphism ring provides a natural splitting of the Selmer group into eigenspaces (under the CM action), and one can attempt to construct independent Kolyvagin classes in each eigenspace. Numerical testing of “rank- r Kolyvagin relations” – verifying the predicted Selmer bounds for specific rank-2 and rank-3 curves with CM – would provide evidence for this approach.

Remark X.9 (Recent progress on rank ≥ 2). Significant progress toward higher-rank BSD has been achieved in recent years. Burungale, Castella, and Kim [10] proved Perrin–Riou’s Heegner point main conjecture, building on Howard’s bipartite Euler systems and Wei Zhang’s work on Kolyvagin’s conjecture, establishing the Iwasawa main conjecture for the Tate–Shafarevich group over anticyclotomic extensions. Kim [5] obtained a higher Gross–Zagier formula by comparing Heegner point Kolyvagin systems with Kurihara numbers, refining the structural understanding of Selmer groups. In 2024, Loeffler and Zerbes [30] constructed a universal Euler system for $\text{GSp}(4)$, showing that all Euler system classes lie in a one-dimensional space. Burungale, Skinner, Tian, and Wan [31] proved the existence of p -adic zeta elements for elliptic curves over imaginary quadratic fields, establishing the Kobayashi conjecture for semistable curves at supersingular primes and presenting the first infinite families of non-CM elliptic curves satisfying the full BSD conjecture.

Threshold axiom: higher Gross–Zagier

Axiom X.10 (HGZ Bridge – HGZ-BSD). *For an elliptic curve $E/\mathbb{Q}$ of analytic rank $r = \text{ord}_{s=1} L(E, s)$, there*

exists:

(HGZ-A) **Canonical cycles:** A family of arithmetic cycles $\mathcal{Z}_r^{(i)}$ ($i = 1, \dots, r$) in the appropriate Chow group or motivic cohomology, functorial under Hecke operators and base change.

(HGZ-B) **Height compatibility:** A canonical height pairing $\langle \cdot, \cdot \rangle_{\text{ht}}$ on these cycles, compatible with the Néron–Tate height for $r = 1$ and satisfying $\det(\langle \mathcal{Z}_r^{(i)}, \mathcal{Z}_r^{(j)} \rangle_{\text{ht}}) = R_E \cdot (\text{local factors})$.

(HGZ-C) **Selmer control:** An Euler system of rank r controlling the Selmer group $\text{Sel}(E/\mathbb{Q})$, yielding $|\text{Sha}(E/\mathbb{Q})| < \infty$ and the reverse inequality $\text{ord} \leq \text{rank}$.

The conjunction (HGZ-A)+(HGZ-B)+(HGZ-C) implies the full BSD formula for rank r .

Remark X.11 (Module status).

Module	Rank 1	Rank
(A) Canonical cycles	Heegner points (GZ 1986)	Diagonal cycles (Darmon–Rotger), Partial
(B) Height compatibility	GZ formula (1986)	Open (higher GZ formula)
(C) Selmer control	Kolyagin (1988)	Partial: $\text{GSp}(4)$ Euler systems [30]

The red block for rank ≥ 2 is primarily **Module C**: existing Euler systems (Loeffler–Zerbes for $\text{GSp}(4)$, Burungale–Skinner–Tian–Wan [31] for imaginary quadratic fields) provide partial control but do not yet yield a complete Kolyagin-type bound on Sha for individual curves of rank ≥ 2 .

Remark X.12 (No-go mechanisms). HGZ-BSD fails if:

1. **Non-canonical cycle choice:** The higher cycles $\mathcal{Z}_r^{(i)}$ depend on auxiliary data (quaternion algebras, test vectors) in a way that cannot be uniformised—this is a language problem, not a computational one.
2. **Degenerate heights:** The height pairing $\langle \cdot, \cdot \rangle_{\text{ht}}$ has a non-trivial kernel for $r \geq 2$, collapsing the determinant. Height saturation (Remark VI.3) guards against this.
3. **Euler system weakness:** The rank- r Euler system has insufficient variability to control all p -primary components of Sha simultaneously.

Remark X.13 (Module C as arithmetic underdetermination). Without rank- r Euler system control (Module C), the Higher Gross–Zagier formula remains *arithmetically underdetermined*: heights can be large without implying Selmer structure. This is the precise content of the red block: the bridge lemma (HGZ-A)+(HGZ-B)+(HGZ-C) $\Rightarrow L^{(r)} \sim \det(\text{Heights})$ requires all three modules, and Module C is the only one without even a partial result for general rank $r \geq 2$ curves. Loeffler–Zerbes [30] provide partial $\text{GSp}(4)$ control, but a full Kolyagin-derivative argument for $|\text{Sha}| < \infty$ at rank ≥ 2 remains open.

3. Arakelov and motivic perspectives

Remark X.14 (Universal Arakelov regulator). The BSD formula involves both p -adic data (Selmer groups, Iwasawa modules) and archimedean data (periods Ω_E , heights $\hat{h}$). The Arakelov intersection theory on the regular model $\mathcal{E} \rightarrow \text{Spec}(\mathbb{Z})$ provides a unified framework: the Arakelov regulator $\hat{R}_E$ combines the archimedean Néron–Tate height with p -adic local contributions into a single adelic height. A “universal Arakelov regulator” as a roof over both p -adic and archimedean regulators would provide the natural bridge for hypothesis (H1): one formula $L^{(r)}(E, 1) = C_r \cdot \hat{R}_E$ valid at all places simultaneously.

Remark X.15 (Arakelov-motivic volume formula). One can attempt to interpret $L^{(r)}(E, 1)$ as the measure of an “arithmetic volume” associated to an Arakelov vector bundle on the arithmetic surface $\mathcal{E}$. In this picture, R_E is the volume of the fundamental domain of the Mordell–Weil lattice (with respect to the Néron–Tate metric), Ω_E is the archimedean volume of the fibre, and $|\text{Sha}|$ is a “defect volume” measuring the failure of local-global compatibility. The BSD formula then asserts that the higher GZ formula (higher arithmetic Euler characteristic – computed via the L -function – equals this motivic volume. Connecting this to the Tamagawa number conjecture [33, 34] would provide a conceptual derivation of the BSD formula. Indeed, the Bloch–Kato conjecture already provides the rigorous motivic framework for the uniqueness question discussed in Proposition IV.4: it defines Tamagawa numbers for general motives and derives the BSD formula as the special case for $h^1(E)(1)$, removing the rational ambiguity of the Beilinson conjectures.

Remark X.16 (Hodge-Arakelov volumes: Sha as topological volume). The Hodge-Arakelov theory of Mochizuki interprets the comparison between de Rham and étale cohomology of elliptic curves as a “discrete analogue” of the Hodge decomposition. In this framework, $|\text{Sha}|$ could be viewed as the measure of a topological volume – the “obstruction to global triviality” – that is forced to be finite by positivity constraints analogous to the Hodge–Riemann relations. If a positivity framework could be applied to this topological volume, finiteness of Sha (Axiom I / hypothesis H3) might follow from positivity alone.

4. Algebraic and categorical approaches

Remark X.17 (Higher Chow groups on Picard modular surfaces). For rank $r \geq 2$, the relevant motives are no longer one-dimensional: the height pairing involves an $r \times r$ regulator matrix. The natural algebraic framework is provided by higher Chow groups $\text{CH}^r(X, r)$ on Picard modular surfaces (Shimura varieties for $\text{U}(2, 1)$) or on self-products of the modular curve. The derived intersection pairing on these higher Chow groups generalizes the Néron–Tate pairing and could provide the $r \times r$ height matrix needed for hypothesis (H1). The connection to L -

functions is via the Beilinson conjectures, which predict that special values of L -functions are related to regulators on higher Chow groups.

Remark X.18 (Derived Hecke algebras and Taylor-Wiles defect). Venkatesh’s derived Hecke algebra program relates the Taylor-Wiles method (patching completed cohomology) to derived algebraic geometry. The “Taylor-Wiles defect” – the discrepancy between the dimension of a deformation ring and the dimension of the cohomology – is conjectured to equal the rank of the Mordell-Weil group. If this could be made precise for elliptic curves, it would provide a new route to the rank part of BSD: $\text{rank } E(\mathbb{Q}) = \text{TW defect} = \text{ord}_{s=1} L(E, s)$.

5. Analytic and structural approaches

Remark X.19 (L -function decomposition into Galois sectors). The L -function $L(E, s)$ decomposes under twists by characters χ of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$: the Hasse-Weil-Artin L -function $L(E, \chi, s)$ encodes the arithmetic of E in the “ χ -sector.” By analogy with the topological sector decomposition in Yang-Mills theory (cf. the companion YM paper), one could decompose the BSD problem sector-by-sector: proving BSD for each twisted L -function $L(E, \chi, s)$ separately, then assembling the full result. Darmon and Rotger [8] pursue exactly this strategy for Artin representations, establishing BSD for Hasse-Weil-Artin L -functions in rank 1 cases.

Remark X.20 (Cohomological sweeping processes for Sha finiteness). The finiteness of Sha (Axiom I / hypothesis H3) can be reformulated as a statement about the global-to-local map in Galois cohomology: $\text{Sha}(E/\mathbb{Q}) = \ker(H^1(\mathbb{Q}, E) \rightarrow \prod_v H^1(\mathbb{Q}_v, E))$. A “sweeping process” approach from convex analysis would attempt to show that this kernel is finite by iteratively “sweeping” local obstructions: at each prime v , the local contribution $H^1(\mathbb{Q}_v, E)$ absorbs a definite proportion of the global classes, leaving only finitely many in the kernel. The key input would be a uniform bound on the “absorption rate” across primes, which is related to the Tamagawa numbers c_v and the Lang-Trotter heuristics.

Remark X.21 (O-minimal structures and Sha finiteness). The o-minimal methods of Bakker-Brunebarbe-Tsimerman [28] (successfully applied to the Hodge conjecture, cf. the companion Hodge paper) could potentially be applied to Sha finiteness. The key idea: the period domain for the family of elliptic curves over $\mathbb{Q}$ is a definable analytic space in an o-minimal structure, and the set of curves with infinite Sha (if any exist) would form a definable subset. The tameness properties of o-minimal structures (finite stratification, controlled topology) could impose constraints on this locus, potentially showing it is empty. This is speculative but connects the BSD and Hodge programs at a structural level.

E. Numerical verification

The BSD formula has been verified to high precision for thousands of elliptic curves up to rank 4 (LMFDB [22]). The positivity framework makes specific numerical predictions:

- For every curve with $\text{rank } E(\mathbb{Q}) = r$, the ratio $L^{(r)}(E, 1)/(r! \cdot \Omega_E \cdot R_E \cdot \prod c_p)$ must equal $|\text{Sha}|/|E_{\text{tors}}|^2$ – a positive rational number, where $|\text{Sha}|$ is itself a perfect square by the alternating Cassels-Tate pairing.
- The regulator R_E – being the Gram determinant of a positive-definite form – provides an a priori *lower bound* on $|L^{(r)}(E, 1)|$ (given the other BSD factors).
- For rank-2 CM curves, the Selmer splitting predicts a factorization of the Kolyvagin bound into eigenspace contributions.

The accompanying Examples V.4–V.6 verify these predictions for specific curves of rank 0, 1, and 2.

F. Post-Zookeeper transfer: a higher-rank module programme

The post-Zookeeper use of the present paper is not to claim a new proof of BSD in higher rank. Rather, it turns the higher-rank obstruction into a modular transfer problem. The relevant five slots are:

In particular, the CORE closure of Riemann- $C2/MS2$ has no automatic BSD consequence. It only fixes the shared normal-form discipline: identify an operator or form, an explicit defect, a finite approximation scheme, a gap, and a limiting passage. For BSD the red block remains higher-rank Selmer control and the higher Gross-Zagier bridge, especially Module C in Remark X.11.

Operator/form.: The Neron-Tate and p -adic height pairings on Mordell-Weil and Selmer directions.

Defect.: The BSD defect

$$\delta_{\text{BSD}}(E) := \text{ord}_{s=1} L(E, s) - \text{rank } E(\mathbb{Q}),$$

together with the residual between the predicted leading term and the regulator expression.

Approximation.: Finite conductor, fixed rank, finite Selmer level, and finite layers in an Iwasawa tower.

Gap.: Non-degeneracy of the archimedean or p -adic regulator.

Limit.: The passage from p -adic leading-term information and Selmer structure to the archimedean BSD leading-term formula.

This suggests a conservative next target. Instead of formulating a general rank- r proof, one should first isolate a rank-two transfer statement of the form

(higher Gross–Zagier input)+(Selmer control)+(p-adic height) $\Rightarrow$ (rank ≤ 1)

Kim’s higher Gross–Zagier and Selmer/Iwasawa work, together with the Castella–et al. p -adic BSD framework, provide the natural literature backbone for this programme. The expected gain is diagnostic rather than final: higher rank is decomposed into four separately testable modules instead of remaining a single monolithic gap.

Result classification

This paper establishes a **structural normal-form characterisation**: under four positivity axioms, the BSD formula is the unique admissible shape for the leading term of $L^{(r)}(E, 1)$. For rank ≤ 1 , this is a genuine reformulation of proven results; for rank ≥ 2 , it is a conditional structural analysis.

XI. CONCLUSION

The Birch and Swinnerton-Dyer conjecture, like the Hodge Conjecture and the Weil conjectures, is fundamentally a statement about the impossibility of discrepancy between analytic and geometric structures. The Néron-Tate height provides the canonical positivity structure: it is a positive-definite form on rational points that detects arithmetic directions via their energy cost.

The BSD formula is the unique “normal form” compatible with four axiomatic constraints: finite arithmetic capacity (Sha finiteness), cooperative height positivity (Néron-Tate), controlled Iwasawa growth (predictable growth in $\mathbb{Z}_p$ -towers), and Euler-system composition coherence (norm-compatibility).

The rank ≤ 1 proof is already a positivity No-Go argument: Gross-Zagier couples an analytic derivative to a non-negative height, and Kolyvagin’s Euler system excludes alternatives. For higher rank, the framework identifies the precise missing ingredient: a universal “higher Gross-Zagier” formula coupling $L^{(r)}(E, 1)$ to the positive-definite $r \times r$ regulator matrix.

For rank ≤ 1 , our contribution is conceptual: a new perspective via positivity normal form. For rank ≥ 2 , the Height Saturation Conjecture identifies the precise mathematical target for future work.

Result	Type	“BSD does not ask us to find rational points. It asks us to show that the L -function has no room for zeros without them.”
<i>Proven (unconditional):</i>		
Néron–Tate positivity (Axiom II)	Classical theorem	
BSD normal form (rank ≤ 1)	Theorem (Gross–Zagier + Kolyvagin)	
Height saturation (rank 2, numerical)	17/17 curves verified	
Regulator scaling $R \sim N^{0.10}$	Numerical	
<i>Structural contributions:</i>		This work was conducted as independent research without institutional funding.
Axiom system (I–IV)	Definition	
Uniqueness of BSD shape	Proposition	<i>AI tools.</i> AI-based tools (Claude by Anthropic) were used for mathematical formalization and text generation.
Gross–Zagier as prototype	Interpretation	
<i>Open (fundamental):</i>		All mathematical content, conjectures, and responsibility for correctness lie solely with the author.
Higher Gross–Zagier ($r \geq 2$)	Open	
Higher Kolyvagin systems	Open	<i>Funding information.</i> This research received no external funding.
Sha finiteness ($r \geq 2$)	Open	

ACKNOWLEDGMENTS

-
- [1] B. J. Birch & H. P. F. Swinnerton-Dyer (1965). Notes on elliptic curves. II. *Journal für die reine und angewandte Mathematik*, 218, 79–108. DOI: 10.1515/crll.1965.218.79.
- [2] B. Gross & D. Zagier (1986). Heegner points and derivatives of L -series. *Inventiones Mathematicae*, 84, 225–320. DOI: 10.1007/BF01388809.
- [3] V. A. Kolyvagin (1990). Euler systems. In *The Grothendieck Festschrift*, Vol. II, pp. 435–483. Birkhäuser. DOI: 10.1007/978-0-8176-4575-5_11.
- [4] K. Kato (2004). p -adic Hodge theory and values of zeta functions of modular forms. *Astérisque*, 295, 117–290.
- [5] C.-H. Kim (2024). A higher Gross-Zagier formula and the structure of Selmer groups. To appear in *Transactions of the AMS*. arXiv:2203.12161.
- [6] C. Skinner & E. Urban (2014). The Iwasawa main conjectures for GL_2 . *Inventiones Mathematicae*, 195(1), 1–277. DOI: 10.1007/s00222-013-0448-1.
- [7] B. Mazur & K. Rubin (2004). *Kolyvagin Systems*. Memoirs of the AMS, 168(799). DOI: 10.1090/memo/0799.
- [8] H. Darmon & V. Rotger (2017). Diagonal cycles and Euler systems II: The Birch and Swinnerton-Dyer conjecture for Hasse-Weil-Artin L -functions. *Journal of the AMS*, 30(3), 601–672. DOI: 10.1090/jams/861.
- [9] M. Bertolini, H. Darmon, & K. Prasanna (2013). Generalized Heegner cycles and p -adic Rankin L -series. *Duke Mathematical Journal*, 162(6), 1033–1148. DOI:

- 10.1215/00127094-2142056.
- [10] A. Burungale, F. Castella, & C.-H. Kim (2021). A proof of Perrin-Riou's Heegner point main conjecture. *Algebra & Number Theory*, 15(7), 1627–1653. DOI: 10.2140/ant.2021.15.1627.
 - [11] D. Burns & T. Sano (2021). On the theory of higher rank Euler, Kolyvagin and Stark systems. *International Mathematics Research Notices*, 2021(13), 10118–10206. DOI: 10.1093/imrn/rnz103.
 - [12] D. Burns, R. Sakamoto, & T. Sano (2025). On the theory of higher rank Euler, Kolyvagin and Stark systems, II: the general theory. To appear in *Algebra & Number Theory*. arXiv:1805.08448.
 - [13] A. Lei, D. Loeffler, & S. L. Zerbes (2014). Euler systems for Rankin-Selberg convolutions of modular forms. *Annals of Mathematics*, 180(2), 653–771. DOI: 10.4007/annals.2014.180.2.6.
 - [14] D. Loeffler, C. Skinner, & S. L. Zerbes (2022). Euler systems for $\mathrm{GSp}(4)$. *Journal of the European Mathematical Society*, 24(2), 669–733. DOI: 10.4171/JEMS/1124.
 - [15] J. H. Silverman (2009). *The Arithmetic of Elliptic Curves*, 2nd ed. Graduate Texts in Mathematics 106, Springer. DOI: 10.1007/978-0-387-09494-6.
 - [16] A. Wiles (1995). Modular elliptic curves and Fermat's Last Theorem. *Annals of Mathematics*, 141(3), 443–551. DOI: 10.2307/2118559.
 - [17] L. Geiger (2026). The Saturation Theorem: Axiomatic Constraints on Quantum Gravity from Cosmological Relaxation. Zenodo preprint. doi:10.5281/zenodo.19036188.
 - [18] L. Geiger (2026). The Renormalized Free Energy Framework. Zenodo preprint. doi:10.5281/zenodo.19036190.
 - [19] L. Geiger (2026). The Riemann Hypothesis Part III: The Analytical Rank-Finite Certificate. Zenodo preprint. doi:10.5281/zenodo.19035640.
 - [20] R. Greenberg & G. Stevens (1993). p -adic L -functions and p -adic periods of modular forms. *Inventiones Mathematicae*, 111(2), 407–447. DOI: 10.1007/BF01231294.
 - [21] M. Bhargava & A. Shankar (2015). Ternary cubic forms having bounded invariants, and the existence of a positive proportion of elliptic curves having rank 0. *Annals of Mathematics*, 181(2), 587–621. DOI: 10.4007/annals.2015.181.2.4.
 - [22] The LMFDB Collaboration (2024). *The L -functions and Modular Forms Database*. <https://www.lmfdb.org>.
 - [23] J. Nekovář (2006). Selmer complexes. *Astérisque*, 310, vi+559.
 - [24] T. Dokchitser & V. Dokchitser (2010). On the Birch-Swinnerton-Dyer quotients modulo squares. *Annals of Mathematics*, 172(1), 567–596. DOI: 10.4007/annals.2010.172.567.
 - [25] X. Yuan, S.-W. Zhang, & W. Zhang (2013). *The Gross-Zagier Formula on Shimura Curves*. Annals of Mathematics Studies 184, Princeton University Press.
 - [26] B. Perrin-Riou (1987). Points de Heegner et dérivées de fonctions L p -adiques. *Inventiones Mathematicae*, 89(3), 455–510. DOI: 10.1007/BF01388982.
 - [27] G. Kings, D. Loeffler, & S. L. Zerbes (2017). Rankin-Eisenstein classes and explicit reciprocity laws. *Cambridge Journal of Mathematics*, 5(1), 1–122. DOI: 10.4310/CJM.2017.v5.n1.a1.
 - [28] B. Bakker, B. Brunebarbe & J. Tsimerman (2023). o -minimal GAGA and a conjecture of Griffiths. *Inventiones Mathematicae*, 232(1), 163–228. DOI: 10.1007/s00222-022-01166-1.
 - [29] A. Smith, 2^∞ -Selmer groups, 2^∞ -class groups, and Goldfeld's conjecture, arXiv:1702.02325v3, 2022.
 - [30] D. Loeffler and S. L. Zerbes, *A universal Euler system for $\mathrm{GSp}(4)$* , arXiv:2411.12576, 2024.
 - [31] A. Burungale, C. Skinner, Y. Tian, and X. Wan, *Zeta elements for elliptic curves and applications*, arXiv:2409.01350, 2024.
 - [32] K. Česnavičius (2018). The Manin constant in the semistable case. *Compositio Mathematica*, 154(9), 1889–1920. DOI: 10.1112/S0010437X18007273.
 - [33] S. Bloch & K. Kato (1990). L -functions and Tamagawa numbers of motives. In *The Grothendieck Festschrift*, Vol. I, pp. 333–400. Birkhäuser. DOI: 10.1007/978-0-8176-4574-8.9.
 - [34] J.-M. Fontaine & B. Perrin-Riou (1994). Autour des conjectures de Bloch et Kato: cohomologie galoisienne et valeurs de fonctions L . In *Motives* (Seattle 1991), Proc. Symp. Pure Math. 55, Part 1, pp. 599–706. AMS.
 - [35] L. Szpiro (1981). Propriétés numériques du faisceau dualisant relatif. In: *Séminaire sur les pinceaux de courbes de genre au moins deux*, Astérisque **86**, 44–78.
 - [36] L. Geiger (2026). Structural limits of the classical route to Corollary 3.12: a theta-product diagnostic. Preprint (Bernau).