

Digital Antianxiety Treatment and Cognitive Performance: an Experimental Study

Elisa Cavatorta ^{*} Simona Grassi [†] Mark Lambiris [‡]

November 6, 2020

Abstract

We study the impact of an innovative antianxiety digital treatment on performance in a cognitive demanding task using a randomized control trial design. We exogenously manipulate a cognitive bias associated with anxiety – the tendency to disproportionately allocate attention to negative stimuli – in a group of young males and females. We observe that the treatment reduces the cognitive bias of treated females and significantly improves their performance in the cognitive task. Evidence from university examination results taken three months after the intervention suggests that the digital treatment may generate long-lasting benefits.

JEL codes: C91; I10; J16.

Keywords: Anxiety; Cognitive Performance; Gender; Randomized Control Trial.

1 Introduction

Anxiety disorders are a significant public health concern across the globe. According to the US National Institute of Mental Health (NIMH), anxiety disorders are the most

^{*}Department of Political Economy, King's College London, United Kingdom. Email: elisa.cavatorta@kcl.ac.uk

[†]King's Business School, Economics, King's College London, United Kingdom. Email: simona.grassi@kcl.ac.uk

[‡]Swiss Tropical and Public Health Institute and University of Basel, Basel, Switzerland. E-mail: mark.lambiris@swisstph.ch

Declarations of interest: none. The study was financed by a Swiss National Science Foundation (SNSF) grant (Project number: PDFMP1137024). This study is registered in the American Economic Association (AEA) RCT Registry and the unique identifying number is: AEARCTR-0005161.

common mental illness in the USA, affecting 19% of the American adult population annually. In England, data from the most recent Adult Psychiatric Morbidity Survey 2014 (McManus et al., 2016) show that about 17% of adults had been affected by a common mental disorder in the previous week,¹ with a mix of anxiety and depression (7.8%) and generalized anxiety disorder (5.9%) being the two most common disorders.² Some longitudinal studies document an increasing prevalence of anxiety and depression, particularly among the young population (Twenge et al., 2010).³ A meta-analysis of anxiety prevalence statistics across countries shows that females are on average twice as likely to suffer from anxiety than males (e.g. Remes et al., 2016), a remarkable difference even when possible gender differences in symptom reporting are taken into account.⁴

Anxiety has important and long-lasting repercussions on individuals' health, social functioning and wellbeing. In the UK, mental illnesses are estimated to cost the economy about 4.5% of GDP every year (Stansfeld et al., 2016). Das-Munshi et al. (2008) estimate that mixed anxiety and depression are responsible for one-fifth of days off work, representing the most frequent cause of sickness absence in England. According to the UK Office for National Statistics (2018) there has been an increase in the proportion of younger workers aged 25-34, who attribute their sickness absence to mental health conditions, rising from 7.2% in 2009 to 9.6% in 2017.⁵

Several studies show that for the same level of ability, anxiety reduces educational performances and professional outcomes (see, e.g. Beard, 2011): anxiety can be thought of as a "tax on performance". Since anxiety prevalence varies across socio-demographic groups, the detrimental effect on performance can be expected to affect some groups more than others. A number of studies find gender differences in performance in high-pressure

¹Common mental disorders include depression, generalized anxiety disorder, panic, phobias, obsessive compulsive disorder, and other not otherwise specified disorders referred to as "mixed anxiety and depression".

²Mental health statistics for the U.S. are taken from NIMH at <http://www.nimh.nih.gov/health/statistics/prevalence/any-anxiety-disorder-among-adults.shtml> and refer to the years 2001-2003; McManus et al. (2016) is available at <https://webarchive.nationalarchives.gov.uk/20180328140249/http://digital.nhs.uk/catalogue/PUB21748>

³Part of the observed increase in prevalence may be due to improvements in diagnosis and changes in the acceptability of reporting mental health disorders, but studies that take these confounders into account still show increases in the prevalence of mental health disorders (Twenge et al., 2010).

⁴According to the Adult Psychiatric Morbidity Survey 2014 Report from England (McManus et al., 2016), for the same level of mental health disorder (and assuming everyone who reports the disorder, receives treatment), for every five females who report it, only two out of five males report the same disorder.

⁵The ONS report and data are available at <https://www.ons.gov.uk/employmentandlabourmarket/peopleinwork/employmentandemployeetypes/articles/sicknessabsencefallstothelowestratein24years/2018-07-30>.

environments where anxiety is likely to arise. For example, Paserman (2007) shows that in tennis tournaments females players make more errors than males as competitive pressure increases. In a controlled experiment Cahliková et al. (2019) find that women under competitive conditions perform worse when they are stressed relative to when they are not stressed. Coffman (2014) shows that females attempt fewer questions in a quiz than males, even after controlling for knowledge and confidence. Anxiety may explain these differences: if anxiety triggers behaviours that are associated with poorer performance in stressful situations (e.g., being less focused), then the anxiety-driven penalty on performance penalizes females more than it does males. Given the prevalence of examination- and quiz-based assessments as screening devices, the anxiety tax on performance may be partly responsible for the gender imbalances observed in quiz-based rankings and the prejudices around females being less skillful in quantitative subjects (e.g. Guiso et al., 2008).⁶

An important question arises: Can specialized interventions help mitigate the detrimental impact of anxiety on performance? Here we provide an answer by evaluating the impact of an innovative form of antianxiety training, the cognitive bias modification (CBM), on performance in a cognitive test in groups of young males and females, using a randomized control trial (RCT) with a difference-in-difference design. Studies in cognitive psychology and neuroscience have shown that anxiety impairs the use of attentional control resources that are necessary inputs for successful performance in cognitive tasks.⁷ CBM treatment aims to restore the use of attentional control resources.

CBM treatment was developed by cognitive psychologists (MacLeod et al., 1986) and is based on the premise that some of the cognitive biases through which an individual perceives reality are a cause of anxiety. CBM addresses attention bias, one of the most widely recognized biases associated with anxiety. Attention bias is the tendency to disproportionately and more rapidly attend to threatening stimuli in comparison with other concurrent neutral stimuli.⁸

⁶For example, in 2016 twice as many male students in Mathematics and Computer Science at the University of Oxford received a first-class honours – the highest grade classification in the UK higher education system – than female students. As a response, the university extended the duration of examinations in the belief that the observed gender gap in academic performance was due to females being more adversely affected by time pressure in examinations than males. See <https://www.telegraph.co.uk/education/2018/02/01/oxford-university-extends-exam-times-womens-benefit/>

⁷Attentional control resources include the ability to shift attention to specific features of the task that are most relevant for its completion, the ability to inhibit attention to distractors and interferences, and the ability to keep track of and to retrieve relevant information (see Derakshan and Eysenck, 2009).

⁸Other biases associated with anxiety are memory bias and interpretative bias. The former is the tendency to retrieve memories selectively favoring negative information, while the latter is the tendency to resolve ambiguity toward a negative direction. The experimental psychopathology literature has focused

Anxiety reduces performance because it subconsciously drives an individual’s attention toward threatening distractors that are irrelevant to the task to be completed (see, e.g. Eysenck and Derakshan, 2011; Wilson and MacLeod, 2003). For example, while taking a test, an anxious student is more likely to be distracted by her worries about the test or other threatening cues in the surrounding environment. Performance will suffer from attentional control resources being diverted away from stimuli relevant to the completion of the task toward task-irrelevant stimuli. Anxious individuals may compensate for decrements in performance by exerting additional effort. However, such effort is costly and its effectiveness is decreasing, especially when the task is demanding.⁹

CBM treatment reduces attention bias by training individual attention. CBM is a computer-based training in which pairs of threatening and neutral stimuli appear sequentially on the screen. The subject is trained to divert attention away from threatening stimuli by requiring repeated identification of the location of the neutral stimulus in a pair of neutral and threatening stimuli. We expect CBM-treated participants to reduce their attention bias and to improve their performance in a cognitive task. The improvement in performance originates from liberating cognitive resources previously trapped by the focus on threatening and anxiety-generating stimuli. Larger reductions in attention bias may be expected in more anxious individuals, such as females, leading in turn to greater improvement in task performance.

CBM treatment has advantages in comparison with alternative treatments such as cognitive behavioural therapy (CBT) and antidepressant medications that aim to reduce the consequences of anxiety or the anxiety itself. The meta-analysis by Hakamata et al. (2010) shows that, on clinical patient samples, CBM treatment yields effect sizes of comparable magnitude to those from CBT and drug treatments. However, CBM is cheaper, easier to administer to patients, amenable to large-scale interventions, non-invasive and, insofar as it has been tested, well accepted by patients (Beard, 2011).

Our results show that CBM reduces attention bias in treated females and produces significant improvements in performance in a cognitive task for this group, measured as the number of attempted questions in the test. Treated females’ performance increases on average by 38% compared with control group females’ performance. Males do not show a significant reduction in bias, and consequently, have no performance improvement. This is

so far mainly on attention bias.

⁹Attentional control theory predicts that anxiety impairs attention control even in cases where there are no threat-related stimuli irrelevant to completion of the task. This is because the optimal strategy of anxious individuals, who often feel under general threat, is to allocate attentional resources widely in order to preempt dangers, rather than concentrating attentional resources on specific focuses. This implies a reduction in the attentional control resources available for any given cognitive task.

not due to a differential engagement with the treatment.¹⁰ A more plausible explanation is that males are originally less anxious than females and may have less to gain from the treatment. Rather, it is the anxious females, who are more likely to be penalized by anxiety, who obtain the largest increases in performance.

This paper makes several contributions. First, our results inform the use of digital health treatments against one of the most common mental health conditions worldwide: anxiety. The results in this paper focus on CBM and extend the recent evidence on its efficacy in two ways. i) The results evidence the impact of CBM on a real behavioral outcome typically hindered by anxiety: performance in a cognitive task. Whereas previous psychological and psychopathological studies have demonstrated encouraging effects of CBM on measures of attention bias and anxiety, this study brings these findings a step forward by analysing the resulting changes in behaviour. ii) The study’s design tests the efficacy of CBM using a novel group of people that is larger and more representative of the levels of anxiety in the general population than previous studies, most of which were confined to small samples of pathologically anxious patients.¹¹ In contrast, this study’s sample comprises 261 university students, with a broad range of anxiety levels reflecting the distribution of anxiety in the general population. For this reason, our findings are the first, as far as we know, to inform of the expected impact of CBM on a ‘typical’ user, as opposed to selective groups of responsive users.

Second, the finding that exogenous manipulation of attention bias can affect performance in cognitive tasks contributes to the understanding of the range of non-cognitive factors affecting performance. Our work adds to the literature on the roles of personality traits (Borghans et al., 2008; Moutafi et al., 2006), grit and determination (Alan et al., 2019). By focusing on a sample of young adults, the paper also contributes to the scant body of evidence on the response of non-cognitive skills to interventions in adulthood (Blattman et al., 2017, e.g.). Despite the recognised malleability of attitudes and behaviours during young adulthood (Krosnick and Alwin, 1989), the empirical evidence is thin in comparison to the large number of works documenting interventions in early childhood.

Third, the paper contributes to the current debate on the limitations of exams and

¹⁰In additional analysis, we have studied whether the difference in the effect of CBM between genders could be attributed to female participants complying with the instructions differently compared to males and/or engaging with the CBM training with a different degree of commitment than males, to an extent sufficient to create different levels of intensity of the CBM treatment. However, we did not find any strong supporting evidence for this hypothesis. The analysis is available upon request.

¹¹For example, among the pool of 12 studies included in the most recent meta-analysis of CBM efficacy by Hakamata et al. (2010), the largest sample size used comprises 94 subjects.

standardised tests as screening devices for cognitive ability (Rear, 2019). Our results show that the performance of anxiety-prone individuals significantly increases following a reduction in attention bias. This is indicative of the extent to which typical assessments of cognitive ability may be poor proxies due to the impact of anxiety.

The remainder of paper is organized as follows: Section 2 explains the CBM treatment; Section 3 describes the experimental design; Section 4 presents the empirical strategy; Section 5 provides evidence that the CBM treatment was effective in reducing attention bias and increasing performance in a cognitive task; Section 6 concludes the paper. The Supplementary Material (SM) discusses robustness tests and contains the experimental instructions.

2 The Treatment: Cognitive Bias Modification

CBM treatment aims to mitigate the consequences of anxiety by re-diverting the sub-conscious focus on threatening stimuli toward neutral or positive ones. The treatment is computer-based and involves the rapid sequential appearance of pairs of visual stimuli on the screen. We adopted the version of CBM that uses pairs of photographs of human faces as visual stimuli. In each pair of photographs, the face of the same person appears with an angry expression (threat stimulus) and an emotionally neutral expression (neutral stimulus). Figure SM.1 in the SM presents an example of the pair of faces used in the experiment.¹²

Participants see each pair of angry and neutral faces at the center of their computer screen for 500 milliseconds (ms), before the faces disappear. The photographs are juxtaposed from left to right with a space in between. The location of the angry and the neutral faces on the right or on the left of the screen is randomized. After the faces have disappeared, a probe (in the form of a small cross) pops up for 500 ms at the location of one of the two images. The task for the participant is to identify the location of the probe as rapidly and accurately as possible by clicking on a prespecified key on the keyboard, one for the left and one for the right position. Then, another screen with another pair of faces appears and the procedure repeats. A CBM session consists of 360 consecutive

¹²Visual stimuli used in CBM procedures include words, such as in the pioneering work by MacLeod et al. (1986), or faces such as in Eldar and Bar-Haim (2010) which is the protocol that we follow, or See et al. (2009) and Li et al. (2008). Faces are taken from validated databases; we used the Karolinska Directed Emotional Faces (KDEF) database (Lundqvist et al., 1998). Validation guarantees that the expressions portrayed by the actors are accurately recognized by third parties as communicating the intended emotions (Goeleven et al., 2008). The KDEF contains seventy different pairs of faces, half of which use male actors and half female actors.

screens. A session lasts for about 10 minutes. The participants in our study completed a minimum of 16 sessions (one per day). Section 3 on experimental design describes how we ensure compliance throughout the study period. Figure SM.2 in the SM presents an example of the CBM task.

CBM trains participants in the treatment group to avoid threats by presenting the probe always at the location of the neutral face. In the control condition the probe is equally likely to replace the neutral or the angry face. The response latency, i.e., the time that passes from the moment when the probe appears to the moment when the participant clicks on the keyboard to indicate its location, is a measure of attention. Participants have an attention bias if they respond considerably faster to probes replacing the threatening stimulus in comparison with the neutral one.

The efficacy of the CBM treatment in reducing attention bias is supported by a variety of studies. A meta-analysis of CBM in 12 RCTs (Hakamata et al., 2010) found significant improvements in attention bias, with a rather large effect size ($d=1.16$, $CI=(0.82,1.5)$), and in various anxiety measures, with a medium effect size ($d=0.61$, $CI=(0.42, 0.81)$) on average, and a larger effect for individuals who were pathologically anxious ($d=0.78$) than for those who were not ($d=0.48$).

Selected CBM sessions are preceded and followed by 60 screens equivalent to the control condition, i.e. the probe replaces either stimulus with equal frequency. Therefore these longer sessions include a sequence of 60 screens (random probe) + 360 screens (CBM treatment/control condition) + 60 screens (random probe) and are called assessment sessions. These sessions are useful to assess the response latency toward threatening stimuli in the treatment group (which otherwise would not be observed) and therefore validate the efficacy of the CBM treatment over different sessions. Selected days for assessment sessions are the same for all participants, whether in the treatment or control conditions.

3 Experimental Design

In this section we describe our sampling frame, the way in which we administered the CBM treatment and the experimental design.

Our sample comes from a database of students from the University of Lausanne and the Swiss Federal Institute of Technology in Lausanne. A total of 6,000 students received an invitation to complete a questionnaire to measure their trait anxiety, the tendency to experience anxiety in general. We adopt an established questionnaire called the State-

Trait Anxiety Inventory (STAI) developed by Spielberger (1980). The STAI is composed of 40 questions addressed at identifying two dimensions of anxiety: a permanent stable trait (trait anxiety) and a temporary stimulus-induced anxiety state (state anxiety). For either dimension, the STAI produces a scale ranging from 20 to 80 points in which a value above 45 is considered to indicate a highly anxious individual (see Salemink et al., 2009; Yiend et al., 2005).¹³ To increase the variation of trait anxiety in the experimental sample while offering the treatment to people who would potentially gain from it, we defined treatment beneficiaries as those who scored a value of at least 35 points in trait anxiety.

The sample includes the first 300 students with trait anxiety above 35 who accepted our invitation. Owing to no-shows ($n = 7$) and a small rate of attrition, the final longitudinal empirical sample consists of 261 subjects. Attrition is not correlated with treatment, and is balanced in terms of observables; see Table SM.1 in SM.¹⁴

The experiment is based on a longitudinal design with randomization of the treatment at the individual level. Participants took part in a baseline laboratory session that included the first CBM intervention and was an assessment session (i.e., 60 + 360 + 60 screens). The treatment was assigned randomly when participants attended their first laboratory session.¹⁵ Part of the CBM treatment was conducted at home, over a period of three weeks, for a total of 14 CBM completed sessions (Figure 1 illustrates the timeline of the experiment).¹⁶ A CBM session is considered completed if the participant identifies the location of the probe correctly in a minimum of 90% of the screens. We extensively

¹³Typical questions to measure trait anxiety in the STAI ask subjects to indicate which answers among “almost never”, “sometimes”, “often”, “almost always” they feel satisfied with themselves, or how frequently they feel as happy as others seem to be, etc.

¹⁴Out of the 300 students, 7 did not show up to the initial laboratory session and were excluded for the study; 32 (15 treated and 17 control subjects) were dropped from the sample because they either did not turn up at the final laboratory session or they did not comply with the rules of the training.

¹⁵Participants were randomly allocated to computer cubicles in the lab. On the cubicle’s desk, they found the CBM instruction pack with an ID number stapled to the first page of the pack. A range of ID numbers were associated with the control condition and a different range was associated with the treatment condition. Nowhere in the instructions was there a reference to the treatment or control condition.

¹⁶The CBM training was programmed using *Inquisit 4*. The instructions to access the CBM training include four simple steps: opening a webpage, entering a unique ID, downloading and running the application that produces the CBM task, and completing the task. The URL link to access the training and the dates on which the home-sessions had to be completed were printed in the instruction pack given to participants in their first laboratory sessions. Access to all CBM sessions occurred in the same manner, whether the CBM session was conducted in the laboratory or from a personal computer at home. Participants could ask questions in regard to the procedure to access the CBM training in person, during the first laboratory session, and by email during the training period. The home-session completion and click accuracy were monitored.

monitored treatment compliance.¹⁷ The experiment concluded with a follow-up laboratory CBM session that was an assessment session. We incentivized participation with a contribution of CHF 110 (approximately €91) to be paid at the end of the study on condition of full compliance.

The baseline and follow-up laboratory sessions included a cognitive task and were programmed using *zTree* (Fischbacher, 2007)). The task consisted in answering a selection of questions taken from the Raven’s Progressive Matrices booklet (Raven et al., 2003). Each question includes a group of logically related geometric designs with one missing piece that the participant had to identify from some available options. Participants were asked to answer correctly as many questions as possible in 10 minutes. Two different sets of questions were drawn at baseline and follow-up to minimize the possibility of memorizing answers. In the first laboratory session, the cognitive task was done before the CBM treatment, whereas in the follow-up laboratory session, the CBM treatment preceded the cognitive task.

Our measure of performance in the cognitive task is the number of attempted questions within the 10 minutes, regardless of whether the answers are correct or incorrect. This choice rests on the purpose of CBM: restoring attentional control resources that enable participants to be more focused and efficient in answering cognitively demanding questions. CBM is not expected to change cognitive ability and therefore we expect the ratio of the number of correct answers to attempted questions to remain constant. An alternative outcome variable would be the number of correct answers. Whereas attempting a question can be directly linked to cognitive resources being freed up, answering a question correctly is a probabilistic outcome conditional on attempting the question. The probability of a correct answer depends on several individual factors (such as subject’s IQ, familiarity with the questions, etc.) and a (random) luck component. This individual heterogeneity is expected to increase the residual variance of the measurement. We report our results with respect to correct answers in the Supplementary Material.¹⁸


¹⁷To ensure that participants completed the necessary sessions of CBM at home, we sent daily reminder e-mails and called participants on the phone if they missed a CBM session. The CBM sessions had to be completed by 10 p.m. every day. If catching up of a missed session was required or the participant did not meet the required 90% score of correct clicks, then the participant was instructed to conduct the CBM session during the weekend. Noncompliers were eliminated from the experiment. In rare instances, subjects completed more than one CBM session on the same day. In these cases, only the first completed session is included in the analysis. Overall average click accuracy per sessions was 95%.

¹⁸The Raven’s questions appeared sequentially on the computer screen. The score in this task – that is, the number of correct answers – was not directly incentivized; however, participants were informed that (i) the experimenter would calculate their score, and (ii) this score would affect successive tasks and the final earnings. We have no reason to believe that subjects did not aim to answer the task questions correctly. The overall ratio of correct answers to attempted questions is 0.65 (pre-treatment); if subjects

For every CBM screen, we collected the length of time (in milliseconds) taken by the participant to identify the location of the probe on the screen. Typically, CBM studies exclude latencies that are too fast or too slow because these are not indicative of sensible response times given the task. We follow Zvielli et al. (2014) and See et al. (2009) and exclude latencies smaller than 200 ms and greater than 1,500 ms, which results in 2.7% of the latencies observations being dropped.

To measure attention bias at baseline we average (within subject) response latencies to threat and neutral stimuli in the first 60 screens of the first laboratory assessment session (in which the probe replaces threat and neutral stimuli with equal frequency for both control and treatment groups). The attention bias measure is $(\bar{N} - \bar{T})_{\text{pre},i}$, where \bar{N} and \bar{T} are averages within subject over the 60 screens and the subscript “pre” indicates that the measurement is pre-treatment. When $(\bar{N} - \bar{T})_{\text{pre},i} > 0$ attention bias is present.

Figure 1: TIMELINE OF EXPERIMENT



4 Empirical Strategy

To examine the causal effect of the CBM training on attention bias and performance in the cognitive test, we adopt an empirical model explaining pre-post treatment changes in attempted questions (a difference-in-difference design). This allows us to control for unobserved (time-invariant) factors that may influence the outcome variable and might be correlated with the CBM treatment condition, despite the treatment being randomised by design. The baseline model specification is:

had answered questions randomly this ratio would be close to 1/6 since there is one correct answer out of six possible options.

$$y_{it} = \beta_0 + a_i + \beta_1 CBM_{it} + \delta_0 t_{endline} + u_{it}, \quad t = 1, 2. \quad (1)$$

where a_i captures unobserved individual-specific traits; CBM_{it} is an indicator variable representing treatment assignment, that equals 1 if the subject is in the treatment group and 0 if the subject is in the control group. $t_{endline}$ is an indicator variable for the post-treatment period. Finally, u_{it} is a disturbance term. We estimate the treatment effect by estimating the following first-differenced equation, which removes a_i :

$$\Delta y_i = \beta_1 \Delta CBM_i + \delta_0 + \Delta u_i \quad (2)$$

The empirical specification (2) further controls for observed heterogeneity due to age, trait anxiety (and its squared value) and type of study (in the robustness analysis) measured at the pre-treatment stage. The causal interpretation of the impact of CBM relies on the assumption that $\mathbb{E}(\Delta CBM_i \Delta u_i) = 0$, i.e., the change in treatment status is assumed to be uncorrelated with the change in the idiosyncratic errors. The ordinary least squares (OLS) estimator of β_1 gives a consistent estimate of the average treatment effect. We bootstrap standard errors at the level of the individual.

We estimate the model by sex. We split the sample by sex because of biological and neurological differences between males and females which may influence our outcome variables differently. The estimation by sex allows the control variables to yield different effects on the outcome in regard to both sign and magnitude (something that we observe empirically, for example for the effect of age).

5 Results

This section discusses two sets of results: first, we evaluate the efficacy of the CBM treatment in reducing attention bias; second, we evaluate the impact on participant performance in the cognitive test.

There are no systematic differences between treatment and control group in terms of individual characteristics, such as age, program of study, years of study, (self-reported) average grades and trait anxiety between the male and female samples (Tables SM.2 for females and SM.3 for males in SM). T-tests of mean differences in attempted questions show statistical significant differences within genders. These differences are due to a small number of ‘extreme’ observations in the lower tail of the distribution of the outcome variable. The SM provides a robustness analysis on a restricted sample where observations

below the 5th and above the 95th percentile of the outcome distribution are excluded (Table SM.7, which we discuss in subsection 5.2). Attention bias at baseline shows significant heterogeneity among participants: attention bias is observed in 54% of female and male samples, with an average attention bias of 21 ms in females and 20 ms in males (see Table SM.4). This means that our average treatment effect (ATE) estimates are lower bound estimates of the CBM impact, since only about 54% of subjects display an observable attention bias.

In the female sample, we observe a mildly higher baseline $(\bar{N} - \bar{T})_{\text{pre},i}$ in the treatment arm on average, but the difference is not statistically significant. The average number of attempted questions and correct answers in the cognitive test (which are outcome variables) at baseline are moderately higher in the control group. The difference-in-difference empirical strategy will correct for initial differences between treatment arms. Across gender, we observe females having higher levels of trait anxiety than males (p -value < 0.01), in line with the higher prevalence of anxiety in females in the population. Surprisingly, we do not find a strong correlation between the measures of trait anxiety and attention bias. We adopted the STAI test to measure trait anxiety to keep with the existing literature, however future research may wish to investigate more sophisticated and objective measures, such as biomarkers.


5.1 Validation of Treatment Efficacy

Participants in the CBM treatment condition, where the probe always replaces the neutral stimulus, are trained to become faster at locating the neutral stimulus. Participants in the control condition, where the probe randomly replaces either the neutral or the threatening stimulus, are not expected to change their response times, at least not in a different way for either one of the stimuli (since, potentially, they can become faster at identifying the location of both, an indication of possible ‘learning effects’). In the spirit of a validation test, the comparison between response latencies to the *neutral* stimuli in the treatment and control groups over CBM sessions provides a first test of the efficacy of the CBM treatment in reducing the attention bias.

In Figure 2 we compare the *median* response time to neutral stimuli between treatment and control group in each session relative to the initial (i.e., the first CBM session in the laboratory) median response time. Figure 2 shows a significant decrease in response latency in the treatment group over sessions. A large proportion of the decrease occurs within the first four sessions, after which the rate of decrease is slower. Reaction times to neutral stimuli in the control group remain stable, except for some variability in the later

sessions.¹⁹ We show the median response latency since it is less sensitive to outliers and therefore is a more conservative measure than the mean. Figure SM.4 in the SM shows that using the mean reaction times does not produce a different pattern.

Figure 2: MEDIAN RESPONSE TIME TO NEUTRAL FACE RELATIVE TO THE FIRST ASSESSMENT SESSION



Note: The figure shows the median reaction time towards neutral stimuli in each session relative to the initial median reaction time (session 1 in the laboratory) in the treatment group (purple) and control group (orange). The 95% confidence intervals are computed using the between-subjects variation.

A second way to test the efficacy of CBM is to measure the extent to which an entire course of CBM training reduces attention bias in the treatment group relative to the control. To measure and compare the attention bias, we need to observe the reaction times to neutral and threatening stimuli. We can compare the response times to neutral stimuli between treatment and control groups *in each CBM session*, but we do not have observations for threatening stimuli for participants in the treatment group since they always observe the probe replacing the neutral stimuli. However, both treatment and control groups see neutral and threatening stimuli in the assessment sessions. We evaluate the impact of the entire course of CBM training on attention bias by measuring the attention bias in the 60 screens preceding the first CBM session at baseline (i.e., first assessment session) and in the 60 screens following the last CBM session in the follow-up (post-treatment) laboratory session (which completes the three-week-long CBM

¹⁹Figure SM.3 in SM shows that, in the control group, the median reaction times to threat stimuli closely mirror the median reaction times to neutral stimuli across sessions. This evidence speaks against the idea that latencies to neutral or threat stimuli change in a different way over time. This also allays the concern that latencies toward neutral stimuli might be easier to modify relative to latencies to threat stimuli.

treatment) for the treatment and control groups.

We calculate the average change in attention bias, that is $\frac{1}{n} \sum_i [(\bar{N} - \bar{T})_{\text{post},i} - (\bar{N} - \bar{T})_{\text{pre},i}]$ where \bar{N} and \bar{T} are the (average) reaction times (in milliseconds, and the average is taken over the 60 screens within subject) to neutral (N) and threat (T) stimuli and the subscripts ‘post’ and ‘pre’ indicate that the measurement is post-treatment and pre-treatment, respectively. If CBM is effective, one would expect a negative overall difference (since attention bias post-treatment will be smaller than the initial bias).

Table 1 reports the CBM average treatment effect on attention bias on all females and males (Panel: Everyone) and on the sample of female and male participants with pre-treatment attention bias (Panel: Conditional on attention bias). The average treatment impact is a reduction in attention bias of -18 ms for females. There is no evidence of attention bias reduction for males. The CBM impact on subjects with (pre-treatment) attention bias gives an indication of the average CBM treatment effect on the *treated*. The unconditional effects are plotted in Figure 3.²⁰ CBM-treated females with attention bias show a reduction of -23 ms on average (p -value = 0.001), as compared with females with attention bias in the control group. Table SM.6 shows that larger decreases in attention bias occur among female participants who start with a higher pre-treatment attention bias. In this category, CBM reduces approximately 60% of the average attention bias.

5.2 Impacts on Cognitive Performance

Table 2 reports the estimated impact of CBM on performance in the cognitive test, measured as the number of attempted questions as described in Section 3. CBM-treated female participants significantly improve their performance; however, there is no significant impact for males on average, although the coefficient is positive. The magnitude of the impact on females is large: the impact equates to approximately one additional attempted question above an average of 2.7 (equivalent to an approximately 37% increase). These results are consistent with the effects of CBM reported in Section 5.1, which suggest


²⁰One may expect the change in attention bias between post- and pre-treatment to be zero for the control group. While theoretically possible, this does not happen empirically. We observe that for control group participants who have attention bias at baseline, the gap $(\bar{N} - \bar{T})_{\text{post},i}$ in the post-treatment session shrinks relative to its pre-treatment value, so that the overall difference post- and pre-treatment is negative on average. Reduction in attention bias in the control group is a feature reported in other evaluations as well (Eldar and Bar-Haim, 2010). It would be concerning if \bar{N} and \bar{T} in the *control* group went in opposite directions (e.g., with \bar{T} increasing and \bar{N} decreasing, or vice versa). Panel A in Table SM.5 in the SM reports \bar{N} and \bar{T} of control participants with baseline attention bias for each session and shows that the two measures move in the same direction. This is reassuring. A similar pattern is observed in the entire sample (panel B of Table SM.5). Furthermore, we use a difference-in-difference approach, which controls for pre-post-treatment changes in the control group.

Table 1: CBM IMPACT ON CHANGES IN ATTENTION BIAS: EVERYONE AND
CONDITIONAL ON ATTENTION BIAS, N-T>0

Panel: Everyone	Female	Male	Female	Male
CBM	-18.369*** (7.008)	3.968 (6.310)	-16.691** (6.953)	3.910 (6.489)
Age			2.655* (1.485)	1.580 (1.286)
Trait anxiety (pre-treatment)			-2.909 (2.942)	0.618 (2.744)
Trait anxiety (squared)			0.033 (0.033)	-0.004 (0.031)
Constant	3.224 (4.650)	-2.639 (4.934)	6.251 (61.042)	-53.295 (68.566)
adj. R^2	0.053	-0.005	0.066	-0.018
N	129	126	129	125
<hr/>				
Panel: Conditional on Attention bias, N-T>0	Female	Male	Female	Male
CBM	-23.823*** (6.870)	8.389 (8.007)	-22.666*** (7.961)	7.056 (7.763)
Age			-0.098 (1.752)	1.201 (1.671)
Trait anxiety (pre-treatment)			-2.829 (3.249)	-0.208 (2.697)
Trait anxiety (squared)			0.031 (0.036)	0.007 (0.030)
Constant	-14.038*** (4.617)	-23.351*** (5.393)	47.857 (67.347)	-53.408 (68.152)
adj. R^2	0.127	0.004	0.102	-0.022
N	69	68	69	67

Note: The table shows the CBM impact on changes in attention bias between pre and post-treatment (equation 2). Attention bias is measured as the difference in response latency to neutral versus threat stimuli. Regressions in panel: Conditional on Attention bias are estimated on the sample of subjects with pre-treatment positive attention bias. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Figure 3: CHANGE IN ATTENTION BIAS, BY TREATMENT AND SEX



Note: The figure shows the reduction in attention bias pre- and post- treatment for the female and male samples in treatment and control groups. The figure plots the average unconditional changes, $\frac{1}{n} \sum_i^n [(\bar{N} - \bar{T})_{post,i} - (\bar{N} - \bar{T})_{pre,i}]$. For the changes conditional on controls, see Table 1.

that CBM reduces attention bias for treated females but not for treated males.

One explanation for the gender heterogeneity of the treatment impact is that the effects of CBM are more visible at higher levels of anxiety, which females display more than males. Consistent with this interpretation, Table 3 confirms that CBM treatment is more effective on those females scoring baseline anxiety levels higher than the females group's median. The impact on anxious females is three times as large than on less anxious females (p -value < 0.000). As a robustness test, in Table SM.9 in the SM we show that the gender heterogeneity of the treatment impact persists when the number of correct answers in the cognitive test is used as the outcome variable. The average treatment effect on the entire female sample (Panel A) is, however, less precisely estimated due to a larger unobserved heterogeneity in the measurement. These estimates match neatly those in the main analysis: treated anxious females, who attempt more questions after treatment, also increase their score of correct answers by a similar proportion. As cognitive attentional resources are restored by the CBM treatment, anxious (female) participants are able to attempt more questions in a given time and consequently achieve better scores in the cognitive test. The results in Table SM.9 (Panel A) suggest that CBM may not change the performance ranking of the average woman but it may improve the performance ranking of anxious women. Taken together, these results suggest that anxious females are those who benefit most from CBM treatment.

We also conduct a number of robustness tests. First, one might worry that students in

Table 2: CBM IMPACT ON CHANGES IN ATTEMPTED QUESTIONS IN COGNITIVE TEST

	Male	Female	Male	Female
CBM	-0.005 (0.462)	0.995** (0.458)	0.148 (0.460)	1.106** (0.458)
Trait anxiety (pre-treatment)			0.412** (0.180)	-0.476** (0.242)
Trait anxiety (squared)			-0.005** (0.002)	0.005* (0.003)
Age			0.340*** (0.092)	-0.016 (0.116)
Constant	2.841*** (0.290)	2.697*** (0.307)	-12.548*** (4.439)	13.070*** (4.778)
R^2	0.000	0.030	0.110	0.076
N	130	131	130	131
Mean Δ Attempted (control)	2.841	2.696	2.841	2.696

Note: The table shows the CBM impact on (changes in) attempted questions in the Raven’s Matrices test between pre- and post-treatment by gender estimated according to equation 2 in columns 1-2 and adjusted for pre-treatment covariates in columns 3-4. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

science- and mathematics-based study programs²¹ are more accustomed to tests similar to the task used in the laboratory sessions and hence perform better than students from other disciplines. This would bias our results *within* gender if the students in science and mathematics programs were also disproportionately represented in the treatment group. This is not the case for either females or males (see Tables SM.2 for females and SM.3 for males in SM). Controlling for enrollment in a science/mathematics program as an indicator in a robustness test does not change the results (Table SM.10, columns 1 and 2, in SM).²²

Second, we cannot exclude that the length of higher education experience has an impact on performance via other channels – through, for example, “training” students to be more resilient to anxiety – which may interact with the effect of CBM and be correlated with participation in the experiment (but not the treatment allocation which was randomized).²³ When we restrict the sample to first-year students, for whom the effect of education training is minimal since they had been in higher education for about

²¹These include Biology, Medicine, Life Sciences, Geology Architecture, Engineering, Computer Science, Business and Economics, Management of Technology.

²²Note that female participants in humanities programs have a higher average number of attempted questions at baseline compared with females in science- and mathematics-based degrees, while it is the opposite for males. Since it is arguably harder to increase performance when starting from a higher baseline value and females are over-represented in humanities degrees, our main results are likely to be attenuated.

²³We observe a higher number of participants from first-year bachelor students (39%) than from second- and third-year and Master’s students.

Table 3: CBM IMPACT ON CHANGES IN ATTEMPTED QUESTIONS IN COGNITIVE TEST
BY TRAIT ANXIETY AND BY GENDER

Female sample	Low anxiety	High anxiety	Low anxiety	High anxiety
CBM	0.458 (0.606)	1.594** (0.728)	0.499 (0.703)	1.642** (0.780)
Age			-0.098 (0.154)	0.038 (0.187)
Trait anxiety (pre-treatment)			-1.157 (0.953)	0.288 (1.075)
Trait anxiety (squared)			0.015 (0.014)	-0.002 (0.010)
Constant	2.971*** (0.392)	2.406*** (0.471)	26.959 (17.308)	-9.064 (28.713)
R^2	0.007	0.068	0.078	0.121
N	69	62	69	62
Mean Δ Attempted (control)	2.97	2.40	2.97	2.40
Male sample	Low anxiety	High anxiety	Low anxiety	High anxiety
CBM	0.242 (0.585)	-0.248 (0.734)	0.698 (0.767)	0.080 (0.736)
Age			0.255** (0.112)	0.457*** (0.171)
Trait anxiety (pre-treatment)			2.004* (1.079)	-0.706 (0.784)
Trait anxiety (squared)			-0.030 (0.017)	0.006 (0.008)
Constant	2.758*** (0.331)	2.933*** (0.482)	-36.158* (17.340)	12.487 (20.327)
R^2	0.002	0.002	0.143	0.149
N	65	65	65	65
Mean Δ Attempted (control)	2.75	2.93	2.75	2.93

Note: The table shows the CBM impact on (changes in) attempted questions in the Raven's Matrices test between pre- and post-treatment estimated according to equation 2 (adjusted for covariates) by anxious and non-anxious type. A subject is classified in the group of 'High anxiety' if her Trait Anxiety score is higher than the median Trait Anxiety within her gender group, otherwise the subject is classified in the group 'Low anxiety'. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

one month at the start of the CBM training, we still observe an impact on treated females (Table SM.10, columns 3 and 4.).

Several additional robustness tests are presented in the SM. We test the possibility that confounding effects attributable to observations coming from the tails of the outcome distribution (within gender) might be impacting the results. Table SM.7 in the SM reports the treatment effect on a ‘trimmed’ sample from which observations equal or below the 5th percentile and equal or above the 95th percentile of the (pre-treatment) outcome distribution (within gender) are excluded. The cost of trimming is a reduction in statistical power. However, the magnitude of the treatment effect is unaltered across specifications and is very similar to the regressions using the full sample.

Despite the treatment randomisation, one may worry about the influence of unaccounted unobservables. Table SM.8 in the SM presents evidence on how strong the selection on unobservables would need to be in order to reduce the coefficient of the treatment effect to zero using the methods suggested by Altonji et al. (2005) and Oster (2019). Both Altonji’s and Oster’s ratios statistics are negative for both female and male samples. Negative ratios arise when the observable controls are on average negatively correlated with the treatment and positively with the outcome (or vice-versa). The test assumes the unobservables have the same pattern of correlation as the included variables. The tests’ negative ratios suggest that our estimates are attenuated towards zero by unobservable characteristics (i.e. the estimates are downward biased).

To determine whether the impact of CBM training is long-lived and improves actual examination performance, we collected the participants’ examination results in the exam session which took place about three months after the follow-up data collection.²⁴ Examination results have a low variability and we do not observe a significant impact on average grade. However, the negative correlation between trait anxiety and university examination results that exists among females in the control group is absent in the treatment group three months after the training. This suggests that the intervention may generate long-lasting benefits (Table 4).

²⁴On approval by the student faculties of affiliation, we were able to retrieve examination data for 136 subjects (52% of the original sample of participants) directly from faculty administration offices. The data were assembled by a member of staff at the University of Lausanne. The authors never had access to any personal data available to the administration offices that could be used to link the students in our sample to their examination data and do not own the examination data.

Table 4: LONG TERM EFFECTS OF CBM ON CORRELATION
BETWEEN UNIVERSITY EXAM GRADES AND ANXIETY TRAIT SCORES

	Male	Female
Treatment group:		
Correlation	0.031	-0.125
(p-value)	(0.877)	(0.475)
Tracked subjects	27	35
Average exam grades	4.117	4.159
SD of exam grades	1.022	1.219
Control Group:		
Correlation	0.152	-0.339
(p-value)	(0.479)	(0.072)
Tracked subjects	24	29
Average exam grade	4.375	4.446
SD of exam grades	0.87	0.551

Note: The table shows the Pearson’s correlation (corr) between trait anxiety scores and university exam grades (averaged across exam sat) by treatment and control group and by sex (sample of N=136). The correlation’s p-values are reported in brackets. ‘tracked subjects’ indicate the number of subject for whom we were able to receive exam data.

6 Conclusions

Lower performance in cognitive tasks is associated with attention bias towards negative stimuli. Attention bias is associated to anxiety and has a detrimental effect on performance because it diverts cognitive resources away from the completion of a task. We randomized an antianxiety digital treatment, cognitive bias manipulation (CBM), that aims at reducing attention bias. Existing studies focused on pathologically anxious subjects, whereas the participants of our experiment are college students.

The results show that CBM reduces attention bias and improves the cognitive performance of females in the treatment group. The magnitude of the impact is large (37% increase in performance) and is associated to the more anxious females in the sample. There is no significant effect for males. Since according to medical statistics anxiety prevalence is higher in females on average, these results suggest the potential for cognitive bias modification to be an effective way to reduce the detrimental effects of anxiety on cognitive performance and the resulting gender disparities in opportunities.

Future studies may seek to replicate these results and analyse hypotheses that we were unable to test due to lack of data. One hypothesis, suggested by an anonymous referee, concerns a differential reaction in latencies to stimuli when the face shown corresponds with the sex of the respondent. This would be a useful addition to the current results be-

cause, if it is the case that the participant's latency responds better to same-gender faces, then CBM training programs can be easily personalised to increase their effectiveness.

CBM interventions are shorter, cheaper and more easily administered than alternative available treatments, such as cognitive behavioral therapy and medications. As an illustration, a private session of CBT currently costs between £40 and £100, according to NHS statistics, whereas the administration of CBM, once programmed, can run at almost zero marginal costs.

Acknowledgments

We thank participants at the 6th Annual Meeting of the French Experimental Economics Association, the IMEBESS conference in Florence, the Barcelona Graduate Summer Forum and seminars at King's College London, the University of Lausanne and Middlesex University of London.

References

- Alan, Sule, Teodora Boneva, and Seda Ertac**, “Ever failed, try again, succeed better: Results from a randomized educational intervention on grit,” *The Quarterly Journal of Economics*, 2019, *134* (3), 1121–1162.
- Altonji, Joseph G, Todd E Elder, and Christopher R Taber**, “Selection on Observed and Unobserved Variables: Assessing the Effectiveness of Catholic Schools.,” *Journal of Political Economy*, 2005, *113* (1), 151.
- Beard, Courtney**, “Cognitive bias modification for anxiety: current evidence and future directions.,” *Expert review of neurotherapeutics*, 2011, *11* 2, 299–311.
- Blattman, Christopher, Julian C Jamison, and Margaret Sheridan**, “Reducing crime and violence: Experimental evidence from cognitive behavioral therapy in Liberia,” *American Economic Review*, 2017, *107* (4), 1165–1206.
- Borghans, Lex, Angela Lee Duckworth, James Heckman, and Bas ter Weel**, “The Economics and Psychology of Personality Traits,” *Journal of Human Resources*, 2008, *43* (4).
- Cahlíková, Jana, Lubomír Cingl, and Ian Levely**, “How Stress Affects Performance and Competitiveness Across Gender,” *Management Science*, *forthcoming*, 2019.
- Coffman, Katherine Baldiga**, “Gender Differences in Willingness to Guess,” *Management Science*, 2014, *60* (2), 434–448.
- Das-Munshi, Jayati, David Goldberg, Paul E. Bebbington, Dinesh K. Bhugra, Traolach S. Brugha, Michael E. Dewey, Rachel Jenkins, Rob Stewart, and Martin Prince**, “Public health significance of mixed anxiety and depression: beyond current classification,” *British Journal of Psychiatry*, 2008, *192* (3), 171–177.
- Derakshan, Nazanin and Michael W Eysenck**, “Anxiety, processing efficiency, and cognitive performance: New developments from attentional control theory,” *European Psychologist*, 2009, *14* (2), 168–176.
- Eldar, Sharon and Y Bar-Haim**, “Neural plasticity in response to attention training in anxiety,” *Psychological medicine*, 2010, *40* (04), 667–677.
- Eysenck, Michael W and Nazanin Derakshan**, “New perspectives in attentional control theory,” *Personality and Individual Differences*, 2011, *50* (7), 955–960.
- Fischbacher, Urs**, “z-Tree: Zurich toolbox for ready-made economic experiments,” *Experimental Economics*, 2007, *10* (2), 171–178.

- Goeleven, Ellen, Rudi De Raedt, Lemke Leyman, and Bruno Verschuere**, “The Karolinska directed emotional faces: a validation study,” *Cognition and Emotion*, 2008, *22* (6), 1094–1118.
- Guiso, Luigi, Ferdinando Monte, Paola Sapienza, and Luigi Zingales**, “Culture, Gender, and Math,” *Science*, 2008, (5880), 1164–1165.
- Hakamata, Yuko, Shmuel Lissek, Yair Bar-Haim, Jennifer C. Britton, Nathan A. Fox, Ellen Leibenluft, Monique Ernst, and Daniel S. Pine**, “Attention bias modification treatment: a meta-analysis toward the establishment of novel treatment for anxiety,” *Biological psychiatry*, 2010, *68* (11), 982–990.
- Krosnick, Jon A and Duane F Alwin**, “Aging and susceptibility to attitude change,” *Journal of personality and social psychology*, 1989, *57* (3), 416.
- Li, Songwei, Jieqing Tan, Mingyi Qian, and Xinghua Liu**, “Continual training of attentional bias in social anxiety,” *Behaviour research and therapy*, 2008, *46* (8), 905–912.
- Lundqvist, Daniel, Anders Flykt, and A Öhman**, “The Karolinska directed emotional faces (KDEF),” *CD ROM from Department of Clinical Neuroscience, Psychology section, Karolinska Institutet*, 1998, pp. 91–630.
- MacLeod, Colin, Andrew Mathews, and Philip Tata**, “Attentional bias in emotional disorders,” *Journal of abnormal psychology*, 1986, *95* (1), 15.
- McManus, Sally, Paul Bebbington, Rachel Jenkins, and Traolach Brugha**, *Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014*, Leeds: NHS Digital, 2016.
- Moutafi, Joanna, Adrian Furnham, and Ioannis Tsaousis**, “Is the relationship between intelligence and trait Neuroticism mediated by test anxiety?,” *Personality and Individual Differences*, 2006, *40* (3), 587–597.
- Oster, Emily**, “Unobservable selection and coefficient stability: Theory and evidence,” *Journal of Business & Economic Statistics*, 2019, *37* (2), 187–204.
- Paserman, M Daniele**, “Gender differences in performance in competitive environments: evidence from professional tennis players,” *IZA Discussion Paper*, 2007.
- Raven, John, J. C. Raven, and John Hugh Court**, “Manual for Raven’s progressive matrices and vocabulary scales: Section 1: general overview,” 2003.

- Rear, David**, “One size fits all? The limitations of standardised assessment in critical thinking,” *Assessment & Evaluation in Higher Education*, 2019, 44 (5), 664–675.
- Remes, Olivia, Carol Brayne, Rianne Linde, and Louise Lafortune**, “A systematic review of reviews on the prevalence of anxiety disorders in adult populations,” *Brain and Behavior*, 2016, 6 (7), e00497.
- Salemink, Elske, Marcel van den Hout, and Merel Kindt**, “Effects of positive interpretive bias modification in highly anxious individuals,” *Journal of anxiety disorders*, 2009, 23 (5), 676–683.
- See, Jacey, Colin Macleod, and Russell Bridle**, “Attentional Bias: A Real-World Study Using a Home-Based Cognitive Bias Modification Procedure,” *Journal of Abnormal Ps*, 2009, 118 (1), 65 –75.
- Spielberger, Charles D**, *Test anxiety inventory*, Wiley Online Library, 1980.
- Stansfeld, Stephen, Charlotte Clark, Paul Bebbington, Michael King, Rachel Jenkins, and Stephen Hinchliffe**, ‘Chapter 2: Common mental disorders’ in *McManus S, Bebbington P, Jenkins R, Brugha T. (eds), Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014*, Leeds: NHS Digital, 2016.
- Twenge, Jean M., Brittany Gentile, C. Nathan DeWall, Debbie Ma, Katharine Lacefield, and David R. Schurtz**, “Birth cohort increases in psychopathology among young Americans, 1938-2007: A cross-temporal meta-analysis of the MMPI,” *Clinical Psychology Review*, 2010, 30 (2), 145–154.
- Wilson, Edward and Colin MacLeod**, “Contrasting two accounts of anxiety-linked attentional bias: Selective attention to varying levels of stimulus threat intensity,” *Journal of abnormal psychology*, 2003, 112 (2), 212–218.
- Yiend, Jenny, Bundy Mackintosh, and Andrew Mathews**, “Enduring consequences of experimentally induced biases in interpretation,” *Behaviour research and therapy*, 2005, 43 (6), 779–797.
- Zvielli, Ariel, Amit Bernstein, and Ernst H. W. Koster**, “Dynamics of Attentional Bias to Threat in Anxious Adults: Bias towards and/or Away?,” *PLoS ONE*, 2014, (8).

SUPPLEMENTARY MATERIAL


FIGURES

Figure SM.1: EXAMPLE OF A FACE PAIR




Note: From left to right, photo with angry expression and photo with neutral expression. The sample consisted of 70 amateur actors, 35 females and 35 males, of ages between 20 and 30 years taken from the Karolinska Directed Emotional Faces database.

Figure SM.2: SEQUENCES OF SCREENS IN THE TREATMENT CONDITION OF CBM




Note: The two stimuli differing in emotional valence - neutral vs. threat - are presented simultaneously following the centering of the field of vision indicated by a fixation cross. Each image appears randomly on either the left or right side of the screen. Both images (a) and (b) here show that probe appears behind the neutral facial expression, but in different locations.

Figure SM.3: RESPONSE TIME TO THREATENING AND NEUTRAL FACES RELATIVE TO THE FIRST ASSESSMENT SESSION, MEDIAN (CONTROL GROUP)



Note: The figure shows the median reaction time towards threatening (orange) and neutral (blue) stimuli in each session relative to the initial median reaction time (session 1 in laboratory) for the control group. Threatening faces appear only to participants in the control group.

Figure SM.4: RESPONSE TIMES RELATIVE TO THE FIRST ASSESSMENT SESSION, MEAN



Note: The figure shows the mean reaction time towards neutral stimuli in each session relative to the initial mean reaction time (session 1) in the treatment group (purple) and control group (orange). It also shows the mean reaction time towards threatening stimuli in each session relative to the initial mean reaction time (session 1) for the control group (blue). Threatening faces appear only to participants in the control group.

TABLES

Table SM.1: ATTRITION ANALYSIS

CBM	-0.286 (0.465)	-0.385 (0.494)	-0.388 (0.561)
Male		0.710 (0.439)	0.705 (0.485)
Age			-0.015 (0.102)
Constant	-2.220*** (0.274)	-2.633*** (0.403)	-2.298 (2.173)
N	289	288	288
R^2	0.003	0.021	0.021

Note: The table shows logistic regressions where the dependent variable is equal to 1 if the subject was present in the first experimental session but drops from the sample afterwards (and zero otherwise). Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table SM.2: BALANCE TABLE: FEMALE SAMPLE

	(1) TG		(2) Control		(3) Test	
	Mean	S.D.	Mean	S.D.	T test/Chi2#	P-value
N	65		66		131	
Age	20.81	(2.105)	21.13	(2.794)	0.741	0.459
Grade (past year)	4.606	(0.540)	4.537	(0.639)	-0.659	0.510
Trait anxiety (pre-treatment)	43.092	(8.742)	43.681	(9.978)	0.3594	0.719
N-T (baseline)	6.968	(29.18)	-2.916	(32.90)	-1.819	0.071
Attempted questions (pre-treatment)	8.738	(3.649)	10.12	(3.764)	2.134	0.034
Correct answers (pre-treatment)	5.908	(2.199)	6.576	(2.113)	1.773	0.078
Field of studies:						
Humanities	0.475	(0.503)	0.428	(0.498)	0.274#	0.600
Economics/Technical	0.229	(0.424)	0.238	(0.429)	0.012#	0.910
Sciences	0.295	(0.459)	0.333	(0.475)	0.210#	0.647
Year of study:						
BSc year 1	0.415	(0.496)	0.424	(0.498)	0.010	0.918
BSc year 2	0.230	(0.424)	0.181	(0.388)	0.479	0.489
BSc year 3	0.076	(0.268)	0.090	(0.289)	0.083	0.773
MSc year 1	0.200	(0.403)	0.136	(0.345)	0.949	0.330
MSc year 2	0.076	(0.268)	0.121	(0.328)	0.718	0.397
Other years	0.000	(0.000)	0.045	(0.209)	3.023	0.082

Note: The table shows the means and standard deviations (S.D.) of listed variables for the female sample. The final columns report the T-tests (for continuous variables) and Chi-squared tests (for binary variables) for equality of means between treatment arms and the test's p-values. Humanities includes degrees in Literature, Law, Political Sciences; Economics/Technical includes degrees in Business and Economics, Management and Technology; Sciences includes degrees in Biology, Medicine, Life Sciences, Geology, Architecture, Engineering, Computer Science.

Table SM.3: BALANCE TABLE: MALE SAMPLE

	(1) TG		(2) Control		(3) Test	
	Mean	S.D.	Mean	S.D.	T-test/Chi2 \ddagger	P-value
N	67		63		130	
Age	20.74	(2.186)	21.13	(2.587)	0.913	0.362
Grade (past year)	4.659	(0.560)	4.568	(0.518)	-0.963	0.337
Trait anxiety (pre-treatment)	41.552	(8.733)	39.301	(8.329)	-1.501	0.135
N-T (baseline)	1.187	(26.926)	1.628	(28.006)	0.091	0.927
Attempted questions (pre-treatment)	9.575	(3.258)	8.190	(2.401)	-2.386	0.0071
Correct answers (pre-treatment)	6.462	(2.30)	6.206	(1.676)	-0.721	0.472
Field of studies:						
Humanities	0.129	(0.337)	0.193	(0.398)	0.953 \ddagger	0.329
Economics/Technical	0.403	(0.494)	0.354	(0.482)	0.308 \ddagger	0.579
Science	0.467	(0.503)	0.451	(0.501)	0.032 \ddagger	0.857
Year of study:						
BSc year 1	0.318	(0.469)	0.412	(0.496)	1.243	0.265
BSc year 2	0.257	(0.440)	0.142	(0.352)	22.635	0.104
BSc year 3	0.181	(0.388)	0.126	(0.335)	0.739	0.390
MSc year 1	0.166	(0.375)	0.158	(0.368)	0.0149	0.903
MSc year 2	0.030	(0.172)	0.126	(0.335)	0.4213	0.040
Other years	0.045	(0.209)	0.0317	(0.176)	0.162	0.687

Note: The table shows the means and standard deviations (S.D.) of listed variables for the male sample. The final columns report the T-tests (for continuous variables) and Chi-squared tests (for binary variables) for equality of means between treatment arms and the test's p-values. Humanities includes degrees in Literature, Law, Political Sciences; Economics/Technical includes degrees in Business and Economics, Management and Technology; Sciences includes degrees in Biology, Medicine, Life Sciences, Geology, Architecture, Engineering, Computer Science.

Table SM.4: DESCRIPTIVE STATISTICS BY SEX

	Females	Males	Total	T-Test/Chi2 \ddagger
Age	20.977	20.908	20.943	0.230
Grade (past year)	4.572	4.615	4.593	-0.622
N-T (pre-treatment)	1.989	1.401	1.696	0.161
Percentage of subjects with (N-T)>0	54.200	53.850	54.020	0.057
Attention bias (ms)	21.090	20.390	20.743	0.232
Trait anxiety (pre-treatment)	43.389	40.462	41.931	2.634***
Attempted questions (pre-treatment)	9.435	8.885	9.161	1.318
Correct answers (pre-treatment)	6.244	6.338	6.291	-0.362
Field of study:				
Humanities	0.452	0.160	0.305	5.246***
Economics/Technical	0.233	0.376	0.014	-2.454**
Sciences	0.314	0.464	0.389	-2.437**

Note: The table shows the means of listed variables for the entire sample by gender. The final columns report the T-tests (for continuous variables) and Chi-squared tests (for binary variables) for equality of means between genders and the conventional stars symbol for test significance. Humanities includes degrees in Literature, Law, Political Sciences; Economics/Technical includes degrees in Business and Economics, Management and Technology; Sciences includes degrees in Biology, Medicine, Life Sciences, Geology, Architecture, Engineering, Computer Science.

Table SM.5: CONTROL GROUP'S REACTION TIMES BY SESSION

Panel A: Subjects in the Control Group with Attention Bias at Baseline, N-T>0

		CBM sessions																Post-treatment	
		Pre-treatment	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15		16
Female	N	371.4	368.3	366.2	363.2	365.6	380.2	369.4	370.8	374.8	374.8	375.6	365.8	370.8	382.9	402.6	378.5	346.2	340.7
	T	356.3	369.1	368.0	362.6	363.2	383.6	367.7	370.5	372.7	371.3	376.3	363.1	371.4	382.5	402.8	376.7	344.0	339.1
	N-T	15.1	-0.8	-1.8	0.6	2.3	-3.4	1.8	0.3	2.1	3.5	-0.8	2.7	-0.6	0.4	-0.2	1.8	2.2	1.6
Male	N	386.6	361.7	370.8	367.2	366.6	372.3	369.7	376.9	372.9	381.2	373.1	374.4	375.1	377.9	391.1	393.9	338.1	338.9
	T	366.1	357.9	370.2	365.4	368.2	367.2	370.4	374.9	371.7	379.5	371.3	371.9	379.9	376.9	390.3	390.2	339.5	343.4
	N-T	20.5	3.8	0.6	1.9	-1.6	5.1	-0.7	2.1	1.2	1.7	1.8	2.5	-4.7	0.9	0.9	3.6	-1.4	-4.4

Panel B: Everyone in the Control Group

		CBM sessions																Post-treatment	
		Pre-treatment	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15		16
Female	N	378.9	378.0	371.4	370.1	369.9	379.0	372.0	375.6	377.7	376.8	377.1	373.1	379.4	384.6	395.0	379.2	349.8	345.1
	T	381.8	378.1	371.3	369.1	368.4	380.5	371.3	374.8	376.8	374.5	375.2	372.8	379.5	383.2	392.8	378.4	348.0	343.0
	N-T	-2.9	-0.1	0.1	1.0	1.5	-1.5	0.7	0.8	0.9	2.3	1.9	0.2	-0.1	1.4	2.2	0.9	1.8	2.1
Male	N	368.8	359.4	371.6	370.3	370.9	375.1	377.9	382.6	378.7	386.7	377.4	374.7	382.3	382.2	389.8	390.5	336.8	333.3
	T	367.1	356.8	370.0	368.6	371.0	372.2	375.6	379.3	378.1	385.3	375.2	371.5	386.5	381.5	389.8	389.6	337.4	335.2
	N-T	1.6	2.6	1.6	1.7	-0.1	2.8	2.3	3.3	0.6	1.4	2.2	3.2	-4.1	0.7	-0.1	0.9	-0.6	-1.9

Note: The table shows the control group's reaction times to neutral (N) and to threat (T) stimuli in the pre-treatment session, each of the CBM sessions, the post-treatment session.

Note: The table shows the control group's reaction times to neutral (N) and to threat (T) stimuli in the pre-treatment session, each of the CBM sessions, the post-treatment session.

Table SM.6: CBM IMPACT ON CHANGES IN ATTENTION BIAS BY SEVERITY OF
(PRE-TREATMENT) ATTENTION BIAS: FEMALE SAMPLE

Female sample	Conditional on Attention Bias, N-T>0			
	Low bias	High bias	Low bias	High bias
CBM	-8.167 (7.511)	-20.837** (9.465)	-7.401 (8.221)	-19.149* (11.241)
Age			-1.269 (1.705)	1.265 (2.023)
Trait anxiety (pre-treatment)			-1.499 (2.889)	-4.229 (6.154)
Trait anxiety (squared)			0.017 (0.032)	0.046 (0.068)
Pre-treatment				
Mean Attention Bias (ms)	7.65	33.43	7.65	33.43
R^2	0.043	0.105	0.091	0.141
N	33	36	33	36

Note: The table shows the CBM impact on changes in attention bias between pre- and post-treatment (equation 2) by severity of attention bias at baseline in the female sample only. Attention bias is measured as the difference in response latency to neutral versus threat stimuli. The sample includes females who display positive (pre-treatment) attention bias. ‘Low bias’ indicates pre-treatment attention bias lower than the median values; ‘high bias’ indicates attention bias higher than the median value. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table SM.7: ROBUSTNESS: CBM IMPACT ON PRE-POST TREATMENT CHANGES IN
ATTEMPTED QUESTIONS, TRIMMED SAMPLE

	(1) Male	(2) Female	(3) Male	(4) Female
CBM	-0.055 (0.515)	0.982* (0.564)	0.189 (0.481)	0.929* (0.514)
Age	0.312*** (0.097)	-0.001 (0.139)	0.287*** (0.098)	-0.033 (0.136)
Trait anxiety (pre-treatment)	0.409* (0.219)	-0.352* (0.198)	0.381** (0.193)	-0.320* (0.194)
Trait anxiety (squared)	-0.005** (0.002)	0.004* (0.002)	-0.005** (0.002)	0.003 (0.002)
Attempted questions (pre-treatment)			-0.396*** (0.140)	-0.337*** (0.096)
Constant	-11.706** (5.040)	10.319** (4.216)	-7.506 (4.819)	13.591*** (4.064)
Adj. R^2	0.079	0.013	0.158	0.104
N	109	106	109	106

Note: The table shows the CBM impact on changes in attempted questions in the Raven’s Matrices test by gender (equation 2 in the manuscript) in a trimmed sample where observations below or equal to the 5th and above or equal the 95th percentile have been dropped. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table SM.8: OMITTED VARIABLE BIAS: OSTER (2019)/ ALTONJI (2005) RATIOS

	Control set	Oster (2019) bias-adjusted test	Oster (2019) bounds for β	Altonji, Elder, Taber (2005) test
Female	as in Table 2	[<0]	[1.10, 3.77]	[<0]
Male	as in Table 2	[<0]	[0.14, 1.59]	[<0]

Note: The table reports the degree of selection on unobservables relative to observables which is required to produce a treatment effect of zero, using the methods of Altonji et al. (2005) and Oster (2019) (the δ in their papers). The entry [<0] indicates that the respective ratios are negative. This is because the observable controls are on average positively correlated with the outcome variable and negatively with the treatment, suggesting a downward bias in the OLS estimates (provided the unobservables have similar correlation patterns with the outcome and the treatment as the included observables). The control variables included are those presented in Table 2 of the paper. The *bounds* for β are calculated according to Oster (2019), page 18, using the conservative assumption that the maximum explainable variation is 1 ($R_{max} = 1$) and $\delta = 1$.

Table SM.9: ROBUSTNESS: CBM IMPACT ON PRE-POST TREATMENT CHANGES IN CORRECT ANSWERS

	(1)	(2)	(3)
Panel A: Female			
	All	Low anxiety	High anxiety
CBM	0.595*	-0.564	1.646***
	(0.412)	(0.550)	(0.619)
Trait anxiety (pre-treatment)	-0.169	-0.692	1.009
	(0.161)	(0.899)	(0.860)
Trait anxiety (squared)	0.002	0.008	-0.009
	(0.002)	(0.013)	(0.008)
Age	-0.080	-0.132	-0.066
	(0.087)	(0.150)	(0.124)
Constant	7.722**	19.167	-24.547
	(3.751)	(15.935)	(23.424)
Adj. R^2	0.005	0.035	0.094
N	131	69	62
Panel B: Male			
	All	Low anxiety	High anxiety
CBM	-0.187	0.019	-0.453
	(0.382)	(0.485)	(0.447)
Trait anxiety (pre-treatment)	0.194	0.124	0.441
	(0.162)	(1.017)	(0.481)
Trait anxiety (squared)	-0.002	-0.001	-0.005
	(0.002)	(0.015)	(0.005)
Age	-0.078	-0.003	-0.154
	(0.061)	(0.112)	(0.105)
Constant	-0.217	-1.050	-4.842
	(4.169)	(16.642)	(12.039)
Adj. R^2	0.002	0.053	0.024
N	130	65	65

Note: The table shows the CBM impact on changes in correct answers among female participants (Panel A) and male participants (Panel B). A subject is classified in the group of ‘High anxiety’ if their trait anxiety score is higher than the median trait anxiety within their gender group, otherwise the subject is classified in the group ‘Low anxiety’. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. The stars indicate the statistical significance of the one-sided test $H_0: \beta_{CBM} \leq 0$, $H_A: \beta_{CBM} > 0$ using the convention * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table SM.10: ROBUSTNESS: CBM IMPACT ON PRE-POST TREATMENT CHANGES IN ATTEMPTED QUESTIONS, DEGREE TYPES AND FIRST YEAR OF STUDIES

	Male	Female	Male First Year	Female First Year
CBM	0.159 (0.499)	1.133** (0.500)	-0.099 (0.731)	2.092** (0.861)
Age	0.292*** (0.111)	-0.011 (0.131)	0.329* (0.179)	0.389 (0.465)
Trait anxiety (pre-treatment)	0.384** (0.188)	-0.453** (0.210)	0.844** (0.355)	-0.391 (0.341)
Trait anxiety (squared)	-0.005** (0.002)	0.005** (0.002)	-0.010** (0.004)	0.005 (0.004)
Science degree (=1)	-1.257** (0.603)	0.529 (0.550)		
Constant	-9.988** (4.934)	12.100** (4.853)	-20.491*** (7.232)	2.530 (10.019)
R^2	0.138	0.084	0.260	0.145
N	130	131	48	55
Mean Δ Attempted (control)	2.84	2.69	3.42	2.57
Mean Attempted at baseline (Science degree)	8.97	8.97		
Mean Attempted at baseline (Humanities)	8.52	9.93		

Note: The table shows the CBM impact on changes in attempted questions in the Raven's Matrices test between pre- and post-treatment by gender estimated according to equation (2). The estimation sample of columns 3 and 4 is restricted to first year bachelor students. 'Science degree' indicates enrollment into a quantitative study program, including Biology, Medicine, Life Sciences, Geology, Architecture, Engineering, Computer Science, Business and Economics, Management and Technology. Estimations are based on OLS regressions. Standard errors are bootstrapped with 150 replications and reported in brackets. Levels of significance: * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$.

Table SM.11: MEAN VALUE OF ATTENTION BIAS, ATTEMPTED QUESTIONS AND CORRECT ANSWERS: PRE-TREATMENT, POST-TREATMENT AND CHANGE

	Attention bias (mean)			Attempted questions (mean)			Correct answers (mean)		
	(1) Pre- treatment	(2) Post- treatment	(3) Change (post-pre)	(1) Pre- treatment	(2) Post- treatment	(3) Change (post-pre)	(1) Pre- treatment	(2) Post- treatment	(3) Change (post-pre)
Panel A: Female									
Treatment group	6.969	-8.176	-15.145	8.738	12.431	3.692	5.908	8.431	2.523
Control group	-2.916	-0.352	3.224	10.121	12.818	2.697	6.576	8.500	1.924
Difference-in-Difference			-18.369			0.995			0.599
Panel B: Male									
Treatment group	1.188	1.718	1.329	9.537	12.373	2.836	6.463	8.075	1.612
Control group	1.629	-0.859	-2.640	8.190	11.032	2.841	6.206	8.000	1.794
Difference-in-Difference			3.969			-0.005			-0.182

EXPERIMENTAL INSTRUCTIONS

ENGLISH INSTRUCTIONS: CBM or CONTROL TREATMENT

Instructions for the experiment on stress reduction (LAGRA)

Welcome to this experiment!

You were selected to participate on the basis of the questionnaire that you have submitted online.



N.B. YOU CANNOT PARTICIPATE TO THIS EXPERIMENT IF YOU USE UNIX ON YOUR COMPUTER. YOU NEED A COMPUTER THAT EXECUTES WINDOWS OR MAC.

You have to complete a task that lasts between 10 and 13 minutes. You have to repeat this task at home on every working day (from Monday to Friday) for a total of 14 days, before you come back to the lab (see date at the end of the document). You must follow some strict rules during these 14 days; the rules are explained here below. Now, we begin by giving you a description of the task:

(1) THE TASK

-First: you will see three fixation crosses (+++) in the middle of the screen. Look at them, please.

-Second: two photos will briefly appear, one to the left and one to the right of the centre of the screen.



-Third: an « X » will appear in the position of one of the two photos.

YOUR TASK:

- If you see that the « X » appears in the position previously taken by the photo on the left, press the key F of your keyboard **as fast as possible**.
- If you see that the « X » appears in the position previously taken by the photo on the right, press the key J of your keyboard **as fast as possible**.

N.B.: the key F is located to the left of the key J – this should make the task intuitive.

→ It's an easy task!

(2) IMPORTANT RULES TO FOLLOW DURING THE 14 SESSIONS AT HOME

It is important to follow these rules.

It is required to complete the daily task every working day (from Monday to Friday) for a total of 14 times. The task lasts between 10 and 13 minutes.

- (a) Try to complete the task in a very quiet environment, with no distractions.
- (b) You need to have started the task by 22h every day (e.g. : it is acceptable if you start the task at 21h59, and finish around 22h10. But it is not acceptable to start at 22h15.)
- (c) As well as the task described in part (1), the task described here above repeats, and you have to complete the sequence of tasks without interruptions (i.e. do not let the task linger around while you do other things – we can detect this behavior from the results).

- (d) Even if the task looks repetitive, you must focus and try to identify the correct location of the « X » each time.
- (e) ***If you do not complete the task every day, you will receive an email from us to remind you to catch up during the week-end. If you do not complete the tasks 14 times before your second session in the lab, you will not receive any payment.***

(3) HOW TO ACCESS THE TASK FROM HOME


We will have seen the stages together in the first session in the lab. Nevertheless, we include some detailed instructions here below, step by step. Section A applies to those who use Windows/PC. Section B applies to those who use a Mac.

SECTION A : Windows

Step 1: Make sure that the date and time of your computer are up to date.


Step 2: Use **INTERNET EXPLORER** to open this webpage:

<http://research.millisecond.com/marklambiris/FINAL.web>



Step 3: Press « Start »; enter your identifier (this is the number that you received on the day of your first session in the lab). Enter the identifier again when we ask you to confirm it. You have to use the same identifier every day.

Step 4: Press « Install ».



After that, you will see the screen below for 1 or 2 minutes, during this time the program is loading. Then, just follow the instructions to complete the task.




SECTION B : Mac

Step 1: Make sure that the date and time of your computer are up to date.

Step 2: Use **SAFARI** to open this webpage:

<http://research.millisecond.com/marklambiris/FINAL.web>



Step 3: Press « Start »; enter your identifier (this is the number that you received on the day of your first session in the lab). Enter the identifier again when we ask you to confirm it. You have to use the same identifier every day.

Step 4: Press « Downloads », as indicated here below. You should see the file **Inquisit.jnlp**. Double-click the file.



Step 5: You will see the screen here below. Press « Open »:



Step 6: You will see the screen here below. Press « Run »:




After that, you will see the screen below for 1 or 2 minutes, during this time the program is loading. Then, just follow the instructions to complete the task.




(4) PROBLEMS WITH THE PROGRAM

Problems with Mac: When I reach step 4 or 5 described above, I see the following screen:




Solution:


Step 1: In the main screen, click the Apple symbol located in the top left, and select « System Preferences », as indicated here below:




Step 2: Click « Security and privacy », as indicated here below:




Step 3: In the section « General », find « Allow apps downloaded from: ». Select the option « Anywhere ». If this is not allowed, click the padlock symbol, as indicated here below:



Enter your password. Now you can select « Anywhere », and save the changes.






ENGLISH INSTRUCTIONS: COGNITIVE TASK


IQ test: The next task is composed of several questions. We will now distribute leaflets with example questions. You will see 14 symbols and an empty square. Underneath you will see six symbols. You will have to choose the one which is most appropriate to fill in the empty square above. Look at the examples and the explanations of the logic underlying the correct choice; this will help you for the task. If you have questions, please signal the assistants. Click on the button when you are ready. [When everyone is ready]: We will now start the test. In 10 minutes the task will automatically end. You have 10 minutes to complete the task.

Leaflet with example questions:




Correct Answer: B

The logic underlying the question is to be found in each line independently. In each line, there is a specific symbol that increases in size from left to right.




Correct Answer: A

In each line, the number of bars increases by one unit in each following box (from left to right). The number of points increases by one unit until the column in the middle, and then decreases moving to the right. Therefore, the points are symmetrically distributed around the middle row.



Correct Answer: F

The logic underlying of the question is to be found in the diagonals running from top left to bottom right. Each diagonal comprises the same symbol in the same color.



Correct Answer: A

The logic of the question is to be found in the arrows: moving from the left to the right in each row, the arrow turns by 45 degree clockwise.

FRENCH INSTRUCTIONS: CBM or CONTROL TREATMENT

Instructions pour l'expérience sur la réduction du stress (LAGRA)


Bienvenue à cette expérience!

Vous avez été sélectionné pour y participer suite à votre questionnaire soumis en ligne. **N.B. VOUS NE POUVEZ PAS PARTICIPER A CETTE EXPERIENCE SI VOUS UTILISER UNIX SUR VOTRE ORDINATEUR. IL FAUT ABSOLUMENT UN ORDINATEUR QUI EXECUTE WINDOWS, OU UN MAC.**

Vous allez effectuer une tâche, qui dure entre 10 et 13 minutes. Vous devrez répéter la tâche, chez vous, chaque jour ouvrable (de lundi à vendredi) pendant un total de 14 jours, avant de revenir au laboratoire (voir dates à la fin du document). Vous devrez adhérer à des critères stricts pendant ces 14 jours ; cela va être expliqué ci-dessous. Mais dans un premier temps, on commence avec un descriptif de la tâche:

(1) LA TACHE

- Dans un premier temps: vous verrez trois fixations (+++) au centre de l'écran. Fixez votre regard là-dessus SVP.
- Dans un second temps: deux photos apparaîtront très brièvement: une à gauche et une à droite du centre de l'écran.



- Dans un troisième temps: un « X » apparaîtra dans une des deux positions, précédemment occupée par une photo.

VOTRE TÂCHE:

- si vous voyez le « X » apparaître dans la position précédemment occupée par la photo à gauche du centre, appuyez sur la touche F de votre clavier **le plus rapidement possible**.
- si vous voyez le « X » apparaître dans la position précédemment occupée par la photo à droite du centre, appuyez sur la touche J de votre clavier **le plus rapidement possible**.

N.B.: la touche F est placée à gauche de la touche J sur le clavier - cela devra rendre la tâche intuitive.

→ Ce n'est pas compliqué !

(2) REGLES IMPORTANTES A SUIVRE PENDANT LES 14 SESSIONS, A DOMICILE

Les règles suivantes sont très importantes à suivre.

- (a) Il faut compléter la tâche quotidienne chaque jour ouvrable (de lundi à vendredi) pour un total de 14 fois. La tâche dure entre 10 et 13 minutes.
- (b) Essayez de compléter la tâche dans un endroit aussi calme que possible, sans distractions.

- (c) Il faut jouer à la tâche jusqu'à 22h chaque jour (ex: si vous le commencez à 21h59, vous le terminerez vers 22h10 : ceci est acceptable. Mais le commencer à 22h15 n'est pas acceptable).
- (d) Comme la tâche décrite dans la partie (1) ci-dessus se répète successivement, il faut absolument compléter la tâche en une fois (c'est-à-dire ne pas laisser traîner la tâche pendant longtemps tandis que vous faites autre chose – nous pourrons le voir dans les résultats).
- (e) Même si la tâche semble répétitive, il faut être concentré et essayer d'identifier la position correcte du « X » à chaque tour.
- (f) ***Si vous n'effectuez pas la tâche un jour, vous recevrez un e-mail de notre part pour vous rappeler de la rattraper en fin de semaine. Si vous n'effectuez pas la tâche 14 fois jusqu'à votre seconde sessions au laboratoire, aucun dédommagement ne vous sera versé.***

(3) COMMENT ACCEDER A LA TACHE A DOMICILE


Nous aurions déjà vu les étapes ensemble dès la première session au laboratoire. Mais on inclut des instructions détaillées ci-dessous, en étapes. La section A est pour les utilisateur de Windows/PC. La section B est pour ceux qui utilisent un Mac.

SECTION A : Windows

Etape 1 : (a) Vérifiez que la date et l'heure de votre ordinateur sont à jour


Etape 2 : Utiliser **INTERNET EXPLORER** pour accéder à la page web :

<http://research.millisecond.com/marklambiris/FINAL.web>



Etape 3 : Cliquez sur « Start » ; entrez votre identifiant (ceci est le numéro que vous avez reçu le jour de votre session au laboratoire). Entrez-le de nouveau quand on vous demande de le confirmer. Il faudra utiliser le même identifiant chaque jour.

Etape 4 : Cliquez sur « Install » (Installer).



Par la suite vous verrez l'écran ci-dessous pendant 1 ou 2 minutes, le temps de préparer le programme. Après il suffit de suivre les instructions et effectuer la tâche.



SECTION B: Mac

Etape 1: Vérifiez que la date et l'heure de votre ordinateur sont à jour.

Etape 2: Utiliser **SAFARI** pour accéder à la page web :

<http://research.millisecond.com/marklambiris/FINAL.web>




Etape 3: Cliquez sur « Start » ; entrez votre identifiant (ceci est le numéro que vous avez reçu le jour de votre session au laboratoire). Entrez-le de nouveau quand on vous demande de le confirmer. Il faudra utiliser le même identifiant chaque jour.

Etape 4: Cliquez sur « Downloads » (Téléchargements), comme indiqué ci-dessous. Vous devriez voir le fichier **Inquisit.jnlp**
Faites un double-clic dessus.



Etape 5: Vous verrez l'écran ci-dessous. Cliquez sur « Open » (Ouvrir) :



Etape 6: Vous verrez l'écran ci-dessous. Cliquez sur « Run » :




Par la suite vous verrez l'écran ci-dessous pendant 1 ou 2 minutes, le temps de préparer le programme. Après il suffit de suivre les instructions et jouer à la tâche.




(4) PROBLEMES D'ACCES AU LOGICIEL

Problème sur Mac : Quand j'arrive à l'étape 4 ou 5 ci-dessus je vois l'écran ci-dessous :




Solution:

Etape 1: Sur l'écran principal, cliquez sur le logo Apple en haut à gauche, et sélectionnez « System Preferences », comme indiqué ci-dessous :




Etape 2: Cliquez sur « Security and privacy », comme indiqué ci-dessous :



Etape 3: Dans la section « General », regardez la rubrique « Allow apps downloaded from : ». Idéalement il faut sélectionner l'option « Anywhere ». Donc si ce n'est pas le cas, comme indiqué ci-dessous, cliquez sur le cadenas :



Entrez votre mot de passe. Vous pouvez maintenant sélectionner « Anywhere », et conserver les changements.




FRENCH INSTRUCTIONS: COGNITIVE TASK

IQ test: La tâche suivante est composée de plusieurs questions. Nous vous distribuons maintenant des questions d'exemples. Vous verrez quatorze symboles et un carré vide. Au-dessous vous verrez six symboles et vous devez choisir celui qui convient le mieux. Regardez les exemples et les explications de la logique derrière ça, cela vous aide après. Cliquez le bouton quand vous êtes prêt.

[Quand ils seront prêts]


Nous commençons maintenant. Après 10 minutes, le questionnaire sera terminé automatiquement. Vous avez 10 minutes pour répondre.

Leaflet with example questions:





Réponse correcte: B

La logique du problème réside ici dans les lignes prises indépendamment. Dans chaque ligne, il y a un symbole spécifique qui grossit de gauche à droite.




Réponse correcte: A

Dans chaque ligne, le nombre de barres augmente de une unité de case en case. Le nombre de points augmente jusqu'à la colonne du milieu et diminue ensuite. Donc, on peut considérer que les points sont distribués de façon symétrique de part et d'autre de la colonne centrale.



Réponse correcte: F

La logique du problème réside dans les diagonales allant d'en haut à gauche jusqu'en bas à droite. Chaque diagonale comporte un même symbole de même couleur.



Réponse correcte: A

La logique réside dans les lignes, de gauche à droite : la flèche tourne à chaque case de 45 degrés dans le sens des aiguilles d'une montre.