Selank Available for Research Use Only

(300 – 900 mcg daily, internasal spray, cycled)

Selank is a synthetic tuftsin-derived heptapeptide with the amino-acid sequence Thr-Lys-Pro-Arg-Pro-Gly-Pro. It was developed in Russia for intranasal administration as an anxiolytic and nootropic peptide. It is used clinically in Russia by intranasal administration to reduce anxiety and produce mild antiasthenic and nootropic effects. Benefits for stress resilience, attention, and working memory are likely.

💉 💉 Probably safe for short courses. Long-term safety is plausible.

⭐ ⭐ Probably effective for anxiety.

⭐ ⭐ Likely effective for cognitive support and stress resilience.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 13-Oct-26. The most recent version of this document is available at https://pdfs.researcher6076.com/pdfs/selank.pdf

Short-term safety is likely based on the favorable tolerability reported in small controlled human trials. Long-term safety is plausible based on Selank's peptide biology, short-term clinical tolerability, and supportive animal evidence.

Dosing Patterns

Intranasal Selank produces acute central effects via direct transport to the central nervous system intranasal application. Community reports commonly describe subjective effects within 15–60 minutes.

Selank dosing is goal-dependent. Common dosing frequencies and cycle lengths are practice-derived combining Russian clinical practice and speculative community practice.

- **Situational stress:** 150–300 mcg per administration, one to three times daily, for a total of 150–900 mcg daily.
- **Daily anxiolysis:** 150–300 mcg per administration, twice daily, for a total of 300–600 mcg daily.
- **Cognitive support:** 150–450 mcg per administration, once or twice daily, for a total of 150–900 mcg daily.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Selank Available for Research Use Only
(300 – 900 mcg daily, internasal spray, cycled)

Safety signals exist from a 2015 clinical study where Selank was administered intranasally at 2.7 mg daily for 14 days, with some participants continuing through day 21.

Benefits

- **Anxiety reduction:** Selank reduces anxiety without clinically significant sedation.
- **Improved stress resilience:** Selank likely reduces behavioral and cognitive disruption during stress.
- **Cognitive support:** Selank likely supports attention, learning, and memory, particularly under stress.
- **Neurotrophic signaling:** Selank likely increases hippocampal BDNF expression. Resulting support for synaptic plasticity is plausible.
- **Immune modulation:** Selank modulates cytokine and immune signaling.
- **Antiviral activity:** Selank likely has antiviral activity against influenza viruses.
- **Enkephalin signaling:** Selank plausibly prolongs endogenous enkephalin activity by inhibiting enkephalin-degrading enzymes.
- **Community reports:** Improved calm, focus, and sleep are widely reported.

Potential Applications

- **Anxiety disorders:** Selank reduces anxiety symptoms in generalized anxiety disorder and in studied phobic-anxiety and somatoform disorders.
- **Situational anxiety:** Selank likely reduces acute situational anxiety.
- **Stress-related cognitive impairment:** Selank likely supports attention, learning, and memory under stress.
- **Post-stroke recovery:** Selank likely supports neurological recovery after stroke.

Cautions

- **Pregnancy and breastfeeding:** Use is precautionary because safety during pregnancy and lactation is not established.
- **Unstable psychiatric conditions:** Use caution in psychosis or bipolar mania because Selank's central nervous system effects have not been adequately studied in these conditions.
- **CNS depressants and psychoactive polypharmacy:** Interactions are poorly characterized, so concurrent use warrants caution.

Side Effects

- **Nasal irritation:** Intranasal administration can cause temporary burning or irritation.
- **Sleep effects:** Community reports describe sleep disturbance or vivid dreams, particularly with late dosing.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- **Activation:** Community reports describe occasional temporary anxiety or activation.

Biological Mechanism

Selank is a tuftsin-derived heptapeptide with multimodal effects. After intranasal administration, direct transport to the central nervous system through olfactory or trigeminal pathways is plausible. Selank plausibly enhances GABAergic signaling through positive allosteric modulation and prolongs endogenous enkephalin activity by inhibiting enkephalin-degrading enzymes. It likely increases hippocampal BDNF expression and modulates monoamine metabolism. It also modulates cytokine signaling. Together, these effects plausibly support synaptic plasticity, neuroprotection, and regulation of anxiety and stress responses.

Stacking Selank and Semax

Combining Selank and Semax is widely reported in community practice. Because the peptides influence partly distinct but overlapping central nervous system pathways, complementary effects on anxiety, stress resilience, attention, and learning are plausible. Additive activation, sleep disturbance, or mood effects are also plausible. Specific combined doses, sequencing strategies, alternating-day schedules, and cycling patterns remain speculative.

References

1. Zozulia AA, Neznamov GG, Siuniakov TS, et al. Efficacy and possible mechanisms of action of a new peptide anxiolytic Selank in the therapy of generalized anxiety disorders and neurasthenia. Zh Nevrol Psikhiatr Im S S Korsakova. 2008;108(4):38-48. PMID: 18454096.
2. Medvedev VE, Tereshchenko ON, Israelian AY, et al. A comparison of the anxiolytic effect and tolerability of Selank and phenazepam in the treatment of anxiety disorders. Zh Nevrol Psikhiatr Im S S Korsakova. 2014;114(7):17-22. PMID: 25176261.
3. Medvedev VE, Tereshchenko ON, Kost NV, et al. Optimization of the treatment of anxiety disorders with Selank. Zh Nevrol Psikhiatr Im S S Korsakova. 2015;115(6):33-40. doi:10.17116/jnevro20151156133-40.
4. Panikratova YR, Lebedeva IS, Sokolov OY, et al. Functional connectomic approach to studying Selank and Semax effects. Dokl Biol Sci. 2020;490(1):9-11. doi:10.1134/S001249662001007X.
5. Uchakina ON, Uchakin PN, Miasoedov NF, et al. Immunomodulatory effects of Selank in patients with anxiety-asthenic disorders. Zh Nevrol Psikhiatr Im S S Korsakova. 2008;108(5):71-75. PMID: 18577961.
6. Zozulya AA, Kost NV, Sokolov OY, et al. Inhibitory effect of Selank on enkephalin-degrading enzymes as a possible mechanism of its anxiolytic activity. Bull Exp Biol Med. 2001;131(4):315-317. doi:10.1023/A:1017979514274.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

7. Zolotarev YA, Dolotov OV, Andreeva LA, et al. Selank and short-lived enkephalins: effects on their degradation in human plasma. Russ J Bioorg Chem. 2004;30(3):207-212. doi:10.1023/B:RUBI.0000030126.09208.C3.
8. Inozemtseva LS, Karpenko EA, Dolotov OV, et al. Intranasal administration of the peptide Selank regulates BDNF expression in the rat hippocampus in vivo. Dokl Biol Sci. 2008;421:241-243. doi:10.1134/S0012496608040066.
9. Volkova AV, Shadrina MI, Kolomin TA, et al. Selank administration affects the expression of some genes involved in GABAergic neurotransmission. Front Pharmacol. 2016;7:31. doi:10.3389/fphar.2016.00031.
10. Vyunova TV, Andreeva LA, Shevchenko KV, Myasoedov NF. Peptide-based anxiolytics: molecular aspects of Selank biological activity. Protein Pept Lett. 2018;25(10):914-923. doi:10.2174/0929866525666180925144642.
11. Volkova AV, Shadrina MI, Kolomin TA, et al. GABA, Selank, and olanzapine affect the expression of genes involved in GABAergic neurotransmission in IMR-32 cells. Front Pharmacol. 2017;8:89. doi:10.3389/fphar.2017.00089.
12. Semenova TP, Kozlovskaya MM, Zakharova NM, et al. Experimental optimization of learning and memory processes by Selank. Eksp Klin Farmakol. 2010;73(8):2-5. PMID: 20919548.
13. Kolik LG, Nadorova AV, Antipova TA, et al. Selank, peptide analogue of tuftsin, protects against ethanol-induced memory impairment by regulating BDNF content in the hippocampus and prefrontal cortex in rats. Bull Exp Biol Med. 2019;167(5):641-644. doi:10.1007/s10517-019-04588-9.
14. Ershov FI, Narovlyansky AN, Mezentseva MV, et al. Antiviral activity of the peptide Selank in experimental influenza infection. Vopr Virusol. 2009;54(5):19-24. PMID: 19882898.
15. Yasenyavskaya AL, Samotrueva MA, Bashkina OA, et al. Effect of Selank on cytokine profile parameters under chronic social stress conditions. Curr Rev Clin Exp Pharmacol. 2021;16(2):162-167. doi:10.2174/1574884715666200704152810.
16. Konstantinopolsky MA, Chernyakova IV, Kolik LG. Selank attenuates behavioral signs of morphine withdrawal in rats. Bull Exp Biol Med. 2022;173(6):730-733. doi:10.1007/s10517-022-05624-x.
17. Sokolov OY, Kost NV, Andreeva LA, et al. Selank and its metabolites alter behavioral phenotype and enkephalinase activity in mice. Bull Exp Biol Med. 2002;133(2):133-135. doi:10.1023/A:1015582302311.
18. Filippenkov IB, Glazova NY, Sebentsova EA, et al. Selank peptide causes changes in gene expression in the hippocampus of rats in the early hours after acute restraint stress. Nanobiotechnology Reports. 2024;19(3):381-392. doi:10.1134/S2635167624601335.
19. Skvortsova VI, Tvorogova TV, Dubina AI, et al. Experimental hemorrhagic stroke: the study of neuropeptides MIF and Selank administered intraperitoneally. Zh Nevrol Psikhiatr Im S S Korsakova. 2009;109(12 Suppl 2):62-66. PMID: 20873405.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

20. Filippenkov IB, Shpetko YY, Stavchansky VV, et al. Transcriptomic effects of peptide drug Selank during early experimental cerebral ischemia in rats. Mol Biol. 2026;60:530-542. doi:10.1134/S002689332670007X.
21. U.S. Food and Drug Administration. Certain bulk drug substances for use in compounding that may present significant safety risks: Selank acetate. Updated September 27, 2024. Accessed September 30, 2026. https://www.fda.gov/drugs/human-drug-compounding/certain-bulk-drug-substances-use-compounding-may-present-significant-safety-risks.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076