Thymulin & Thymosin alpha-1 for Healthspan

(Thymulin 1.0 mg 5 days/week, plus Tα1 1.0 mg 2 days/week)

Improved responses to vaccines and new pathogens, immune recovery after stress, and reduced inflammatory tissue injury are plausible healthspan objectives of combining Zn-thymulin with thymosin alpha-1 (Tα1). Thymulin links thymic endocrine function to T-cell regulation and, in experimental aging models, suppression of inflammatory myeloid signaling; Tα1 modulates dendritic-cell, innate and adaptive responses. The combination is a mechanistically plausible healthspan hypothesis informed by component studies. [1–8,13–15]

💉 💉 Tα1 has reassuring tolerability findings over studied disease-treatment courses; thymulin has a smaller human evidence base that includes serious events of uncertain causality. Longer-term safety of the proposed is unestablished. [9,15,18–24,29]

⭐ Plausibly effective for healthspan extension. The biological rationale is meaningful but unevenly supported. Thymulin has a close relationship to thymic endocrine aging and experimental evidence in age-associated myeloid inflammation. [3–8] Tα1 improved monocyte HLA-DR in the smaller ETASS sepsis trial. [23,24]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Aging Healthspan and the Immune System

The aging immune system is characterized by reduced production of naïve T cells, changes in T-cell repertoire and differentiation, altered innate-cell function, and impaired responses to some infections and vaccines. Thymic involution contributes by reducing the environment available for new T-cell development. Immune aging also involves peripheral tissues, chronic antigen exposure, metabolism, and the inflammatory environment. [1]

Inflammaging is the persistent, low-grade inflammation associated with aging. Senescent cells, tissue injury, changes in the microbiome, chronic infections, and altered innate signaling all contribute. In aging, reduced protective immunity and excessive inflammation often coexist: an older immune system responds poorly to a new pathogen while maintaining inappropriate background inflammatory activity. **The objective of this protocol is to improve regulation and useful responses**, not to generally increase immune activation. [1,2]

Thymic endocrine decline supplies an additional rationale. A thymus-derived circulating signal can influence cells outside the gland. Administering such a signal could potentially affect peripheral immune regulation despite substantial thymic involution. [3–8]

Better vaccine responses, fewer clinically important infections, improved recovery from illness, and reduced inflammatory tissue injury are plausible intermediate objectives. Sustained preservation of function and freedom from disease would be healthspan outcomes. [1,2,6,13–15]

Thymulin Biology and Evidence for Supporting Aging

Thymulin is a thymic epithelial-cell-derived peptide. The zinc-free peptide is often called facteur thymique sérique (FTS); its zinc-bound form, Zn-thymulin, has the characteristic biological activity described in classical thymulin assays. [3] *(In this survey, "thymulin" is the general name used in the literature. "Zinc-free FTS" and "Zn-thymulin" distinguish between the two when that difference matters.)*

Thymulin Human Observations and Zinc interventions

In a crossover study of 44 older adults, 20 mg/day of supplemental zinc partially restored serum thymulin activity. The study supports a contribution from zinc status to diminished activity in that population. [5] The replacement-like hypothesis is that aging can reduce an endocrine signal involved in immune regulation and that exogenous active thymulin might compensate for part of that loss.

Thymulin Aging-specific Experimental Findings

Kanemaru 2026 examined age-associated inflammatory myeloid-cell activity using human immune-cell experiments and interventions in aged mice. Ex vivo thymulin treatment reduced cytokine-producing myeloid cells from older donors. Mouse macrophage experiments implicated canonical NF-κB signaling through reduced p65 DNA binding and IκBα phosphorylation. In aged tumor-bearing mice, thymulin plus zinc improved tumor control and responsiveness to anti-PD-L1 treatment relative to zinc-containing vehicle. The tumor regimen is summarized below as an illustrative HED. [6]

The human-cell component strengthens relevance to human aging biology but remains an ex vivo intervention. The animal component provides causal intervention evidence in the models tested. [6]

Thymulin Peripheral and Intrathymic Actions

Zn-thymulin restored impaired NK activity in cells from old mice in vitro; zinc-free FTS and zinc alone did not reproduce that result. [7] In hydrocortisone-treated aged mice, five days

of Zn-thymulin modestly improved subsequent splenocyte responses, but not thymocyte responses. [8]

These findings suggest administered thymulin has effects beyond T-cell maturation inside the thymus. Peripheral T-cell, NK-cell and myeloid regulation are relevant to the plausible support thymulin might offer for aging. [3,4,6–8]

Tα1 Biology and Evidence for Supporting Aging

Tα1 has effects on innate and adaptive immunity. Experimental work identifies dendritic-cell and Toll-like-receptor-associated signaling, including MyD88-dependent pathways. Other work shows induction of dendritic-cell tryptophan catabolism through indoleamine 2,3-dioxygenase, contributing to a regulatory environment. These observations explain why it is simplistic to describe Tα1 as just a generic immune stimulant. [13,14]

Tα1's relevant proposed actions include antigen presentation, T-cell responsiveness, NK-cell function and regulation of inflammatory responses. These involve peripheral cells as well as pathways associated with thymic immunity, with effects dependent on the experimental or disease context. [13,14,17,28]

Human evidence relevant to older adults

A double-blind trial in 90 men aged 65–99 years evaluated Tα1 with influenza vaccination, using approximately 1.6 mg subcutaneously twice weekly for eight doses. This is direct human intervention evidence in older adults, but its scope is vaccine antibody response. [15]

Human hepatitis, cancer and critical-illness studies provide additional information about exposure and disease-specific outcomes. They have different baseline immune abnormalities, concomitant treatments and therapeutic objectives. [18–24]

The human evidence for Tα1 is broader than for thymulin, while its application to healthspan remains a plausible extrapolation from immune biology and limited older-adult intervention research. [9,13–15,18–24,29]

Thymulin Dose and Frequency Evidence

Animal Regimens and Illustrative Human Equivalent Doses

A study of inflammatory mice with positive inflammatory and signaling outcomes used about HED 8.3 mg on days 1 and 5 of an eight-day experiment. [12] A study of aged tumor-bearing mice found improved tumor control and anti-PD-L1 responsiveness used about HED 8.3 mg daily for four weeks. [6]

Human treatment experience

In humans, two six-month randomized rheumatoid-arthritis trials evaluated subcutaneous thymulin combined with zinc at doses of 1, 5, and 10 mg. The 1 mg and 5 mg regimens used daily injections for two weeks, followed by three injections weekly. The 5 mg regimen produced significant improvements in overall clinical assessment and four objective clinical measures. Differences in dosing schedules, placebo responses, and baseline immune characteristics prevent establishing an optimal dose or concluding that higher doses reduce efficacy. [9] Translation of these anti-inflammatory and functional benefits to a healthspan rationale remains plausible.

Interpretation of the proposed 1.0 mg daily dose

The proposed 1.0 mg daily dose is a speculative, conservative selection informed by experimental activity and the precedent for frequent, continuous administration. Human thymulin PK is insufficiently characterized to optimize dose and interval. Frequent small pulses remain a mechanistic preference rather than an exposure-matched translation of the animal studies. [6,8–12,26]

Tα1 PK

PK and the Limits of the Pulse Model

In a nine-volunteer crossover study, Tα1 given subcutaneously 1.6 mg HED reached peak concentrations in approximately one to two hours. The elimination half-life was less than three hours, and single-dose and five-day multiple-dose exposures showed no apparent accumulation. Formulation affected exposure. These data support transient circulating exposure, rather than sustained concentrations throughout a several-day dosing interval. [16]

A short plasma residence time is not the same as cessation of biological activity. Gene expression and immune-cell responses can outlast measurable circulating peptide. Short PK does not establish the optimal interval or prove that three injections are better than two. [13,14,16,22]

Concentration response and clinical dose comparisons

An experiment in normal human lymphocytes found a bimodal concentration-response pattern, with activity around 10^{-12} M and 10^{-8} M and variability between donors, but this is not a universal human therapeutic curve or informative of a subcutaneous dose. [17]

The Arase pilot randomized 16 HBV patients to 0.8 or 1.6 mg using the same schedule. The larger Iino trial enrolled 316 HBV patients and reported broadly similar overall efficacy

between doses, with a better response to 1.6 mg in an advanced-fibrosis subgroup that was not stratified at randomization. These findings do not establish dose equivalence or select a healthspan dose. [18,19]

In metastatic melanoma, a randomized study compared Tα1 doses of 1.6, 3.2, and 6.4 mg with chemotherapy-based combinations. Outcomes did not establish a simple proportional improvement with increasing dose. [21]

A review describes 8- and 16-mg doses twice weekly for four weeks as well tolerated in healthy volunteers. This is review-level evidence about tolerability under those conditions, not evidence of added healthspan benefit. [28]

Study and status	***Tα1 dose and schedule***	***Duration***	***Relevance and limitation***
Arase 2003 [18]; randomized HBV pilot	0.8 or 1.6 mg, 6 times weekly for 2 weeks, then twice weekly for 22 weeks	24 weeks of treatment	Direct 0.8-mg exposure; frequent dosing was limited to induction
Iino 2005 [19]; randomized HBV trial	0.8 or 1.6 mg, same induction and maintenance pattern	24 weeks of treatment	Larger human dose comparison; not an healthspan trial
Rodríguez-Torres 2008 [20]; published secondary HCV analysis	Methods list 0.8, 1.6, or 3.2 mg or placebo 3 times weekly with peginterferon	Planned course of 48 weeks	Specific 0.8-mg schedule precedent; thyroid-focused report does not establish dose-specific efficacy or completion for every participant
Gravenstein 1989 [15]; randomized vaccine trial	900 micrograms/m^2 subcutaneously twice weekly	8 doses	Older-adult intervention; different dose basis and outcome
ETASS 2013 [23]; randomized sepsis trial	1.6 mg every 12 hours for 5 days, then daily for 2 days	7 days	Acute intensive exposure; not long-term healthspan evidence
TESTS 2025 [24]; phase 3 sepsis trial	1.6 mg every 12 hours	Up to 7 days	No clear reduction in primary mortality outcome; short-term safety context
Qiu 2015 [27]; published HCC trial protocol	1.6 mg twice weekly after resection	Planned 12 months	Describes intended treatment; a protocol alone supplies no completed-outcome evidence

In early advanced-cancer research, immune-assay changes after a single intramuscular Tα1 dose often emerged after 24–48 hours and seldom persisted beyond 72 hours, motivating the authors' proposal for treatment every two to three days. Responses were not

uniformly favorable, and no tumor responses were observed. This supplies a mechanistic timing hypothesis for the proposed schedule. [22]

Mechanistic Rationale for Healthspan

The combination hypothesis is that thymulin could address diminished thymic endocrine signaling and maladaptive peripheral inflammatory activity, while Tα1 could support antigen presentation and coordination of innate and adaptive responses. Both may affect inflammation and T-cell or NK-cell function. [3–8,13,14,17]

The proposed benefits are improved responses to new antigens or vaccines, reduced clinically important infection burden, better immune recovery after stress and less damaging chronic inflammatory activity. Improved surveillance of abnormal cells is another mechanistic possibility. [6–8,10,13–15]

Proposed Healthspan Dose

The following subcutaneous (SC) amounts and patterns are speculative experimental proposals for healthspan research. [9,16,18–22]

- **Tα1 1.0 mg SC two times weekly**
- **Thymulin 1.0 mg SC five times weekly**

No loading phase, dose escalation, fixed one-day sequencing rule or drug holiday is selected by the component studies for this combination. Animal experiments and disease-treatment schedules do not determine a human healthspan course. [6,9,12,18–24]

Plausible Healthspan Benefits

- **Plausibly improved adaptive responses to new pathogens and vaccines:** Tα1 supports antigen-presentation and T-cell pathways in experimental systems, while Zn-thymulin's role in T-cell regulation could complement these effects. [3,4,13–15,17]
- **Plausibly reduced susceptibility to or severity of some infections:** Complementary effects on T-cell regulation, antigen presentation and immune-cell responses could support pathogen clearance and limit infection-related inflammatory damage. Clinical sepsis findings are mixed and context-specific. [13,14,23,24]
- **Plausibly improved immune surveillance of abnormal cells:** Complementary effects on antigen presentation and immune-cell function could support recognition and elimination of abnormal cells; thymulin enhanced antitumor immunity in aged mouse tumor models. [6,13,14,21]

- **Plausibly reduced age-related chronic inflammation and associated tissue damage:** Zn-thymulin and Tα1 could complement one another in regulating inflammatory signaling and immune responses. [6,11–14]
- **Plausible support for immune recovery after acute physiological stress:** Complementary effects on T-cell regulation and peripheral immune responses could help restore disrupted immune function. [8,10,13,23]

Drivers of Diminished Response and Mitigation Strategies

Tα1 has been administered over weeks and months in clinical studies, but these trials generally did not measure responsiveness to each successive dose. Persistence of a clinical or immune endpoint does not by itself establish absence of desensitization. The twelve-month HCC regimen in reference [27] is a planned protocol, not completed tolerability or adaptation evidence.

For thymulin, biological effects were observed after repeated administration in animal studies over days to weeks. Older human trials add six-month treatment experience. Demonstrating an endpoint effect is weaker evidence about adaptation than measuring responsiveness to successive doses.

Available evidence does not show clinically important progressive desensitization for either proposed component, but it is insufficient to exclude it. [6,8-12,15,18–24,27,29]

Zinc Supplementation

Zinc is an essential cofactor for thymulin's classical biological activity and contributes to immune-cell function. [3,4,25] Adequate zinc availability could plausibly support formation of active Zn-thymulin after zinc-free thymulin administration, although formulation and binding conditions matter. [3,4]

Older adults may have marginal zinc intake, making dietary assessment and correction of deficiency a logical adjunct to immune support. The US RDA is 11 mg/day for adult men and 8 mg/day for adult women; the adult upper intake level is 40 mg/day of elemental zinc from all sources. Excessive chronic intake can impair copper status and cause hematologic, neurological and immune problems. [25] Serum copper declined even in the older-adult study using 20 mg/day supplemental zinc, reinforcing attention to zinc–copper balance. [5]

Serum or plasma zinc can help assess status but varies with illness, collection timing and other factors. Dietary history and clinical context are important; copper assessment may be relevant when excess zinc exposure is suspected. [5,25]

References

[1] Nikolich-Žugich J. The twilight of immunity: emerging concepts in aging of the immune system. *Nat Immunol*. 2018;19(1):10–19. doi:10.1038/s41590-017-0006-x.

[2] Franceschi C, Bonafè M, Valensin S, et al. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci*. 2000;908:244–254. doi:10.1111/j.1749-6632.2000.tb06651.x.

[3] Dardenne M, Pléau JM, Nabarra B, et al. Contribution of zinc and other metals to the biological activity of the serum thymic factor. *Proc Natl Acad Sci U S A*. 1982;79(17):5370–5373. doi:10.1073/pnas.79.17.5370.

[4] Prasad AS, Meftah S, Abdallah J, et al. Serum thymulin in human zinc deficiency. *J Clin Invest*. 1988;82(4):1202–1210. doi:10.1172/JCI113717.

[5] Boukaïba N, Flament C, Acher S, et al. A physiological amount of zinc supplementation: effects on nutritional, lipid, and thymic status in an elderly population. *Am J Clin Nutr*. 1993;57(4):566–572. doi:10.1093/ajcn/57.4.566.

[6] Kanemaru H, Luong S, Yamamoto Y, Mizukami Y, Ito F. Thymulin restrains age-associated myeloid inflammation and enhances cancer immunotherapy. *Nat Commun*. 2026;17:6534. doi:10.1038/s41467-026-75383-0.

[7] Muzzioli M, Mocchegiani E, Bressani N, Bevilacqua P, Fabris N. In vitro restoration by thymulin of NK activity of cells from old mice. *Int J Immunopharmacol*. 1992;14(1):57–61. doi:10.1016/0192-0561(92)90105-T.

[8] Saha AR, Hadden EM, Hadden JW. Zinc induces thymulin secretion from human thymic epithelial cells in vitro and augments splenocyte and thymocyte responses in vivo. *Int J Immunopharmacol*. 1995;17(9):729–733. doi:10.1016/0192-0561(95)00061-6.

[9] Amor B, Dougados M, Mery C, Dardenne M, Bach JF. Nonathymulin in rheumatoid arthritis: two double blind, placebo controlled trials. *Ann Rheum Dis*. 1987;46(7):549–554. doi:10.1136/ard.46.7.549.

[10] Okamoto M, Morishita M, Setoguchi C, Nakata K. Restorative effect of short term administration of thymulin on thymus-dependent antibody production in restraint-stressed mice. *Int J Immunopharmacol*. 1993;15(6):757–762. doi:10.1016/0192-0561(93)90149-S.

[11] Henriques-Coelho T, Oliveira SM, Moura RS, et al. Thymulin inhibits monocrotaline-induced pulmonary hypertension modulating interleukin-6 expression and suppressing p38 pathway. *Endocrinology*. 2008;149(9):4367–4373. doi:10.1210/en.2008-0018.

[12] Novoselova EG, Lunin SM, Glushkova OV, et al. Thymulin, free or bound to PBCA nanoparticles, protects mice against chronic septic inflammation. *PLoS One*. 2018;13(5):e0197601. doi:10.1371/journal.pone.0197601.

[13] Romani L, Bistoni F, Gaziano R, et al. Thymosin alpha 1 activates dendritic cells for antifungal Th1 resistance through toll-like receptor signaling. *Blood*. 2004;103(11):4232–4239. doi:10.1182/blood-2003-11-4036.

[14] Romani L, Bistoni F, Perruccio K, et al. Thymosin alpha1 activates dendritic cell tryptophan catabolism and establishes a regulatory environment for balance of inflammation and tolerance. *Blood*. 2006;108(7):2265–2274. doi:10.1182/blood-2006-02-004762.

[15] Gravenstein S, Duthie EH, Miller BA, et al. Augmentation of influenza antibody response in elderly men by thymosin alpha one. A double-blind placebo-controlled clinical study. *J Am Geriatr Soc*. 1989;37(1):1–8. doi:10.1111/j.1532-5415.1989.tb01561.x.

[16] Rost KL, Wierich W, Masayuki F, Tuthill CW, Horwitz DL, Herrmann WM. Pharmacokinetics of thymosin alpha1 after subcutaneous injection of three different formulations in healthy volunteers. *Int J Clin Pharmacol Ther*. 1999;37(1):51–57. PMID:10027483.

[17] Leichtling KD, Serrate SA, Sztein MB. Thymosin alpha 1 modulates the expression of high affinity interleukin-2 receptors on normal human lymphocytes. *Int J Immunopharmacol*. 1990;12(1):19–29. doi:10.1016/0192-0561(90)90064-T.

[18] Arase Y, Tsubota A, Suzuki Y, et al. A pilot study of thymosin alpha1 therapy for chronic hepatitis B patients. *Intern Med*. 2003;42(10):941–946. doi:10.2169/internalmedicine.42.941.

[19] Iino S, Toyota J, Kumada H, et al. The efficacy and safety of thymosin alpha-1 in Japanese patients with chronic hepatitis B; results from a randomized clinical trial. *J Viral Hepat*. 2005;12(3):300–306. doi:10.1111/j.1365-2893.2005.00633.x.

[20] Rodríguez-Torres M, Ríos-Bedoya CF, Ortiz-Lasanta G, Marxuach-Cuétara AM, Jiménez-Rivera J. Thyroid dysfunction (TD) among chronic hepatitis C patients with mild and severe hepatic fibrosis. *Ann Hepatol*. 2008;7(1):72–77. https://www.medigraphic.com/pdfs/hepato/ah-2008/ah081k.pdf

[21] Maio M, Mackiewicz A, Testori A, et al. Large randomized study of thymosin alpha 1, interferon alfa, or both in combination with dacarbazine in patients with metastatic melanoma. *J Clin Oncol*. 2010;28(10):1780–1787. doi:10.1200/JCO.2009.25.5208.

[22] Dillman RO, Beauregard JC, Zavanelli MI, Halliburton BL, Wormsley S, Royston I. In vivo immune restoration in advanced cancer patients after administration of thymosin fraction 5 or thymosin alpha 1. *J Biol Response Mod*. 1983;2(2):139–149. PMID:6315897.

[23] Wu J, Zhou L, Liu J, et al. The efficacy of thymosin alpha 1 for severe sepsis (ETASS): a multicenter, single-blind, randomized and controlled trial. *Crit Care*. 2013;17(1):R8. doi:10.1186/cc11932.

[24] Wu J, Pei F, Zhou L, et al. The efficacy and safety of thymosin α1 for sepsis (TESTS): multicentre, double blinded, randomised, placebo controlled, phase 3 trial. *BMJ*. 2025;388:e082583. doi:10.1136/bmj-2024-082583. Correction: BMJ. 2025;389:r1098. doi:10.1136/bmj.r1098.

[25] National Institutes of Health Office of Dietary Supplements. Zinc fact sheet for health professionals. Accessed November 10, 2026. https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/

[26] US Food and Drug Administration. Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers. Guidance for Industry. July 2005. https://www.fda.gov/media/72309/download

[27] Qiu SJ, Zhou ZG, Shen F, et al. A multicenter, randomized, observation-controlled clinical trial to evaluate the efficacy and safety of thymalfasin adjuvant therapy in patients with HBV-related HCC after curative resection—first announcement of the protocol. *Expert Opin Biol Ther*. 2015;15(suppl 1):S133–S137. doi:10.1517/14712598.2015.1039979.

[28] Garaci E, Paci M, Matteucci C, Costantini C, Puccetti P, Romani L. Phenotypic drug discovery: a case for thymosin alpha-1. *Front Med (Lausanne)*. 2024;11:1388959. doi:10.3389/fmed.2024.1388959.

[29] Roullet E, Cesaro P, Simon-Lavoine N, Degos JD, Marteau R. Nonathymulin treatment of multiple sclerosis: double-blind pilot study. *Acta Neurol Scand*. 1989;80(6):575–578. doi:10.1111/j.1600-0404.1989.tb03929.x.

[30] Negrev N, Hristosov K, Hachmeryan A, Nyagolov Y. Thymus peptides (thymulin, thymosin alpha 1 and thymosin beta 4) inhibiting effects on the intrinsic blood coagulation pathway in rats. *J of IMAB*. 2017;23(3):1641–1645. doi:10.5272/jimab.2017233.1641.