Thymulin

Healthspan: 1.0 mg daily; Acute infection, esp aged populations: 5.0 daily limited to several weeks

Thymulin is an endogenous epithelial-cell-derived zinc-dependent thymic hormone with actions on thymic and peripheral immune cells. It. It is investigated for restoration of age-associated immune dysfunction, inflammatory regulation and potential adjunctive immune support. Its healthspan rationale draws on declining endogenous activity, zinc biology, and experiments in aged human cells and mice. Specific binding sites have been demonstrated in selected T-cell lines, although a complete molecular receptor model remains unresolved. [5,9,12,19,23,24,31]

💉 Human studies provide limited condition-specific exposure data; they do not establish safety of long-term use. Interactions with immunosuppressants and biologics remain insufficiently characterized. [1–3]

⭐ ⭐ Likely immune and inflammatory support relevant to healthspan; [19,22–25] Human research links reduced endogenous active thymulin with aging, malnutrition, and zinc deficiency. The July 2026 aging experiments strengthen the rationale for immune regulation and healthspan. [9,19,24]

⭐ Plausible adjunctive immune support during acute infections [19,22,27,29]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Zn-thymulin vs Zinc-free FTS

Zinc-free FTS denotes the uncomplexed peptide; a product labeled “thymulin” should not be assumed to contain preformed Zn-thymulin. [5,20] Thymulin biological activity requires one bound zinc ion per peptide molecule. In old-mouse spleen cells, Zn-thymulin restored impaired NK-cell activity in vitro, whereas zinc-free FTS and separately zinc alone did not. [20,22]

Aging, Immunosenescence, and Inflammaging

Active thymulin declines with age, with contributions from thymic involution, reduced peptide production, and altered zinc availability or saturation. Adding zinc to serum or restoring deficient zinc status can recover part of the activity. Thus, replacing an age-depleted regulatory signal is a particularly direct aging hypothesis. [9,23,24]

Kanemaru et al., published July 21, 2026 in Nature Communications, found an age-associated inflammatory myeloid phenotype in humans and mice. Thymulin treatment of human peripheral blood mononuclear cells (PBMCs) ex vivo for 24 hours reduced inflammatory cytokine expression in an age-dependent manner, including samples from adults aged 52–86. [19]

In aged mice, thymulin suppressed inflammatory myeloid-cell activity through canonical NF-κB inhibition and enhanced T-cell IFN-γ responses and cytotoxic function. In aged tumor-bearing mice it improved tumor control and response to anti-PD-L1 treatment, with much smaller effects in young animals. The short immune experiments used daily dosing for one week; tumor experiments extended daily exposure to four weeks. These findings support a restorative, age-dependent immune hypothesis; the survival endpoint was studied in tumor-bearing mice. [19]

Older experiments support peripheral immune restoration: Zn-thymulin restored NK cytotoxicity in old-mouse cells in vitro, and a five-day course in hydrocortisone-treated aged mice modestly augmented splenocyte responses, without improving thymocyte responses. The effects depend on immune context and are not evidence of thymic regrowth. [22,27]

Experimental Dose Range and Frequency

The most relevant animal doses and protocols for a healthspan rationale include an HED of 1.1 µg daily for 21 days, which prevented inflammatory cardiopulmonary damage [28], and an HED of 8.3 mg administered twice over eight days, which reduced experimentally induced inflammation, preserved immune cells, and partially restored disrupted neuroendocrine signaling. [29] The most directly relevant aging study used an HED of 8.3 mg daily, reducing age-associated inflammatory signaling after seven days and improving antitumor immune competence, tumor control, and survival in aged tumor-bearing mice treated for 28 days. [19] Importantly, these benefits were observed primarily in aged mice, with little or no corresponding improvement in young mice. [19]

An unresolved issue is the approximately 7,500-fold difference between the HED doses of 1.1 µg and 8.3 mg.

In humans, two six-month randomized rheumatoid-arthritis trials evaluated subcutaneous thymulin combined with zinc at doses of 1, 5, and 10 mg. The 1 mg and 5 mg regimens used daily injections for two weeks, followed by three injections weekly. The 5 mg regimen produced significant improvements in overall clinical assessment and four objective clinical measures. Differences in dosing schedules, placebo responses, and baseline immune characteristics prevent establishing an optimal dose or concluding that higher doses reduce efficacy. [1] Translation of these anti-inflammatory and functional benefits to a healthspan rationale remains plausible.

Dose & Patterns

The proposed 1.0 mg daily dose is a speculative, conservative selection informed by experimental activity and the precedent for frequent, continuous administration. Human thymulin PK is insufficiently characterized to optimize dose and interval. Comparing widely separated animal experiments and limited human trials does not define a human therapeutic window. [1,19,26–30]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Thymulin

Healthspan: 1.0 mg daily; Acute infection, esp aged populations: 5.0 daily limited to several weeks

Speculative experimental pattern for immune support and healthspan:

- **Thymulin 1.0 mg daily for immune support / aging / healthspan**

Small, repeated doses could plausibly restore an age-depleted regulatory signal. Daily administration has precedent; continued administration is a mechanistic hypothesis, while endogenous origin alone does not establish the safety of chronic exogenous exposure. [19,23,28]

A dose for acute infection is informed by likely benefits in in T-cell maturation and function. In **infection accompanied by immune exhaustion, restoring these functions could theoretically improve pathogen clearance**. The rationale may be stronger in older patients, because aging is associated with lower thymulin and impaired cellular immunity. [5,8,19,22,27–29] The dose size is informed the RA trials where 5.0 daily was well tolerated. [1]

- **Thymulin 5.0 mg daily for immune support with acute infections, esp senior populations**

Combining Thymulin and Thymosin Alpha-1

Thymulin and thymosin alpha-1 (Tα1) are potentially complementary rather than redundant. Tα1 has substantially stronger human clinical, dose, PK, and safety evidence and broadly modulates innate/adaptive immune coordination. Thymulin has weaker human evidence but a more direct relationship to age-related thymic endocrine decline, zinc biology, and inflammatory regulation. [5,9,19,23,35]

A combined plausible hypothesis would pair Tα1's effects on antigen presentation and immune responsiveness with thymulin's potential restoration of deficient T/NK-cell function and restraint of excessive myeloid inflammation. Both peptides also engage regulatory pathways; their roles overlap and should not be divided into exclusive "stimulation" and "suppression" functions. [19,22,35]

The combination of Tα1 and thymulin is mechanistically plausible. [19,22,35]

Thymulin Benefits

These benefits, particularly reductions in chronic inflammation and restoration of immune competence, are likely to be more pronounced in older individuals with age-associated immune dysfunction. In animal studies, thymulin substantially improved inflammatory regulation and antitumor immunity in aged mice, with minimal corresponding benefits in young mice. [19] This suggests that thymulin may primarily restore dysregulated immune function rather than enhance already healthy immune activity. Benefits may also occur in younger individuals experiencing pathological inflammation or immune dysfunction. [1,28,29]

- **Likely modulation of T-cell function and thymic signaling.** Zn-thymulin influences thymocyte maturation and peripheral T-cell markers in animal and in vitro studies. [5,6,12]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- **Likely reduction of NF-κB-related inflammatory signaling in appropriate inflammatory contexts.** Zn-thymulin can modulate this pathway and reduce production of selected proinflammatory cytokines. [8,19,29]
- **Plausible antiviral immune support**, based on T/NK-cell biology. [19,22,27]
- **Plausible support for recovery of immune function after an immune stressor.** Zn-thymulin's effects on T-cell maturation and immune signaling provide a mechanistic rationale. [12,26,27]
- **Plausible complementary effects with thymosin alpha-1.** Zn-thymulin and thymosin alpha-1 have partly distinct immunomodulatory actions, providing a mechanistic rationale for complementary effects; reported combined use is anecdotal. [19,22,35]

Indications

- **Adjunctive immune support for recurrent infections**. [12,22,27]
- **Adjunctive support during postviral recovery**. [19,22,27]
- **Age-related immune dysfunction**. [9,19,22–25]
- **Age-associated myeloid inflammatory signaling**. [19]
- **Adjunctive use with thymosin alpha-1 for complementary immunomodulation**. [19,22,35]
- **Adjunct in selected chronic inflammatory or autoimmune contexts.** [4,6,17,28]

Cautions

- **Active or recent malignancy:** use should be reviewed with the treating oncologist. Improved tumor control and checkpoint-treatment response in aged mice do not establish safety or benefit in human cancer or justify modifying cancer treatment. [19]
- **Unstable or uncontrolled autoimmune disease with active severe flares:** avoid unsupervised immune-active interventions.
- **Unsupervised use in people on potent immunosuppressants or biologics:** risk of unpredictable interactions.
- **Pregnancy and breastfeeding:** insufficient safety data.
- **Uncorrected zinc deficiency** may impair active thymulin biology. Adequate zinc status does not establish that an injected zinc-free product is equivalent to preformed Zn-thymulin. [9,20]
- **Anticoagulants, bleeding disorders or procedures**: a short rat experiment found prolonged aPTT and reduced intrinsic-pathway factor activity after thymulin; gross or histological hemorrhage was not observed. Clinical bleeding relevance is uncertain. [33]

Adequate Zinc Status and Thymulin

Zinc deficiency can reduce endogenous active thymulin even when zinc-free peptide remains present. Correcting inadequate intake or deficiency can restore some activity; this is distinct

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

from proving activation of an injected product. Zinc supplementation should address nutritional needs, not be assumed to guarantee quantitative conversion of zinc-free FTS to Zn-thymulin. [9]

Zinc intake should generally meet established dietary requirements, which are 8 mg daily for adult women and 11 mg daily for adult men. The adult tolerable upper intake level of 40 mg daily includes zinc from food, supplements, and medications. [10]

Serum zinc, serum copper, and ceruloplasmin may help assess zinc-copper balance, interpreted in clinical context. Serum zinc is influenced by inflammation and other factors. Excess zinc can impair copper absorption and immune function; increasing zinc beyond nutritional needs is not an established way to improve thymulin effects. [10]

Personal Observation: Morning Congestion

I experienced chronic low-grade postnasal drip and morning congestion for as long as I can remember. Since beginning Thymulin, those symptoms resolved. This is an uncontrolled personal observation.

A speculative explanation is reduced mucosal inflammatory signaling. Thymulin reduced IL-1β release and NF-κB activation in fetal alveolar epithelial cells; a thymulin-analog gene delivered locally by DNA nanoparticles prevented airway inflammation and remodeling in an allergic-asthma mouse model. These are different experimental systems from the personal nasal symptoms and subcutaneous product described here. [8,18]

Why Zn-thymulin Might Help with Serious Infection

Zn-Thymulin has a plausible rationale in severe infection because it is involved in T-cell maturation and function. It supports Th1 immunity, including IL-2 production, which in turn supports cytotoxic T cells and NK cells needed to eliminate infected cells. In **severe infection accompanied by immune exhaustion, restoring these functions could theoretically improve pathogen clearance**. Zn-thymulin also has anti-inflammatory actions, including suppression of NF-κB, p38 MAPK and pro-inflammatory cytokines, so it could potentially strengthen effective cellular immunity without simply increasing systemic inflammation. **The rationale may be stronger in older patients**, because aging is associated with lower thymulin and impaired cellular immunity. [5,8,19,22,27–29]

LPS experiments concern inflammatory modulation in endotoxemia rather than live-pathogen infection. Severe infection requires established medical care. [29]

Why Zn-thymulin Might Help with Minor Infection

For a minor viral infection, Zn-thymulin could theoretically help by strengthening cellular antiviral immunity. It promotes T-cell differentiation and function and is associated with Th1 signaling, including IL-2 and IFN-γ; these pathways support NK cells and cytotoxic T cells that recognize and destroy virus-infected cells. **This rationale is probably weakest in a healthy young person**

whose thymus and cellular immune response are already functioning well, but becomes more plausible in older people, because of thymic involution, declining Zn-thymulin activity, reduced NK-cell activity and weaker cell-mediated immunity accompany aging. [5,19,22,23,27]

Why Zn-thymulin Might Help with Autoimmune Diseases

Thymulin may also make biological sense in autoimmune disease because its function is immunoregulatory rather than simply immunostimulatory. The autoimmune rationale is supported by the RA trails endpoints. [1] Thymulin influences differentiation and activity of multiple T-cell populations, with particularly **notable effects historically reported on suppressor/regulatory T-cell function, which helps prevent immune responses against the body's own tissues**. Thymulin also has anti-inflammatory effects, including inhibition of NF-κB and p38 MAPK signaling and reductions in pro-inflammatory cytokine activity. The plausible result is therefore **better immune regulation: strengthening deficient T-cell functions where necessary while restraining inappropriate inflammatory or autoreactive responses**. Altered or reduced Thymulin activity has been reported in autoimmune diseases, stregthing the mechanistic justification for this application. [4–6,8,17,28]

Thymulin reduced anti-DNA antibodies and glomerular deposits in lupus-prone mice. A separate study used severe experimental autoimmune encephalomyelitis and found reduced disease severity and inflammatory imbalance; it was not a lupus model. These findings support a qualified immunoregulatory rationale. [4,17]

Caution is warranted for patients on potent immunosuppressants or biologics due to risk of unpredictable interactions.

Hashimoto’s thyroiditis

Thymulin’s effects on T-cell maturation and NF-κB-related inflammatory signaling provide a plausible mechanistic rationale for modulating autoimmune inflammation in Hashimoto’s thyroiditis. Anecdotal users report reduced inflammatory symptoms and improved resilience. [5,8]

Rheumatoid arthritis (RA)

In vitro thymulin improved abnormal T-cell subset ratios in eight of nine rheumatoid-arthritis patients, with no significant change in four lupus samples. In two small six-month randomized RA trials, the 5 mg schedule gave the strongest clinical signal. [1]

Rheumatoid arthritis with interstitial lung disease (RA-ILD)

Modulation of the inflammatory component of RA-ILD is mechanistically plausible. In a bleomycin lung-injury mouse model, early thymulin treatment reduced subsequent fibrosis, whereas treatment begun later did not show the same benefit. This supports an early inflammatory-injury hypothesis rather than reversal of established lung scarring. [16]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Thymulin

Healthspan: 1.0 mg daily; Acute infection, esp aged populations: 5.0 daily limited to several weeks

Biological Mechanism

Thymic epithelial cells produce unbound thymulin. Binding zinc forms biologically active Zn-thymulin, an endocrine and paracrine regulator with both intrathymic and extrathymic actions. Exogenous effects on peripheral lymphocytes and myeloid cells do not depend on thymic regrowth. [5,7,19–21]

Mechanistically, Zn-thymulin exerts multiple effects:

Modulation of thymocyte maturation and peripheral T-cell phenotype: Zn-thymulin influences thymocyte maturation and can alter peripheral T-cell markers. Effects on thymic selection, the peripheral T-cell repertoire, and central tolerance are plausible, but the specific receptor and signaling pathways remain incompletely defined. [5,6,12,31]

Thymulin modulates NF-κB-related inflammatory signaling in experimental systems: In fetal alveolar epithelial cells it reduced IL-1β release and NF-κB nuclear translocation; thymulin alone did not reduce IL-6 or TNF-α in that experiment, whereas zinc had additional effects. Other animal models show broader context-dependent cytokine changes. [8,14,19,29]

Historical suppressor-cell findings, T-cell differentiation, and modulation of inflammatory signaling support immune regulation rather than uniform stimulation: Older suppressor-cell assays should not automatically be equated with a defined modern regulatory T-cell phenotype. cAMP-dependent and other downstream pathways have been described in experimental systems, without a complete unified receptor model. [5,8,31]

Endogenous thymulin participates in bidirectional communication between the thymus, immune system, and neuroendocrine system: Experimental effects involve pituitary signaling and vary with age and context. [11,30]

References

[1] Amor B, Dougados M, Mery C, Dardenne M, Bach JF. Nonathymulin in rheumatoid arthritis: two double blind, placebo controlled trials. *Ann Rheum Dis*. 1987;46(7):549–554. doi:10.1136/ard.46.7.549. PMID: 3310925. [Source]

[2] Amor B, Dougados M, Mery C, et al. Thymuline (FTS) in rheumatoid arthritis. *Arthritis Rheum*. 1984;27(1):117–118. doi:10.1002/art.1780270124. PMID: 6362675. [Source]

[3] Faure GC, Bene MC, Thomas P, Tamisier JN. An open trial of thymulin (FTS-Zn) in rheumatoid arthritis patients: sequential clinical and immunological follow-up. *Clin Exp Rheumatol*. 1987;5(3):291–293. PMID: 3427844. [Source]

[4] Israel-Biet D, Noel LH, Bach MA, Dardenne M, Bach JF. Marked reduction of DNA antibody production and glomerulopathy in thymulin (FTS-Zn) or cyclosporin A-treated (NZB × NZW) F1 mice. *Clin Exp Immunol*. 1983;54(2):359–365. PMID: 6360436. PMCID: PMC1535908. [Source]

[5] Bach JF, Dardenne M. Thymulin, a zinc-dependent hormone. *Med Oncol Tumor Pharmacother*. 1989;6(1):25–29. doi:10.1007/BF02985220. PMID: 2657247. [Source]

[6] Faure G, Bene MC, Tamisier JN, Gaucher A, Duheille J. Thymulin (FTS-Zn) induced in vitro modulation of T cell subsets markers on lymphocytes from rheumatoid arthritis and systemic lupus erythematosus patients. *Int J Immunopharmacol*. 1984;6(4):381–388. doi:10.1016/0192-0561(84)90058-4. PMID: 6332789. [Source]

[7] Dardenne M, Savino W, Berrih S, Bach JF. A zinc-dependent epitope on the molecule of thymulin, a thymic hormone. *Proc Natl Acad Sci U S A*. 1985;82(20):7035–7038. doi:10.1073/pnas.82.20.7035. PMID: 2413455. PMCID: PMC391304. [Source]

[8] Haddad JJ. Thymulin and zinc (Zn^{2+})-mediated inhibition of endotoxin-induced production of proinflammatory cytokines and NF-κB nuclear translocation and activation in the alveolar epithelium: unraveling the molecular immunomodulatory, anti-inflammatory effect of thymulin/Zn^{2+} in vitro. *Mol Immunol*. 2009;47(2–3):205–214. doi:10.1016/j.molimm.2009.09.034. PMID: 19850345. [Source]

[9] Prasad AS, Meftah S, Abdallah J, et al. Serum thymulin in human zinc deficiency. *J Clin Invest*. 1988;82(4):1202–1210. doi:10.1172/JCI113717. PMID: 3262625. PMCID: PMC442670. [Source]

[10] National Institutes of Health, Office of Dietary Supplements. Zinc: Fact Sheet for Health Professionals. Updated January 6, 2026. https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/. Accessed November 11, 2026. [Source]

[11] Reggiani PC, Morel GR, Cónsole GM, et al. The thymus-neuroendocrine axis: physiology, molecular biology, and therapeutic potential of the thymic peptide thymulin. *Ann N Y Acad Sci*. 2009;1153:98–106. doi:10.1111/j.1749-6632.2008.03964.x. PMID: 19236333. PMCID: PMC2688715. [Source]

[12] Auger G, Blanot D, Magnin M, et al. Synthesis and biological activity of eight thymulin analogues. *Biol Chem Hoppe Seyler*. 1987;368(5):463–470. doi:10.1515/bchm3.1987.368.1.463. PMID: 3497643. [Source]

[13] Bach JF, Dardenne M, Pleau JM, Rosa J. Biochemical characterisation of a serum thymic factor. *Nature*. 1977;266:55–57. doi:10.1038/266055a0. [Source]

[14] Lunin SM, Khrenov MO, Novoselova TV, Parfenyuk SB, Novoselova EG. Thymulin, a thymic peptide, prevents the overproduction of pro-inflammatory cytokines and heat shock protein Hsp70 in inflammation-bearing mice. *Immunol Invest*. 2008;37(8):858–870. doi:10.1080/08820130802447629. [Source]

[15] Zhao XH, Awaya A, Kobayashi H, Ohnuki T, Tokumitsu Y, Nomura Y. Effects of repeated administrations of facteur thymique sérique (FTS) on biochemical changes related to aging in

senescence-accelerated mouse (SAM). *Jpn J Pharmacol*. 1990;53(3):311–319. doi:10.1254/jjp.53.311. [Source]

[16] Yara S, Kawakami K, Kudeken N, et al. FTS reduces bleomycin-induced cytokine and chemokine production and inhibits pulmonary fibrosis in mice. *Clin Exp Immunol*. 2001;124(1):77–85. doi:10.1046/j.1365-2249.2001.01493.x. [Source]

[17] Lunin SM, Khrenov MO, Novoselova TV, et al. Modulation of inflammatory response in mice with severe autoimmune disease by thymic peptide thymulin and an inhibitor of NF-kappaB signalling. *Int Immunopharmacol*. 2015;25(2):260–266. doi:10.1016/j.intimp.2015.01.021. [Source]

[18] da Silva AL, Martini SV, Abreu SC, et al. DNA nanoparticle-mediated thymulin gene therapy prevents airway remodeling in experimental allergic asthma. *J Control Release*. 2014;180:125–133. doi:10.1016/j.jconrel.2014.02.010. [Source]

[19] Kanemaru H, Luong S, Yamamoto Y, Mizukami Y, Ito F. Thymulin restrains age-associated myeloid inflammation and enhances cancer immunotherapy. *Nat Commun*. 2026;17:6534. doi:10.1038/s41467-026-75383-0. [Source]

[20] Dardenne M, Pléau JM, Nabarra B, et al. Contribution of zinc and other metals to the biological activity of the serum thymic factor. *Proc Natl Acad Sci U S A*. 1982;79(17):5370–5373. doi:10.1073/pnas.79.17.5370. [Source]

[21] Dardenne M, Pleau JM. Interactions between zinc and thymulin. *Met Based Drugs*. 1994;1(2–3):233–239. doi:10.1155/MBD.1994.233. [Source]

[22] Muzzioli M, Mocchegiani E, Bressani N, Bevilacqua P, Fabris N. In vitro restoration by thymulin of NK activity of cells from old mice. *Int J Immunopharmacol*. 1992;14(1):57–61. doi:10.1016/0192-0561(92)90105-T. [Source]

[23] Mocchegiani E, Fabris N. Age-related thymus involution: zinc reverses in vitro the thymulin secretion defect. *Int J Immunopharmacol*. 1995;17(9):745–749. doi:10.1016/0192-0561(95)00064-9. [Source]

[24] Boukaïba N, Flament C, Acher S, et al. A physiological amount of zinc supplementation: effects on nutritional, lipid, and thymic status in an elderly population. *Am J Clin Nutr*. 1993;57(4):566–572. doi:10.1093/ajcn/57.4.566. [Source]

[25] Prasad AS, Fitzgerald JT, Hess JW, Kaplan J, Pelen F, Dardenne M. Zinc deficiency in elderly patients. *Nutrition*. 1993;9(3):218–224. PMID:8353362. [Source]

[26] Okamoto M, Morishita M, Setoguchi C, Nakata K. Restorative effect of short term administration of thymulin on thymus-dependent antibody production in restraint-stressed

mice. *Int J Immunopharmacol*. 1993;15(6):757–762. doi:10.1016/0192-0561(93)90149-S. [Source]

[27] Saha AR, Hadden EM, Hadden JW. Zinc induces thymulin secretion from human thymic epithelial cells in vitro and augments splenocyte and thymocyte responses in vivo. *Int J Immunopharmacol*. 1995;17(9):729–733. doi:10.1016/0192-0561(95)00061-6. [Source]

[28] Henriques-Coelho T, Oliveira SM, Moura RS, et al. Thymulin inhibits monocrotaline-induced pulmonary hypertension modulating interleukin-6 expression and suppressing p38 pathway. *Endocrinology*. 2008;149(9):4367–4373. doi:10.1210/en.2008-0018. [Source]

[29] Novoselova EG, Lunin SM, Glushkova OV, et al. Thymulin, free or bound to PBCA nanoparticles, protects mice against chronic septic inflammation. *PLoS One*. 2018;13(5):e0197601. doi:10.1371/journal.pone.0197601. [Source]

[30] Buckingham JC, Safieh B, Singh S, Arduino LA, Cover PO, Kendall MD. Interactions between the hypothalamo-pituitary adrenal axis and the thymus in the rat: a role for corticotrophin in the control of thymulin release. *J Neuroendocrinol*. 1992;4(3):295–301. doi:10.1111/j.1365-2826.1992.tb00171.x. PMID:21554609. [Source]

[31] Pléau JM, Fuentes V, Morgat JL, Bach JF. Specific receptors for the serum thymic factor (FTS) in lymphoblastoid cultured cell lines. *Proc Natl Acad Sci U S A*. 1980;77(5):2861–2865. doi:10.1073/pnas.77.5.2861. [Source]

[32] Roullet E, Cesaro P, Simon-Lavoine N, Degos JD, Marteau R. Nonathymulin treatment of multiple sclerosis: double-blind pilot study. *Acta Neurol Scand*. 1989;80(6):575–578. doi:10.1111/j.1600-0404.1989.tb03929.x. [Source]

[33] Negrev N, Hristosov K, Hachmeryan A, Nyagolov Y. Thymus peptides (thymulin, thymosin alpha 1 and thymosin beta 4) inhibiting effects on the intrinsic blood coagulation pathway in rats. *J of IMAB*. 2017;23(3):1641–1645. doi:10.5272/jimab.2017233.1641. [Source]

[34] US Food and Drug Administration. Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers. Guidance for Industry. July 2005. https://www.fda.gov/media/72309/download. Accessed November 11, 2026. [Source]

[35] Garaci E, Pica F, Matteucci C, Costantini C, Puccetti P, Romani L. Phenotypic drug discovery: a case for thymosin alpha-1. *Front Med (Lausanne)*. 2024;11:1388959. doi:10.3389/fmed.2024.1388959. [Source]