

When acute soft-tissue injury occurs, like a torn tendon, strained ligament, or severe connective-tissue trauma, the body initiates a complex repair cascade involving inflammation, fibroblast migration, angiogenesis, collagen deposition, and matrix remodeling. TB-500 and BPC-157 are frequently discussed because they act at different, complementary points in this cascade.

💉💉 Probably safe (some human evidence)

★ ★ Probably effective (some positive human signals)

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 30-Aug-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/acute-injury-repair.pdf>



Long-term safety for non-continuous TB-500 use is likely. Long-term safety for continuous use is less certain because of plausible pro-tumor concerns associated with angiogenesis. Long-term safety for non-continuous BPC-157 use is also likely. Long-term safety for continuous use is likewise less certain because of plausible pro-tumor concerns associated with angiogenesis.

TB-500 is likely to improve wound healing. BPC-157 is likely to improve soft-tissue healing, including tendon and muscle repair.

This document presents a plausible protocol and doses for TB-500 and BPC-157 for supporting healing acute soft tissue injuries. There is no evidence-based protocol or dose for these. Instead we synthesis and extrapolate from published clinical notes and forum discussions.

TB-500 as a Systemic Repair Mobilizer

TB-500 plausibly supports systemic repair signaling. Its proposed effects include:

- **Actin cytoskeleton remodeling**
- **Cell migration** (endothelial, progenitor, and stem cells)
- **Angiogenesis** (new blood vessel formation)
- **Improved microcirculation**
- **Reduced fibrosis**
- **Enhanced extracellular matrix reorganization**

BPC-157 as a Repair Accelerator

BPC-157 likely supports several repair processes, including:

- **Fibroblast migration** to the injury site
- **Collagen synthesis** and improved tendon/ligament fiber alignment
- **Angiogenesis** (microvascular repair)
- **Growth-factor modulation** (VEGF, FGF)
- **Reduced inflammatory cytokines** (TNF- α , IL-1 β , IL-6)
- **Improved epithelial and endothelial repair**

Twice-daily dosing of BPC-157 during the acute phase plausibly provides repeated support for repair signaling.

Which Injury Type Suggests Conservative vs Aggressive Treatment?

Conservative Protocol

Lower doses plausibly reduce exposure and theoretical risk.

- Chronic tendonitis
- Mild strains
- Overuse injuries
- Small partial tears
- Cautious users

Moderate Protocol

Organized around the approximate natural repair timeline.

- Typical acute tendon or ligament injuries
- Partial tears confirmed by imaging
- Post-surgical tendon/ligament repair
- Moderate rotator cuff injuries
- Patellar tendon injuries
- Hamstring tendon injuries

Aggressive Protocol

- Severe acute injuries
- Large tendon injuries (Achilles, quadriceps tendon)
- Major ligament injuries (MCL/LCL partial tears)
- Situations requiring accelerated recovery

Injury Repair Phases

Injury repair occurs in overlapping phases, and this protocol organizes dosing around their approximate progression. The body's repair goals in early healing (weeks 1–4) is rapid fibroblast recruitment, angiogenesis, inflammation control, and early collagen organization. The goals in intermediate healing (weeks 5–8) is strengthen collagen crosslinking, stabilize microcirculation, and support tendon/ligament remodeling. The goals of maintenance (weeks 9–10) are to support latestage remodeling and prevent reinjury. (Timelines are approximate, minor injuries heal faster, severe injuries take longer.)

Doses

		Conservative	Moderate	Aggressive
Early healing ~weeks 1-4	TB-500	2 mg 2x weekly	2.5 mg 2x weekly	5 mg 2x weekly
	BPC-157	250 mcg daily	250 mcg 2x daily preferred, or 500 mcg daily	500 mcg 2x daily preferred, or 750 mcg daily
Intermediate healing ~weeks 5-8	TB-500	1 mg 2x weekly, or 2 mg weekly	1.25 mg 2x weekly, or 2.5 mg weekly	2.5 mg 2x weekly, or 5 mg weekly
	BPC-157	250 mcg daily	250 mcg 2x daily, or 500 mcg daily	500 mcg 2x daily, or 750 mcg daily
Maintenance ~weeks 9+	TB-500	n/a	n/a	1 mg weekly
	BPC-157		250 mcg daily	250 mcg daily

TB-500 + BPC-157 Benefits (Abbreviated)

- Likely faster tendon and ligament healing
- Likely improved collagen organization
- Likely improved tissue strength during repair-
- Plausibly reduced reinjury risk

Cautions

Because these peptides may promote angiogenesis, caution may be warranted in the following situations:

- Active cancer
- Unexplained masses
- Pregnancy
- Systemic infection
- Severe uncontrolled inflammation

Other Peptides

Other peptides are discussed in the context of repair.

KPV likely has **anti-inflammatory effects** and is plausibly **relevant during acute injury repair**, although its additional benefit in this protocol is uncertain. Doses of 250 to 750 mcg are often discussed.

ARA-290 likely **supports nerve repair**. If the acute injury involves nerve damage, ARA-290 is plausibly beneficial. Doses of 2 to 4 mg daily are often discussed for nerve repair.

Although **GHK-Cu** is often discussed in the context of skin and hair, it likely **supports tissue repair and wound remodeling**. It is plausibly appropriate to include GHK-Cu when scar remodeling is an issue. Doses of 2 mg daily are often discussed for tissue repair.

For more detailed information

TB-500 - <https://pdfs.researcher6076.com/pdfs/TB-500.pdf>

BPC-157 - <https://pdfs.researcher6076.com/pdfs/BPC-157.pdf>

KPV - <https://pdfs.researcher6076.com/pdfs/KPV.pdf>

ARA-290 - <https://pdfs.researcher6076.com/pdfs/ARA-290.pdf>

GHK-Cu - <https://pdfs.researcher6076.com/pdfs/GHK-Cu.pdf>

References

BPC-157

1. Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157: novel therapy in gastrointestinal tract. *Curr Pharm Des.* 2011;17(16):1612–1632. doi:10.2174/138161211796196954.
2. Staresinic M, Sebecic B, Patrlj L, et al. Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth. *J Orthop Res.* 2003;21(6):976–983. doi:10.1016/S0736-0266(03)00110-4.
3. Chang CH, Tsai WC, Lin MS, Hsu YH, Su Pang JH. The promoting effect of pentadecapeptide BPC 157 on tendon healing involves tendon outgrowth, cell survival, and cell migration. *J Appl Physiol* (1985). 2011;110(3):774–780. doi:10.1152/jappphysiol.00945.2010.
4. Gwyer D, Wragg NM, Wilson SL. Gastric pentadecapeptide body protection compound BPC 157 and its role in accelerating musculoskeletal soft tissue healing. *Cell Tissue Res.* 2019;377:153–159. doi:10.1007/s00441-019-03016-8.
5. Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157-NO-system relation. *Curr Pharm Des.* 2014;20(7):1126–1135. doi:10.2174/13816128113190990411.
6. Sever AZ, Sever M, Vidovic T, et al. Stable gastric pentadecapeptide BPC 157 in the therapy of the rats with bile duct ligation. *Eur J Pharmacol.* 2019;847:130–142. doi:10.1016/j.ejphar.2019.01.030.
7. Lee E, Padgett B. Intra-articular injection of BPC 157 for multiple types of knee pain. *Altern Ther Health Med.* 2021;27(4):8–13. PMID:34324435.

8. Vasireddi N, Hahamyan H, Salata MJ, et al. Emerging use of BPC-157 in orthopaedic sports medicine: a systematic review. HSS J. 2025;21(4):485–495. doi:10.1177/15563316251355551.
9. U.S. Food and Drug Administration. FDA Briefing Document for BPC-157-Related Bulk Drug Substances (BPC-157 [free base] and BPC-157 acetate). Pharmacy Compounding Advisory Committee Meeting; July 23–24, 2026. <https://www.fda.gov/media/193343/download>.

KPV

10. Getting SJ, Schiöth HB, Perretti M. Dissection of the anti-inflammatory effect of the core and C-terminal (KPV) alpha-melanocyte-stimulating hormone peptides. J Pharmacol Exp Ther. 2003;306(2):631–637. doi:10.1124/jpet.103.051623.
11. Kannengiesser K, Maaser C, Heidemann J, et al. Melanocortin-derived tripeptide KPV has anti-inflammatory potential in murine models of inflammatory bowel disease. Inflamm Bowel Dis. 2008;14(3):324–331. doi:10.1002/ibd.20334.
12. U.S. Food and Drug Administration. FDA Briefing Document for KPV-Related Bulk Drug Substances (KPV [free base] and KPV acetate). Pharmacy Compounding Advisory Committee Meeting; July 23–24, 2026. <https://www.fda.gov/media/193346/download>.

ARA-290

13. Liu G, Li W, Jiang S, et al. ARA290, an alternative of erythropoietin, inhibits activation of NLRP3 inflammasome in Schwann cells after sciatic nerve injury. Eur J Pharmacol. 2025;997:177610. doi:10.1016/j.ejphar.2025.177610.
14. Brines M, Dunne AN, van Velzen M, et al. ARA 290, a nonerythropoietic peptide engineered from erythropoietin, improves metabolic control and neuropathic symptoms in patients with type 2 diabetes. Mol Med. 2015;20(1):658–666. doi:10.2119/molmed.2014.00215.

GHK-Cu

15. Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. FEBS Lett. 1988;238(2):343–346. doi:10.1016/0014-5793(88)80509-X.
16. Maquart FX, Bellon G, Chaqour B, et al. In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ in rat experimental wounds. J Clin Invest. 1993;92(5):2368–2376. doi:10.1172/JCI116842.

17. Siméon A, Emonard H, Hornebeck W, Maquart FX. The tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ stimulates matrix metalloproteinase-2 expression by fibroblast cultures. *Life Sci.* 2000;67(18):2257–2265. doi:10.1016/S0024-3205(00)00803-1.
18. U.S. Food and Drug Administration. Certain Bulk Drug Substances for Use in Compounding That May Present Significant Safety Risks: GHK-Cu for Injectable Routes of Administration. <https://www.fda.gov/drugs/human-drug-compounding/certain-bulk-drug-substances-use-compounding-may-present-significant-safety-risks>.

Thymosin β 4 / TB-500 mechanistic literature

TB-500-specific evidence

7. Sosne G, Qiu P, Goldstein AL, Wheeler M. Biological activities of thymosin beta4 defined by active sites in short peptide sequences. *FASEB J.* 2010;24(7):2144–2151. doi:10.1096/fj.09-142307.
8. U.S. Food and Drug Administration. FDA Briefing Document for TB-500-Related Bulk Drug Substances (TB-500 [free base] and TB-500 acetate). Pharmacy Compounding Advisory Committee Meeting; July 23–24, 2026. <https://www.fda.gov/media/193349/download>.
9. Bock-Marquette I, Saxena A, White MD, DiMaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. *Nature.* 2004;432(7016):466–472. doi:10.1038/nature03000.
10. Smart N, Risebro CA, Melville AAD, Moses K, Schwartz RJ, Chien KR, Riley PR. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. *Nature.* 2007;445(7124):177–182. doi:10.1038/nature05383.
11. Evans MA, Smart N, Dubé KN, et al. Thymosin β 4-sulfoxide attenuates inflammatory cell infiltration and promotes cardiac wound healing. *Nat Commun.* 2013;4:2081. doi:10.1038/ncomms3081.
12. Maar K, Thatcher JE, Karpov E, et al. Thymosin Beta-4 modulates cardiac remodeling by regulating ROCK1 expression in adult mammals. *Int J Mol Sci.* 2025;26(9):4131. doi:10.3390/ijms26094131.
13. Srivastava D, Ieda M, Fu J, Qian L. Cardiac repair with thymosin β 4 and cardiac reprogramming factors. *Ann N Y Acad Sci.* 2012;1270:66–72. doi:10.1111/j.1749-6632.2012.06696.x.
14. Maar K, Hetenyi R, Maar S, Faskerti G, Hanna D, Lippai B, Takatsy A, Bock-Marquette I. Utilizing developmentally essential secreted peptides such as thymosin Beta-4 to remind the adult organs of their embryonic state—new directions in anti-aging regenerative therapies. *Cells.* 2021;10(6):1343. doi:10.3390/cells10061343.

15. Xing Y, Ye Y, Zuo H, Li Y. Progress on the function and application of thymosin β 4. *Front Endocrinol (Lausanne)*. 2021;12:767785. doi:10.3389/fendo.2021.767785.

Thymosin β 4 — wound and tissue repair

14. Malinda KM, Sidhu GS, Mani H, Banaudha K, Maheshwari RK, Goldstein AL, Kleinman HK. Thymosin beta4 accelerates wound healing. *J Invest Dermatol*. 1999;113(3):364–368. doi:10.1046/j.1523-1747.1999.00708.x.
15. Philp D, Nguyen M, Scheremeta B, St-Surin S, Villa AM, Orgel A, Kleinman HK, Elkin M. Thymosin beta4 increases hair growth by activation of hair follicle stem cells. *FASEB J*. 2004;18(2):385–387. doi:10.1096/fj.03-0244fje.
16. Sosne G, Szliter EA, Barrett R, Kernacki KA, Kleinman HK, Hazlett LD. Thymosin beta 4 promotes corneal wound healing and decreases inflammation in vivo following alkali injury. *Exp Eye Res*. 2002;74(2):293–299. doi:10.1006/exer.2001.1125.
17. Sosne G, Ousler GW. Thymosin beta 4 ophthalmic solution for dry eye: a randomized, placebo-controlled, Phase II clinical trial conducted using the controlled adverse environment (CAE™) model. *Clin Ophthalmol*. 2015;9:877–884. doi:10.2147/OPTH.S80954.
18. Conte E, Genovese T, Gili E, et al. Thymosin β 4 protects C57BL/6 mice from bleomycin-induced damage in the lung. *Eur J Clin Invest*. 2013;43(3):309–315. doi:10.1111/eci.12048.

Thymosin β 4 — neurological repair

19. Xiong Y, Zhang Y, Mahmood A, et al. Neuroprotective and neurorestorative effects of thymosin beta4 treatment initiated 6 hours post injury following traumatic brain injury in rats. *J Neurosurg*. 2012;116(5):1081–1092. doi:10.3171/2012.1.JNS111729.
20. Morris DC, Chopp M, Zhang L, Lu M, Zhang ZG. Thymosin β 4 improves functional neurological outcome in a rat model of embolic stroke. *Neuroscience*. 2010;169(2):674–682. doi:10.1016/j.neuroscience.2010.05.017.