

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

BPC-157 is likely to accelerate soft-tissue healing in tendons, ligaments, and muscles, support repair of gastrointestinal injury, and improve outcomes after traumatic brain injury. It is one of the most widely used peptides, with few serious safety concerns reported.

💉 Possibly safe. Limited human exposure and very widespread community use with few serious adverse events reported are reassuring. Long-term safety remains plausible.

★ ★ Probably effective based on strong animal evidence.

C Pro-angiogenic: Use particular caution with active or recently cancer. BPC-157 promotes angiogenic signaling in experimental models, making increased tumor vascularization or growth a plausible risk.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.



This document was updated 26-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/bpc-157.pdf>

BPC-157 is a synthetic 15-amino-acid peptide whose sequence was described as a fragment of a protein originally isolated from human gastric juice. It has been widely studied in preclinical models for its ability to accelerate healing of tendons, muscles, nerves, and the gastrointestinal tract through coordinated vascular and anti-inflammatory mechanisms.

Long-term safety is plausible based on preclinical findings, limited human exposure, and very widespread community use with relatively few serious adverse effects reported.

Improved soft-tissue healing is likely based on reproducible tendon and muscle repair findings in animal models, supported by effects on angiogenesis, fibroblast migration, collagen organization, and nitric-oxide signaling.

Dosing Patterns

Human dosing remains speculative. The following doses and schedules reflect commonly reported practitioner and community practice, informed in part by preclinical research.

For specific injury repair

- **100–500 mcg SC**, with **250 mcg** being most common, **once or twice daily**

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

- **Subcutaneous administration is commonly used, with any additional benefit from injecting near the injured area is speculative.**
- **2–8 weeks** based on injury

For chronic injuries and age-related stiffness

- **100–500 mcg SC**, with **250 mcg** being most common, **daily to once weekly**
- **Continuously**

Potential for Diminishing Benefit

BPC-157 has no established primary receptor and appears to influence several downstream pathways. Classical receptor desensitization has not been demonstrated, while other biological adaptation during continuous exposure remains plausible.

Animal pharmacokinetic studies found an elimination half-life of less than 30 minutes after intravenous or intramuscular administration. The duration of downstream biological effects and the value of cycling remain unknown.

Community cycling patterns

As an injury progresses from active repair into later remodeling, the additional benefit of continuing BPC-157 may plausibly diminish. This provides a **mechanistic rationale for higher dose, time-limited courses**, but does not establish that cycling is required.

Continuous daily dosing, weekly maintenance dosing, and periodic breaks are reported community practices. Their relative effectiveness and optimal timing remain speculative.

Benefits

Taken together, the preliminary human findings and broader experimental evidence suggest that BPC-157 plausibly has local tissue-protective, repair-supporting, and symptom-modulating effects across injured connective and epithelial tissues.

- **Tendon and ligament repair:** BPC-157 is likely to support collagen organization, fibroblast migration, tissue outgrowth, and biomechanical recovery based on rodent models of Achilles-tendon and medial-collateral-ligament injury.
- **Muscle repair:** BPC-157 is likely to improve myofiber regeneration, reduce tissue necrosis, and support recovery of contractile function after crush or transection injury based on rodent models.
- **Intra-articular knee-pain relief:** A small retrospective human series reported significant knee-pain improvement in 11 of 12 patients who received BPC-157 alone. Structural joint or cartilage repair remains speculative.

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

- **Gastrointestinal protection and repair:** BPC-157 is likely to protect injured gastrointestinal mucosa and support healing of ulcers, intestinal anastomoses, and fistulas based on rodent models.
- **Interstitial-cystitis symptoms:** An uncontrolled 12-patient human pilot reported substantial symptom improvement after a single procedure administering BPC-157 around inflamed areas of the bladder. Ten patients reported complete symptom resolution, and two reported approximately 80% improvement. This preliminary finding applies specifically to local bladder-wall administration.
- **Microvascular repair:** BPC-157 is likely to support angiogenesis, endothelial nitric-oxide signaling, and restoration of tissue perfusion in injured or ischemic tissue based on animal evidence.
- **Acute neurologic injury:** BPC-157 is likely to reduce tissue damage and support functional recovery after traumatic brain injury, spinal-cord injury, and ischemic injury based on rodent models.
- **Inflammatory and oxidative signaling:** BPC-157 is likely to reduce inflammatory-mediator activity and oxidative damage within injured tissues based on animal models.
- **Wound and skin repair:** BPC-157 is likely to support re-epithelialization, granulation-tissue formation, angiogenesis, fibroblast activity, and collagen deposition based on rodent wound models.
- **Acute incisional pain:** BPC-157 is likely to provide short-term antinociceptive effects after acute incisional injury based on controlled rat evidence.
- **General recovery and nonspecific pain:** Faster overall recovery and broader nonspecific pain relief are widely reported in peptide communities but remain speculative.

Potential Applications

- **Tendon or ligament injury:** Likely to support repair, with potential applications to strains, sprains, partial tears, and chronic tendinopathy.
- **Muscle injury:** Likely to support repair after muscle tears and other direct muscle injury.
- **Post-surgical tissue repair:** Likely to support healing in comparable soft-tissue and gastrointestinal repair settings.
- **Gastrointestinal injury:** Likely to support mucosal protection and repair in NSAID-associated injury, ulcerative injury, and intestinal inflammation.
- **Peripheral nerve injury:** Likely to support axonal regeneration and functional recovery after direct nerve injury.
- **Neuropathic symptoms:** Improvement in neuropathic pain or sensory symptoms is plausible.

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

Cautions

- **Active or recent cancer:** Use particular caution because BPC-157 promotes angiogenic signaling in experimental models, making increased tumor vascularization or growth a plausible concern.
- **Pregnancy or breastfeeding:** Use caution because safety during pregnancy and breastfeeding is unknown.
- **Unstable blood pressure or vascular disease:** Use caution because BPC-157 may plausibly alter nitric-oxide-mediated vascular tone.
- **Agents with overlapping repair effects:** Additive angiogenic, vasomotor, or tissue-remodeling effects are speculative when BPC-157 is combined with other regenerative peptides or related experimental agents.

Side Effects

- **Community-reported symptoms:** Headache, dizziness, lightheadedness, changes in appetite or digestion, fatigue, and lethargy are occasionally reported but remain speculative adverse effects.
- **Blood-pressure effects:** Changes in blood pressure are plausible because BPC-157 influences nitric-oxide-mediated vascular tone in experimental models.

Biological Mechanism

Experimental studies indicate that BPC-157 influences multiple repair pathways. It increases VEGFR2 expression and internalization and activates VEGFR2–Akt–eNOS signaling in endothelial and animal models. These effects provide a plausible mechanism for angiogenesis, nitric-oxide-mediated vascular tone, and microvascular repair.

In tendon models, BPC-157 promotes tendon-fibroblast migration, spreading, survival under oxidative stress, and explant outgrowth, with FAK–paxillin activation providing a plausible cellular mechanism. Animal injury studies also show improved collagen organization and biomechanical repair, supporting likely tendon and ligament healing.

In gastrointestinal animal models, BPC-157 modulates nitric-oxide and prostaglandin signaling, preserves microvascular and endothelial integrity, reduces inflammatory injury, and improves mucosal, anastomotic, and fistula healing. These findings support likely gastrointestinal protection and repair.

In animal central-nervous-system injury models, BPC-157 reduces tissue damage and improves functional recovery, supporting likely neuroprotective effects. Reduced oxidative stress and inflammatory signaling, together with preserved neurovascular function, provide plausible contributing mechanisms.

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

Overall, BPC-157 plausibly functions as a pleiotropic repair peptide by influencing vascular, inflammatory, and cellular repair pathways across multiple tissues.

BPC-157 and Cancer Risk

BPC-157 promotes angiogenesis and VEGFR2–Akt–eNOS signaling in experimental models. These effects make increased tumor vascularization or growth a plausible concern in active or recently treated cancer. Preliminary in-vitro evidence suggests possible anti-tumor effects, including reduced proliferation and VEGF signaling in a human melanoma cell line. The net effect of BPC-157 on cancer remains unknown.

BPC-157 and Autoimmune Conditions

BPC-157 is likely to reduce experimental inflammatory arthritis based on animal evidence. Its effects on inflammatory signaling, oxidative stress, endothelial function, and gastrointestinal-barrier repair provide plausible mechanisms for reducing inflammatory tissue injury in some autoimmune settings. Efficacy for rheumatoid arthritis, psoriasis, autoimmune gastritis, or other human autoimmune conditions remains speculative.

Conclusions

Animal evidence indicates that BPC-157 is likely to support repair of tendon, ligament, muscle, gastrointestinal, dermal, vascular, peripheral-nerve, and central-nervous-system injuries. Short-term safety is possible, and long-term safety is plausible based on preclinical findings, limited human exposure, and very widespread community use. Human dosing, continuous use, and cycling practices remain speculative.

References

- U.S. Food and Drug Administration. Pharmacy Compounding Advisory Committee briefing document: BPC-157-related bulk drug substances (BPC-157 free base and BPC-157 acetate). July 23, 2026.
- Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157: novel therapy in gastrointestinal tract. *Curr Pharm Des.* 2011;17(16):1612–1632. doi:10.2174/138161211796196954.
- Staresinic M, Petrovic I, Novinscak T, et al. Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth. *J Orthop Res.* 2003;21(5):976–983.
- Ceroveckí T, Bojanic I, Brcic L, et al. Pentadecapeptide BPC 157 (PL 14736) improves ligament healing in the rat. *J Orthop Res.* 2010;28(9):1155–1161. doi:10.1002/jor.21107.
- Pevec D, Novinscak T, Brcic L, et al. Impact of pentadecapeptide BPC 157 on muscle healing impaired by systemic corticosteroid application. *Med Sci Monit.* 2010;16(3):BR81–BR88.

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

- Chang CH, Tsai WC, Lin MS, Hsu YH, Pang JHS. The promoting effect of pentadecapeptide BPC 157 on tendon healing involves tendon outgrowth, cell survival, and cell migration. *J Appl Physiol* (1985). 2011;110(3):774–780. doi:10.1152/japplphysiol.00945.2010.
- Gwyer D, Wragg NM, Wilson SL. Gastric pentadecapeptide body protection compound BPC 157 and its role in accelerating musculoskeletal soft tissue healing. *Cell Tissue Res*. 2019;377(2):153–159. doi:10.1007/s00441-019-03016-8.
- Hsieh MJ, Liu HT, Wang CN, et al. Therapeutic potential of pro-angiogenic BPC157 is associated with VEGFR2 activation and up-regulation. *J Mol Med (Berl)*. 2017;95(3):323–333. doi:10.1007/s00109-016-1488-y.
- Hsieh MJ, Lee CH, Chueh HY, et al. Modulatory effects of BPC 157 on vasomotor tone and the activation of Src–caveolin-1–endothelial nitric oxide synthase pathway. *Sci Rep*. 2020;10(1):17078. doi:10.1038/s41598-020-74022-y.
- He L, Feng D, Guo H, et al. Pharmacokinetics, distribution, metabolism, and excretion of body-protective compound 157, a potential drug for treating various wounds, in rats and dogs. *Front Pharmacol*. 2022;13:1026182. doi:10.3389/fphar.2022.1026182.
- Xu C, Sun L, Ren F, et al. Preclinical safety evaluation of body protective compound-157, a potential drug for treating various wounds. *Regul Toxicol Pharmacol*. 2020;114:104665. doi:10.1016/j.yrtph.2020.104665.
- Klicek R, Sever M, Radic B, et al. Pentadecapeptide BPC 157, in clinical trials as a therapy for inflammatory bowel disease (PL14736), is effective in the healing of colocutaneous fistulas in rats: role of the nitric oxide-system. *J Pharmacol Sci*. 2008;108(1):7–17. doi:10.1254/jphs.FP0072161.
- Sikiric P, Seiwerth S, Grabarevic Z, et al. Pentadecapeptide BPC 157 positively affects both non-steroidal anti-inflammatory agent-induced gastrointestinal lesions and adjuvant arthritis in rats. *J Physiol Paris*. 1997;91(3–5):113–122. doi:10.1016/S0928-4257(97)89474-0.
- Huang T, Zhang K, Sun L, et al. Body protective compound-157 enhances alkali-burn wound healing in vivo and promotes proliferation, migration, and angiogenesis in vitro. *Drug Des Devel Ther*. 2015;9:2485–2499. doi:10.2147/DDDT.S82030.
- Jung YH, Kim H, Kim H, Kim E, Baik J, Kang H. The anti-nociceptive effect of BPC-157 on the incisional pain model in rats. *J Dent Anesth Pain Med*. 2022;22(2):97–105. doi:10.17245/jdapm.2022.22.2.97.
- Gjurasin M, Miklic P, Zupancic B, et al. Peptide therapy with pentadecapeptide BPC 157 in traumatic nerve injury. *Regul Pept*. 2010;160(1–3):33–41. doi:10.1016/j.regpep.2009.11.005.
- Tudor M, Jandric I, Marovic A, et al. Traumatic brain injury in mice and pentadecapeptide BPC 157 effect. *Regul Pept*. 2010;160(1–3):26–32. doi:10.1016/j.regpep.2009.11.012.
- Perovic D, Kolenc D, Bilic V, et al. Stable gastric pentadecapeptide BPC 157 can improve the healing course of spinal cord injury and lead to functional recovery in rats. *J Orthop Surg Res*. 2019;14(1):199. doi:10.1186/s13018-019-1242-6.

BPC-157 Available for Research Use Only

(0.1 mg – 0.5 mg daily, increase to 1 mg daily as needed)

- Vukojevic J, Sikiric P, Seiwerth S, et al. The effect of pentadecapeptide BPC 157 on hippocampal ischemia/reperfusion injuries in rats. *Brain Behav.* 2020;10(8):e01726. doi:10.1002/brb3.1726.
- Sever AZ, Sever M, Vidovic T, et al. Stable gastric pentadecapeptide BPC 157 in the therapy of the rats with bile duct ligation. *Eur J Pharmacol.* 2019;847:130–142. doi:10.1016/j.ejphar.2019.01.030.
- Radeljak S, Seiwerth S, Sikiric P. BPC 157 inhibits cell growth and VEGF signaling via the MAPK kinase pathway in the human melanoma cell line. *Melanoma Res.* 2004;14:A14–A15. Conference abstract.
- Lee E, Padgett B. Intra-articular injection of BPC 157 for multiple types of knee pain. *Altern Ther Health Med.* 2021;27(4):8–13.
- Lee E, Walker C, Ayadi B. Effect of BPC-157 on symptoms in patients with interstitial cystitis: a pilot study. *Altern Ther Health Med.* 2024;30(10):12–17.
- Lee E, Burgess K. Safety of intravenous infusion of BPC157 in humans: a pilot study. *Altern Ther Health Med.* 2025;31(5):20–24.