

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

(Metformin 500 mg, Tadalafil 5 mg, daily)

Metformin and tadalafil are prescription drugs with distinct but complementary mechanisms relevant to experimental healthspan use. Metformin improves cellular energy sensing, glucose regulation, and mitochondrial function. Tadalafil sustains nitric-oxide and cyclic-guanosine-monophosphate signaling and is likely to improve endothelial function and cerebral perfusion. Their effects on complementary metabolic and vascular pathways make a combined healthspan benefit plausible.

💉💉 Safety is likely. Both drugs have established safety profiles for long-term use within approved prescribing parameters, and current prescribing information identifies no specific interaction between them.

★ Plausibly effective for improving healthspan.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 02-Oct-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/metform-plus-tadalafil-for-longevity.pdf>



Why Signaling Changes Matter for Healthspan

Conserved aging pathways: Many interventions that extend healthspan in model organisms converge on a small set of conserved pathways: **reduced insulin/IGF signaling, activation of AMPK/sirtuins, suppression of mTORC1, enhanced autophagy, improved mitochondrial function, reduced chronic inflammation, and preserved proteostasis.**

Biomarker → mechanism → outcome logic: If a therapy reliably **activates or restores** these pathways in tissues relevant to aging (metabolic organs, vasculature, brain), it provides **biologically plausible** grounds to hypothesize improved healthspan, especially when multiple complementary pathways are affected.

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

(Metformin 500 mg, Tadalafil 5 mg, daily)

Metformin: Links to Healthspan

Primary cellular effects

- **Mitochondrial and redox signaling:** Metformin can modulate mitochondrial complex I and the cellular redox state, sometimes increasing the AMP-to-ATP ratio and activating energy-stress signaling. This could plausibly promote adaptive stress responses and cellular maintenance relevant to age-related resilience.
- **AMPK and mTOR signaling:** Metformin can activate AMP-activated protein kinase and reduce mTORC1 signaling in relevant metabolic contexts, supporting glucose uptake, fatty-acid oxidation, and cellular maintenance. These effects could plausibly improve metabolic resilience and slow aspects of age-related functional decline.
- **Gut-mediated effects:** Metformin increases intestinal glucose utilization and GLP-1 signaling and can alter bile-acid metabolism and the gut microbiome, including enrichment of *Akkermansia* and short-chain-fatty-acid-producing species in some studies. These changes plausibly contribute to improved insulin sensitivity, lower postprandial glucose excursions, and broader metabolic and anti-inflammatory effects.

Downstream maintenance programs

- **Autophagy and proteostasis:** By influencing AMPK and mTORC1 signaling, metformin could plausibly support autophagy and the clearance of damaged proteins and organelles. Greater autophagic and proteostatic activity supports improved healthspan in multiple experimental organisms.
- **Glucose and insulin regulation:** Metformin reduces hepatic glucose production and improves glucose regulation. In people with insulin resistance, prediabetes, or diabetes, this can reduce chronic hyperinsulinemia. Lower insulin exposure could plausibly support healthspan, while broader suppression of insulin-like growth factor signaling should not be assumed.
- **Inflammatory regulation:** Metformin can reduce selected inflammatory biomarkers in some metabolically affected populations. A broader reduction in chronic low-grade inflammaging and its contribution to age-related disease is plausible.
- **Senescence-associated signaling:** Metformin is likely to modulate senescence-associated signaling and the senescence-associated secretory phenotype in tissues and models where these effects have been observed. A broader reduction in human tissue dysfunction through these pathways is plausible.

Why these map to healthspan: AMPK activation, mTOR suppression, and enhanced autophagy are canonical cellular-maintenance signals in model organisms. Metformin's

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
2-Oct-26*

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack (Metformin 500 mg, Tadalafil 5 mg, daily)

ability to engage these programs provides a mechanistic bridge from drug action to cellular processes that influence healthspan.

Tadalafil: Links to Healthspan

Primary vascular / signaling effects

- **Endothelial signaling:** Tadalafil inhibits phosphodiesterase type 5, sustaining cyclic guanosine monophosphate and nitric-oxide signaling and producing vasodilation. Daily tadalafil can improve endothelial function in some human populations. These effects could plausibly protect vascular-dependent tissues from aspects of age-related decline.
- **Microvascular perfusion:** Tadalafil is likely to improve perfusion and oxygen delivery in tissues responsive to nitric-oxide and cyclic-guanosine-monophosphate signaling. Animal and limited human evidence make support for cardiac, renal, skeletal-muscle, and cerebral perfusion likely, while preservation of multi-organ function and slower age-related decline remain plausible.

Downstream maintenance and protective programs

- **Mitochondrial protection:** Tadalafil is likely to improve mitochondrial efficiency, reduce reactive-oxygen-species production, and influence autophagy and mitophagy in tissues studied in animal models. Comparable systemic cytoprotective effects in humans are plausible.
- **Vascular inflammation and platelet signaling:** Human studies support favorable effects on endothelial and inflammatory biomarkers, while limited evidence suggests reduced platelet activation in some populations. Observational outcome studies make a reduction in atherothrombotic events possible.
- **Neurovascular support:** Tadalafil is likely to improve cerebral perfusion in responsive populations. Through improved blood flow and effects on cyclic-guanosine-monophosphate, CREB, and BDNF signaling, support for metabolite clearance, synaptic function, and protection from neurodegenerative processes is plausible.
- **Lymphatic and interstitial flow:** PDE5 inhibition could plausibly modulate lymphatic smooth-muscle tone and facilitate lymphatic drainage in some contexts. Resulting support for interstitial-fluid clearance and immune-cell trafficking is plausible.

Why these map to healthspan: Vascular health is a major determinant of multi-organ aging; preserving endothelial function and microcirculation reduces the burden of ischemia, inflammation, and organ dysfunction that drive morbidity and mortality.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
2-Oct-26*

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack (Metformin 500 mg, Tadalafil 5 mg, daily)

Sustained NO–cGMP signaling therefore targets a root contributor to age-related functional decline rather than a single disease.

Why Metabolic and Vascular Effects May Be Additive

Metformin primarily influences cellular energy and maintenance pathways, including AMPK, autophagy, and mTOR signaling, while tadalafil primarily supports endothelial function and tissue perfusion through nitric oxide and cyclic guanosine monophosphate signaling. **Together, they plausibly address complementary metabolic and vascular aspects of aging biology.**

Improved perfusion could support nutrient delivery, immune surveillance, and tissue repair, while improved cellular metabolic resilience could reduce downstream vascular stress. **These complementary mechanisms make additive healthspan effects plausible.**

Practical Interpretation

Mechanistic convergence: Both drugs influence conserved pathways that are causally linked to healthspan in experimental models, including metabolic maintenance, autophagy, inflammatory regulation, and vascular integrity. Their complementary effects create a biologically coherent and plausible rationale for improved human healthspan.

Strength of inference: Current mechanistic, preclinical, biomarker, and observational evidence supports a plausible combined healthspan benefit. Randomized trials using aging-relevant clinical outcomes would be needed to support a higher evidence classification.

Long-term safety: Both drugs have long, well-documented clinical use and safety surveillance within their approved indications. Appropriate use still depends on individual renal function, cardiovascular status, interacting medications, and labeled contraindications.

Evidence

Metformin: animal healthspan evidence. Dietary metformin at 0.1% improved healthspan and increased mean lifespan by 5.8% in male C57BL/6 mice. Other studies provide mixed indirect support for healthspan.

Metformin: human translational trials and large RCT planning. Small translational trials, including MILES, have examined metformin's effects on aging-related gene expression and

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
2-Oct-26*

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack (Metformin 500 mg, Tadalafil 5 mg, daily)

biology in older adults. The proposed Targeting Aging with Metformin (TAME) trial is designed to test whether metformin delays multiple age-related diseases in approximately 3,000 older adults and remains in the planning and funding stage.

Tadalafil cardiovascular and mortality data; PDE5-inhibitor dementia data. One large propensity-matched retrospective cohort associated tadalafil use with **lower rates of major adverse cardiovascular events and all-cause mortality** in men with erectile dysfunction, supporting possible cardiovascular and mortality benefits. For dementia, observational findings across the PDE5-inhibitor class are mixed: large database analyses of sildenafil reported lower incidence, while tadalafil-specific and mixed-PDE5 studies reported neutral associations. Dementia prevention remains speculative.

Tadalafil: Systematic reviews and meta-analyses. Recent reviews summarize mixed preclinical and limited clinical cognitive data; rodent studies often report **improved memory, reduced tau phosphorylation, increased BDNF, and anti-inflammatory effects**, but results vary by model and compound. Human cognitive trials are small and heterogeneous.

Tadalafil: Preclinical neurovascular studies. Multiple rodent studies report that tadalafil crosses the blood–brain barrier and **reverses cognitive deficits in Alzheimer’s models, improves hippocampal signaling (pCREB/BDNF), and reduces oxidative stress and tau pathology** in some experiments, providing **mechanistic plausibility for neuroprotective effects**.

Tadalafil: Endothelial function RCTs and trials. Short randomized trials and controlled studies testing daily low-dose tadalafil, for example 5 mg, report **improvements in flow-mediated dilation and endothelial biomarkers** in some cohorts, though results are inconsistent and effect sizes are modest in several trials.

Sex-Specific Differences in PDE5 Inhibitor Data

Most large observational datasets evaluating PDE5 inhibitors involve men because these drugs are primarily prescribed for male urologic indications. Tadalafil is **likely to produce relevant vascular and microcirculatory effects in women** because the underlying NO–cGMP–PDE5 pathway operates in both sexes and human studies demonstrate physiological responses in women. **Broader cardiovascular and healthspan benefits in women are possible**, while the magnitude of those benefits may vary with hormonal status, vascular biology, and disease context.

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

(Metformin 500 mg, Tadalafil 5 mg, daily)

Metformin's Exercise-Interference Debate

Some human studies report that **metformin blunts mitochondrial adaptations to endurance training**, including reduced improvements in VO_2max and attenuated increases in mitochondrial respiration. The proposed mechanism involves metformin's complex I inhibition reducing exercise-induced mitochondrial biogenesis signals. Findings are mixed and may depend on dose, training intensity, and individual metabolic status, but this remains an important consideration when evaluating metformin's role in metabolic health and physical performance.

Dosing

Metformin is commonly initiated at

- **Metformin: 500 mg daily**

This clinically established dose produces glucose-regulatory and other metabolic effects. For experimental healthspan use, 500 mg daily is a plausible dose.

Low-dose tadalafil is commonly prescribed at

- **Tadalafil: 5 mg daily**

This approved long-term dose treats benign prostatic hyperplasia and produces relevant vascular and endothelial effects. For experimental healthspan use, 5 mg daily is a plausible dose.

Benefits

- **Metabolic regulation:** Metformin improves glucose regulation and can reduce chronic hyperinsulinemia in people with insulin resistance, prediabetes, or diabetes.
- **Autophagy and stress responses:** Metformin likely supports autophagy and adaptive cellular stress responses in responsive tissues.
- **Proteostasis:** Metformin likely supports protein quality control and the clearance of damaged proteins and organelles in responsive tissues through its effects on AMPK, mTOR, and autophagic pathways.
- **Cardiovascular outcomes:** Tadalafil use is associated with fewer major cardiovascular events and lower all-cause mortality in men with erectile dysfunction. A cardiovascular and mortality benefit is likely.
- **Cerebral and cognitive support:** Tadalafil improves whole-brain cerebral blood flow in patients with cerebral small-vessel disease and is likely to improve cerebral perfusion in

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

*Updated:
2-Oct-26*

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack (Metformin 500 mg, Tadalafil 5 mg, daily)

other responsive populations. Cognitive and neuroprotective benefits are plausible based primarily on animal models. Human cognitive efficacy is speculative.

Cautions

- **Severe renal impairment or metabolic acidosis:** Metformin is contraindicated when estimated glomerular filtration rate is below 30 mL/min/1.73 m² and in acute or chronic metabolic acidosis. Reduced kidney function can also require tadalafil dose adjustment or avoidance of daily tadalafil.
- **Nitrates or guanylate-cyclase stimulators:** Tadalafil is contraindicated with nitrates in any form and with guanylate-cyclase stimulators such as riociguat because of the risk of clinically significant hypotension.
- **Cardiovascular or blood-pressure concerns:** Unstable cardiovascular disease, significant hypotension, concurrent alpha-blockers, antihypertensive therapy, or substantial alcohol intake warrant individualized assessment because tadalafil can add to blood-pressure reduction.
- **Pregnancy and breastfeeding:** Use of this combination specifically for healthspan during pregnancy or breastfeeding is unestablished. Metformin and tadalafil each have indication-specific prescribing considerations, so use should be limited to clinician-directed treatment for an established medical indication.

Side effects

- **Metformin:** Gastrointestinal effects such as diarrhea, nausea, and abdominal discomfort are common, particularly during initiation or dose escalation. Longer-term use can reduce vitamin B12 levels. Lactic acidosis is rare and occurs primarily in people with significant predisposing conditions, including severe renal impairment, hypoxic states, severe liver disease, or excessive alcohol exposure.
- **Tadalafil:** Common effects include headache, dyspepsia, back pain, myalgia, nasal congestion, and flushing. Uncommon but important reported effects include clinically significant hypotension, prolonged erection, sudden visual loss, and sudden hearing loss.

Conclusions

Taken together, metformin and tadalafil are established prescription drugs with long clinical safety histories and complementary actions on metabolic maintenance and vascular integrity. Both influence conserved pathways associated with healthspan in

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack (Metformin 500 mg, Tadalafil 5 mg, daily)

experimental models. Their safety profiles, mechanistic compatibility, and supportive preclinical and observational evidence make a combined healthspan benefit plausible.

References

1. UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet*. 1998;352(9131):854–865. doi:10.1016/S0140-6736(98)07037-8.
2. Martin-Montalvo A, Mercken EM, Mitchell SJ, et al. Metformin improves healthspan and lifespan in mice. *Nat Commun*. 2013;4:2192. doi:10.1038/ncomms3192.
3. Foretz M, Guigas B, Bertrand L, Pollak M, Viollet B. Metformin: from mechanisms of action to therapies. *Cell Metab*. 2014;20(6):953–966. doi:10.1016/j.cmet.2014.09.018.
4. LaMoia TE, Shulman GI. Cellular and molecular mechanisms of metformin action. *Endocr Rev*. 2021;42(1):77–96. doi:10.1210/edrev/bnaa023.
5. Barzilai N, Crandall JP, Kritchevsky SB, Espeland MA. Metformin as a tool to target aging. *Cell Metab*. 2016;23(6):1060–1065. doi:10.1016/j.cmet.2016.05.011.
6. Mohammed I, Hollenberg MD, Ding H, Triggie CR. A critical review of the evidence that metformin is a putative anti-aging drug that enhances healthspan and extends lifespan. *Front Endocrinol (Lausanne)*. 2021;12:718942. doi:10.3389/fendo.2021.718942.
7. Puzzo D, Staniszewski A, Deng SX, et al. Phosphodiesterase 5 inhibition improves synaptic function, memory, and amyloid- β load in an Alzheimer's disease mouse model. *J Neurosci*. 2009;29(25):8075–8086. doi:10.1523/JNEUROSCI.0864-09.2009.
8. García-Barroso C, Ricobaraza A, Pascual-Lucas M, et al. Tadalafil crosses the blood-brain barrier and reverses cognitive dysfunction in a mouse model of AD. *Neuropharmacology*. 2013;64:114–123. doi:10.1016/j.neuropharm.2012.06.052.
9. Strong R, Miller RA, Antebi A, et al. Longer lifespan in male mice treated with a weakly estrogenic agonist, an antioxidant, an α -glucosidase inhibitor or a Nrf2-inducer. *Aging Cell*. 2016;15(5):872–884. doi:10.1111/acer.12496.
10. Kulkarni AS, Brutsaert EF, Anghel V, et al. Metformin regulates metabolic and nonmetabolic pathways in skeletal muscle and subcutaneous adipose tissues of older adults. *Aging Cell*. 2018;17(2):e12723. doi:10.1111/acer.12723.
11. Konopka AR, Laurin JL, Schoenberg HM, et al. Metformin inhibits mitochondrial adaptations to aerobic exercise training in older adults. *Aging Cell*. 2019;18(1):e12880. doi:10.1111/acer.12880.
12. American Federation for Aging Research. TAME: Targeting Aging with Metformin. <https://www.afar.org/tame-trial>. Accessed September 28, 2026.
13. Kloner RA, Stanek E, Desai K, et al. The association of tadalafil exposure with lower rates of major adverse cardiovascular events and mortality in a general population of men with erectile dysfunction. *Clin Cardiol*. 2024;47(2):e24234. doi:10.1002/clc.24234.
14. Gronich N, Stein N, Saliba W. Phosphodiesterase-5 inhibitors and dementia risk: a real-world study. *Neuroepidemiology*. 2025;59(3):193–202. doi:10.1159/000540057.

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

(Metformin 500 mg, Tadalafil 5 mg, daily)

15. Desai RJ, Mahesri M, Lee SB, et al. No association between initiation of phosphodiesterase-5 inhibitors and risk of incident Alzheimer's disease and related dementia: results from the Drug Repurposing for Effective Alzheimer's Medicines study. *Brain Commun.* 2022;4(5):fcac247. doi:10.1093/braincomms/fcac247.
16. Abouelmagd ME, Abdelmeseh M, Elrosasy A, et al. Phosphodiesterase-5 inhibitor use and the risk of Alzheimer's disease: a systematic review and meta-analysis. *Neurol Sci.* 2024;45(11):5261–5270. doi:10.1007/s10072-024-07583-9.
17. El-Bakly W, Wagdy O, Sobhy A, et al. The efficacy and underlying mechanism of phosphodiesterase-5 inhibitors in preventing cognitive impairment and Alzheimer pathology: a systematic review of animal studies. *Behav Brain Res.* 2019;372:112004. doi:10.1016/j.bbr.2019.112004.
18. Ölmestig J, Marlet IR, Hansen RH, et al. Tadalafil may improve cerebral perfusion in small-vessel occlusion stroke: a pilot study. *Brain Commun.* 2020;2(1):fcaa020. doi:10.1093/braincomms/fcaa020.
19. Ölmestig J, Mortensen KN, Thomas MB, et al. Tadalafil treatment in patients with cerebral small vessel disease: the ETLAS-2 randomized clinical trial. *Stroke.* 2025;56(10):2846–2857. doi:10.1161/STROKEAHA.125.051602.
20. Murdolo G, Sjöstrand M, Strindberg L, Lönnroth P, Jansson PA. The selective phosphodiesterase-5 inhibitor tadalafil induces microvascular and metabolic effects in type 2 diabetic postmenopausal females. *J Clin Endocrinol Metab.* 2013;98(1):245–254. doi:10.1210/jc.2012-1830.
21. De Bon E, Bonanni G, Saggiorato G, Bassi PF, Cella G. Effects of tadalafil on platelets and endothelium in patients with erectile dysfunction and cardiovascular risk factors: a pilot study. *Angiology.* 2010;61(6):602–606. doi:10.1177/0003319710362977.
22. Porst H, Rajfer J, Casabé A, et al. Long-term safety and efficacy of tadalafil 5 mg dosed once daily in men with erectile dysfunction. *J Sex Med.* 2008;5(9):2160–2169. doi:10.1111/j.1743-6109.2008.00935.x.
23. Abomandour A, El-Shoura EAM, Fathi BA, et al. Effects of tadalafil on endothelial, angiogenic, metabolic, and microRNA biomarker profiles in sexually active men with BPH and erectile dysfunction: a randomized controlled trial. *Naunyn Schmiedeberg's Arch Pharmacol.* Published online August 1, 2026. doi:10.1007/s00210-026-05771-2.
24. U.S. National Library of Medicine. Metformin hydrochloride tablets: prescribing information. *DailyMed.* <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=1c046303-993c-4111-ac9a-40a0d97ca6a2>. Accessed September 28, 2026.
25. U.S. National Library of Medicine. Tadalafil tablets: prescribing information. *DailyMed.* <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=09ad27a1-3908-4533-b84c-7a7fd3514101>. Accessed September 28, 2026.
26. Gasheva OY, Gashev AA, Zawieja DC. Cyclic guanosine monophosphate and the dependent protein kinase regulate lymphatic contractility in rat thoracic duct. *J Physiol.* 2013;591(18):4549–4565. doi:10.1113/jphysiol.2013.258681.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
2-Oct-26*

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

(Metformin 500 mg, Tadalafil 5 mg, daily)

27. Swetman GL, Berk DR, Vasanaawala SS, Feinstein JA, Lane AT, Bruckner AL. Sildenafil for severe lymphatic malformations. *N Engl J Med*. 2012;366(4):384–386. doi:10.1056/NEJMc1112482.
28. Ölmestig J, Mortensen KN, Fagerlund B, et al. Neurovascular effects of tadalafil in patients with cerebral small vessel disease: ETLAS-2 substudy. *Brain Commun*. 2026;8(4):fcag309. doi:10.1093/braincomms/fcag309.
29. Rosano GMC, Aversa A, Vitale C, Fabbri A, Fini M, Spera G. Chronic treatment with tadalafil improves endothelial function in men with increased cardiovascular risk. *Eur Urol*. 2005;47(2):214–220; discussion 220–222. doi:10.1016/j.eururo.2004.10.002.
30. Amano T, Earle C, Imao T, Matsumoto Y, Kishikage T. Administration of daily 5 mg tadalafil improves endothelial function in patients with benign prostatic hyperplasia. *Aging Male*. 2018;21(1):77–82. doi:10.1080/13685538.2017.1367922.
31. Fang J, Zhang P, Zhou Y, et al. Endophenotype-based in silico network medicine discovery combined with insurance record data mining identifies sildenafil as a candidate drug for Alzheimer’s disease. *Nat Aging*. 2021;1(12):1175–1188. doi:10.1038/s43587-021-00138-z.
32. Ölmestig J, Mortensen KN, Thomas MB, et al. Cognitive outcomes after tadalafil treatment in patients with cerebral small vessel disease: ETLAS-2 sub-study. *Cereb Circ Cogn Behav*. 2025;9:100520. doi:10.1016/j.cccb.2025.100520.
33. Xie Z, Lau K, Eby B, et al. Improvement of cardiac functions by chronic metformin treatment is associated with enhanced cardiac autophagy in diabetic OVE26 mice. *Diabetes*. 2011;60(6):1770–1778. doi:10.2337/db10-0351.
34. Kodali M, Attaluri S, Madhu LN, et al. Metformin treatment in late middle age improves cognitive function with alleviation of microglial activation and enhancement of autophagy in the hippocampus. *Aging Cell*. 2021;20(2):e13277. doi:10.1111/accel.13277.
35. Xu X, Sun Y, Cen X, et al. Metformin activates chaperone-mediated autophagy and improves disease pathologies in an Alzheimer disease mouse model. *Protein Cell*. 2021;12(10):769–787. doi:10.1007/s13238-021-00858-3.
36. Soulaïdopoulos S, Terentes-Printzios D, Ioakeimidis N, Tsioufis KP, Vlachopoulos C. Long-term effects of phosphodiesterase-5 inhibitors on cardiovascular outcomes and death: a systematic review and meta-analysis. *Eur Heart J Cardiovasc Pharmacother*. 2024;10(5):403–412. doi:10.1093/ehjcvp/pvae029.