

Tesamorelin + Ipamorelin For Research Use Only

(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

Tesamorelin is a growth hormone-releasing hormone (GHRH) analog that reduces visceral fat and fatty liver disease. Ipamorelin is a selective ghrelin-receptor agonist that produces a discrete growth-hormone pulse when administered intravenously in humans. Because the two peptides stimulate growth-hormone release through complementary receptors, combining them is likely to increase endogenous growth hormone and insulin-like growth factor 1. Community reports often describe the combination as producing a stronger, smoother, and more physiologic-feeling response, with improved recovery, body composition, and tolerability, but these combination-specific effects remain speculative.

 Short-term safety is likely. Long-term safety is plausible.

★ ★ ★ Tesamorelin increases GH and IGF-1 and reduces visceral fat.

★ ★ The combination is likely to increase GH and IGF-1 because both components independently do so through complementary receptors. This plausibly creates benefits for recovery, fitness, and generalized body recomposition.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 13-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/tesamorelin-plus-ipamorelin.pdf>



Dosage & Patterns

The pivotal phase 3 tesamorelin trials used 2 mg subcutaneously once daily for 26 weeks to reduce visceral fat in adults with HIV-associated abdominal fat accumulation.

The following speculative dosing approaches are commonly discussed:

- **Tesamorelin 2 mg with ipamorelin 250 mcg.**
- **Tesamorelin 0.5 - 1 mg with ipamorelin 100 - 250 mcg**, often selected in community practice in an effort to improve tolerability.
- Reported schedules include **once daily, five days per week**, or **every other day**.
- Cycles are often **12-weeks ON**, followed by **6-weeks OFF**, but other cycling patterns are also used.

Some community protocols use 12–24-week cycles followed by 4–12-week breaks; this cycling practice is speculative. Clinical trials also evaluated continuous tesamorelin treatment for up to 52 weeks, during which visceral-fat reductions were maintained. Anti-tesamorelin antibodies

Tesamorelin + Ipamorelin For Research Use Only

(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

developed in approximately half of treated participants, while mean visceral-fat reductions and IGF-1 responses remained similar in participants with and without antibodies.

Benefits

Tesamorelin has demonstrated human benefits for visceral fat, liver fat, lipid profiles, lean body mass, muscle composition, and cognition in the populations studied. Ipamorelin is justified mechanistically to likely increase the endogenous GH pulse.

A 12-week randomized, placebo-controlled trial in adults with type 2 diabetes compared tesamorelin 1 mg and 2 mg subcutaneously once daily. Both doses significantly increased IGF-1 relative to placebo, with mean increases of 33 ng/mL at 1 mg and 66 ng/mL at 2 mg. Insulin response, fasting glucose, overall diabetes control, and body-composition measures remained similar among the groups over 12 weeks.

It is likely these benefits will apply to the general population.

Intravenous ipamorelin produces a discrete, concentration-dependent GH pulse in humans.

- **Visceral fat reduction:** In adults with HIV-associated abdominal fat accumulation, tesamorelin 2 mg subcutaneously once daily produced a pooled placebo-adjusted 15.4% reduction in visceral adipose tissue at 26 weeks. It also reduced waist circumference and waist-to-hip ratio while largely preserving abdominal subcutaneous fat.
- **Improved lipid profile:** In pooled phase 3 trials involving adults with HIV-associated abdominal fat accumulation, tesamorelin reduced triglycerides and improved the ratio of total cholesterol to high-density lipoprotein cholesterol.
- **Reduced liver fat:** In a 12-month randomized trial involving people with HIV and nonalcoholic fatty liver disease, tesamorelin reduced liver fat and reduced progression of liver fibrosis.
- **Improved executive function:** In a 20-week randomized trial involving healthy older adults and older adults with mild cognitive impairment, tesamorelin 1 mg daily improved overall cognition and executive function. Delayed verbal recall also improved in the subgroup with mild cognitive impairment.
- **Increased lean body mass:** In a 12-week randomized dose-ranging trial involving adults with HIV-associated abdominal fat accumulation, tesamorelin 1 mg and 2 mg daily increased lean body mass relative to placebo.
- **Improved muscle composition:** Among adults with HIV-associated abdominal fat accumulation who responded to tesamorelin, trunk-muscle density and lean muscle area increased over 26 weeks.
- **Physiologic GH stimulation:** Tesamorelin increases basal and pulsatile endogenous growth-hormone secretion in humans, while intravenous ipamorelin produces a discrete growth-

Tesamorelin + Ipamorelin **For Research Use Only** (0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

hormone pulse. Their complementary receptor actions are likely to produce pulsatile endogenous growth-hormone stimulation when combined.

Potential Applications

- **Excess visceral fat:** Tesamorelin reduces visceral fat in adults with HIV-associated abdominal fat accumulation and in adults with abdominal obesity and reduced growth-hormone secretion.
- **Fatty liver:** Tesamorelin reduces liver fat and fibrosis progression in people with HIV and nonalcoholic fatty liver disease.
- **Broader GH-axis support:** Adding ipamorelin is likely to increase growth hormone and insulin-like growth factor 1 through complementary receptor activity and could plausibly **support recovery, fitness, and generalized body recomposition.**

Cautions

- **Previous malignancy:** A preexisting malignancy should be inactive and its treatment complete before tesamorelin is considered.
- **Pregnancy or Breastfeeding:** Tesamorelin should be avoided during pregnancy or breastfeeding.
- **Diabetes or glucose intolerance:** Tesamorelin can cause glucose intolerance or diabetes, so glucose status should be assessed before and during use.
- **Persistently elevated IGF-1:** Reassess use when insulin-like growth factor 1 remains above the age-adjusted range. Discontinuation should be considered when it remains more than three standard deviations above the age-adjusted mean.
- **Concerning growth-related symptoms:** Unexplained masses, rapid tissue growth, or other concerning symptoms warrant clinical assessment and reassessment of continued use.

Side Effects

- **Fluid retention:** Peripheral edema can occur and is often most apparent early in treatment.
- **Musculoskeletal symptoms:** Joint pain and muscle pain can occur.
- **Headache and nausea:** Both have been reported in clinical studies.
- **Changes in sleep:** Deeper sleep and vivid dreams are reported in peptide communities and remain speculative as combination-specific effects.

Cancer Risk

Tesamorelin increases circulating insulin-like growth factor 1. After 26 weeks of treatment, 47% of participants had levels more than two standard deviations above the age-adjusted mean and 36% had levels more than three standard deviations above the mean. At 52 weeks, these proportions were 34% and 23%, respectively. Insulin-like growth factor 1 participates in cellular proliferation, and higher circulating levels are epidemiologically associated with several cancers.

Tesamorelin + Ipamorelin For Research Use Only

(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

It is therefore plausible that sustained growth-hormone and insulin-like-growth-factor-1 elevation could promote the growth of an occult or recurrent malignancy.

Drivers of Diminished Response and Mitigation Strategies

For **Receptor desensitization** both GHRH receptor agonism and ghrelin receptor stimulation can produce reduced responsiveness with continuous exposure as described in pituitary physiology literature. For **Neutralizing antibodies** tesamorelin clinical trial reports document anti-drug antibodies including neutralizing antibodies in a minority of subjects, so antibody mediated loss is a real documented risk. For **downstream pathway adaptation** IGF-1 feedback and intracellular signaling adjustments are expected and reduce efficacy over time. For **System Level Physiological Adaptation** compensatory metabolic and endocrine changes further limit sustained benefit. **Overall likelihood of clinically meaningful waning is high.**

Mitigation supported by the evidence is **cycling** therapy, monitoring **IGF-1**, and testing for **Neutralizing antibodies** if clinical efficacy declines.

Biological Mechanism

Tesamorelin and ipamorelin stimulate pituitary somatotrophs through distinct receptors. Tesamorelin is a stabilized analog of growth-hormone-releasing hormone that activates the growth-hormone-releasing-hormone receptor. Ipamorelin is a selective agonist of growth-hormone-secretagogue receptor 1a. Because both agents independently stimulate growth-hormone release in humans, combining them is likely to increase endogenous growth-hormone output through complementary receptor activity. Preservation of normal endocrine feedback during combined use is mechanistically plausible.

Receptor Binding and Intracellular Signaling

Tesamorelin activates growth-hormone-releasing-hormone receptors on pituitary somatotrophs. These receptors couple primarily to the stimulatory G protein Gs, increasing adenylyl cyclase activity, cyclic adenosine monophosphate, and protein kinase A signaling that supports growth-hormone synthesis and secretion. Ipamorelin activates growth-hormone-secretagogue receptor 1a, which couples primarily to Gq/11 proteins and activates phospholipase C, inositol trisphosphate, and intracellular calcium signaling. These pathways converge on the cellular machinery controlling growth-hormone release, making an additive secretory effect from combined activation mechanistically plausible.

Pituitary Dynamics and GH Pulsatility

Tesamorelin increases basal and pulsatile endogenous growth-hormone secretion in humans. Intravenous ipamorelin produces a discrete growth-hormone pulse that peaks rapidly and then declines. Combining the two agents is likely to increase total endogenous growth-hormone output through complementary receptor activity. A larger pulse amplitude, modestly extended

Tesamorelin + Ipamorelin For Research Use Only

(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

pulse duration, and continued regulation by insulin-like growth factor 1 and hypothalamic somatostatin are mechanistically plausible.

Peripheral Growth-Hormone and IGF-1 Effects

Tesamorelin-induced growth-hormone pulses activate hepatic growth-hormone receptors and increase circulating insulin-like growth factor 1. Human trials show reductions in visceral fat, liver fat, and triglycerides, together with increases in lean body mass. These effects could plausibly involve increased lipolysis and lipid oxidation, altered very-low-density-lipoprotein metabolism, growth-hormone-receptor signaling through Janus kinase 2, signal transducer and activator of transcription 5, and mitogen-activated protein kinase pathways, and insulin-like-growth-factor-1 signaling that supports protein synthesis and mitochondrial function.

Pharmacokinetics and Enzymatic Stability

Tesamorelin contains an amino-terminal trans-3-hexenoyl modification that increases resistance to enzymatic degradation and improves stability relative to native growth-hormone-releasing hormone. In human intravenous studies, ipamorelin had an approximately two-hour terminal half-life and produced a growth-hormone response that peaked at about 40 minutes and declined to low levels within six hours. These distinct properties make a temporally complementary pharmacodynamic profile during combined use plausible.

Conclusions

Tesamorelin increases growth hormone and insulin-like growth factor 1 and has demonstrated human benefits for visceral fat, liver fat, lipid profiles, lean body mass, muscle composition, and cognition in the populations studied. Intravenous ipamorelin produces a discrete growth-hormone pulse, and combining the two agents is likely to increase endogenous growth hormone and insulin-like growth factor 1 through complementary receptor activity. Greater overall efficacy, a smoother pulse pattern, and better tolerability are widely reported in peptide communities but remain speculative as combination-specific effects.

References

Tesamorelin clinical trials and human studies

Falutz J, Allas S, Kotler D, et al. A placebo-controlled, dose-ranging study of a growth hormone-releasing factor in HIV-infected patients with abdominal fat accumulation. *AIDS*. 2005;19(12):1279-1287. doi:10.1097/01.aids.0000180099.35146.30.

Falutz J, Allas S, Blot K, et al. Metabolic effects of a growth hormone-releasing factor in patients with HIV. *N Engl J Med*. 2007;357(23):2359-2370. doi:10.1056/NEJMoa072375.

Falutz J, Allas S, Mamputu JC, et al. Long-term safety and effects of tesamorelin, a growth hormone-releasing factor analogue, in HIV patients with abdominal fat accumulation. *AIDS*. 2008;22(14):1719-1728. doi:10.1097/QAD.0b013e32830a5058.

Tesamorelin + Ipamorelin For Research Use Only
(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

Falutz J, Potvin D, Mamputu JC, et al. Effects of tesamorelin, a growth hormone-releasing factor, in HIV-infected patients with abdominal fat accumulation: a randomized placebo-controlled trial with a safety extension. J Acquir Immune Defic Syndr. 2010;53(3):311-322.

doi:10.1097/QAI.0b013e3181cbdaff.

Falutz J, Mamputu JC, Potvin D, et al. Effects of tesamorelin in human immunodeficiency virus-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind, placebo-controlled phase 3 trials with safety-extension data. J Clin Endocrinol Metab. 2010;95(9):4291-4304. doi:10.1210/jc.2010-0490.

Stanley TL, Feldpausch MN, Oh J, et al. Effect of tesamorelin on visceral fat and liver fat in HIV-infected patients with abdominal fat accumulation: a randomized clinical trial. JAMA. 2014;312(4):380-389. doi:10.1001/jama.2014.8334.

Stanley TL, Fourman LT, Feldpausch MN, et al. Effects of tesamorelin on non-alcoholic fatty liver disease in HIV: a randomised, double-blind, multicentre trial. Lancet HIV. 2019;6(12):e821-e830. doi:10.1016/S2352-3018(19)30338-8.

Makimura H, Feldpausch MN, Rope AM, et al. Metabolic effects of a growth hormone-releasing factor in obese subjects with reduced growth hormone secretion: a randomized controlled trial. J Clin Endocrinol Metab. 2012;97(12):4769-4779. doi:10.1210/jc.2012-2794.

Koutkia P, Canavan B, Breu J, Torriani M, Kissko J, Grinspoon S. Growth hormone-releasing hormone in HIV-infected men with lipodystrophy: a randomized controlled trial. JAMA. 2004;292(2):210-218. doi:10.1001/jama.292.2.210.

Stanley TL, Chen CY, Branch KL, et al. Effects of a growth hormone-releasing hormone analog on endogenous GH pulsatility and insulin sensitivity in healthy men. J Clin Endocrinol Metab. 2011;96(1):150-158. doi:10.1210/jc.2010-1587.

Clemmons DR, Miller S, Mamputu JC. Safety and metabolic effects of tesamorelin, a growth hormone-releasing factor analogue, in patients with type 2 diabetes: a randomized, placebo-controlled trial. PLoS One. 2017;12(6):e0179538. doi:10.1371/journal.pone.0179538.

Adrian S, Scherzinger A, Sanyal A, et al. The growth hormone-releasing hormone analogue, tesamorelin, decreases muscle fat and increases muscle area in adults with HIV. J Frailty Aging. 2019;8(3):154-159. doi:10.14283/jfa.2018.45.

Lake JE, La K, Erlandson KM, et al. Tesamorelin improves fat quality independent of changes in fat quantity. AIDS. 2021;35(9):1395-1402. doi:10.1097/QAD.0000000000002897.

Stanley TL, Falutz J, Mamputu JC, Soulban G, Potvin D, Grinspoon SK. Effects of tesamorelin on inflammatory markers in HIV patients with excess abdominal fat: relationship with visceral adipose reduction. AIDS. 2011;25(10):1281-1288. doi:10.1097/QAD.0b013e31828347f3f1.

Tesamorelin + Ipamorelin For Research Use Only

(0.5 - 2 mg, + 100 - 250 mcg 3 – 7 days / week, fasted, subq)

Fourman LT, Billingsley JM, Agyapong G, et al. Effects of tesamorelin on hepatic transcriptomic signatures in HIV-associated NAFLD. JCI Insight. 2020;5(16):e140134. doi:10.1172/jci.insight.140134.

Russo SC, Ockene MW, Arpante AK, et al. Efficacy and safety of tesamorelin in people with HIV on integrase inhibitors. AIDS. 2024;38(12):1758-1764. doi:10.1097/QAD.0000000000003965.

Baker LD, Barsness SM, Borson S, Merriam GR, Friedman SD, Craft S, Vitiello MV. Effects of growth hormone-releasing hormone on cognitive function in adults with mild cognitive impairment and healthy older adults: results of a controlled trial. Arch Neurol. 2012;69(11):1420-1429. doi:10.1001/archneurol.2012.1970.

Ipamorelin human evidence

Gobburu JV, Agersø H, Jusko WJ, Ynddal L. Pharmacokinetic-pharmacodynamic modeling of ipamorelin, a growth hormone-releasing peptide, in human volunteers. Pharm Res. 1999;16(9):1412-1416. doi:10.1023/A:1018955126402.

Beck DE, Sweeney WB, McCarter MD; Ipamorelin 201 Study Group. Prospective, randomized, controlled, proof-of-concept study of the ghrelin mimetic ipamorelin for the management of postoperative ileus in bowel resection patients. Int J Colorectal Dis. 2014;29(12):1527-1534. doi:10.1007/s00384-014-2030-8.

Ipamorelin animal and mechanistic evidence

Raun K, Hansen BS, Johansen NL, et al. Ipamorelin, the first selective growth hormone secretagogue. Eur J Endocrinol. 1998;139(5):552-561. doi:10.1530/eje.0.1390552.

Svensson J, Lall S, Dickson SL, et al. The GH secretagogues ipamorelin and GH-releasing peptide-6 increase bone mineral content in adult female rats. J Endocrinol. 2000;165(3):569-577. doi:10.1677/joe.0.1650569.

Andersen NB, Malmlöf K, Johansen PB, Andreassen TT, Ørtoft G, Oxlund H. The growth hormone secretagogue ipamorelin counteracts glucocorticoid-induced decrease in bone formation of adult rats. Growth Horm IGF Res. 2001;11(5):266-272. doi:10.1054/ghir.2001.0239.

Venkova K, Mann W, Nelson R, Greenwood-Van Meerveld B. Efficacy of ipamorelin, a novel ghrelin mimetic, in a rodent model of postoperative ileus. J Pharmacol Exp Ther. 2009;329(3):1110-1116. doi:10.1124/jpet.108.149211.

Related IGF-1 epidemiology

Qian F, Huo D. Circulating insulin-like growth factor-1 and risk of total and 19 site-specific cancers: cohort study analyses from the UK Biobank. Cancer Epidemiol Biomarkers Prev. 2020;29(11):2332-2342. doi:10.1158/1055-9965.EPI-20-0743.