

KLOW Blend (GHK-CU / BPC-157 / TB-500 / KPV) Available for Research Use Only

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

KLOW is a peptide blend that combines GHK-Cu, BPC-157, TB-500, and KPV, intended to support skin quality, soft tissue healing, mucosal integrity, and a low-noise systemic anti-inflammatory tone. Because TB-500 promotes angiogenesis and cell migration, KLOW is contraindicated in subjects with active cancer or elevated cancer risk. For subjects with elevated cancer risk, consider using GHK-Cu with or without BPC-157 and KPV instead of the full KLOW blend.

💉💉 Probably safe (some human evidence)

C Angiogenic – caution for subjects at elevated risk for cancers, Do NOT use with active cancer

★ ★ Probably effective (strong animal signals and/or some positive human signals)

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 31-Jul-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/klow-blend.pdf>



KLOW typically uses a 50/10/10/10 ratio of **GHK-Cu**, **TB-500**, **BPC-157**, and **KPV** respectively. The rationale is to combine **GHK-Cu** gene-level regenerative signaling and copper delivery, with **TB-500** actin and migration support, **BPC-157** endothelial and gut-vascular stabilization, and **KPV** anti-inflammatory and epithelial barrier support. Together these signals aim to improve extracellular matrix quality, microvascular perfusion, tissue remodeling, and mucosal integrity.

The evidence for Klow's long term human safety is moderately weak because there are **no peer-reviewed subcutaneous human trials; Safety confidence rests on mechanistic plausibility, in-vitro and animal tolerability, anecdotal clinical use, and widespread community use.**

The evidence base for KLOW as a subcutaneous therapy is limited because no peer reviewed human trials evaluate the full blend, or most of its components. Confidence rests on mechanistic plausibility, in vitro and animal data for each component, and experiential reports from clinical compounding and research use. **GHK-Cu** has the strongest translational human signal for skin and topical effects and robust mechanistic gene expression data. **BPC-157** and **TB-500** have strong animal and mechanistic evidence for tissue repair and angiogenesis but limited controlled human data. **KPV** has reproducible preclinical anti-inflammatory and barrier data and growing experiential use. The combined blend is mechanistically coherent but lacks controlled human subcutaneous trials.

KLOW Blend (GHK-CU / BPC-157 / TB-500 / KPV) **Available for Research Use Only** (1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

Dosage & Patterns

In the 50/10/10/10 blend KLOW dosing ratio of GHK-cu, TB-500, BPC-157, and KPV are limited to that ratio. There is no human testing to define least effective dose or maximum safe dose. Instead, we rely on common practices at med spas and reported by forum users of FRU peptides. The following pattern emerges

- **1.6 mg / daily KLOW, for skin & hair**
 - ~ 1 mg GHK-CU / daily
 - ~ 200 mcg BPC-157 / daily
 - ~ 200 mcg TB-500 / daily
 - ~ 200 mcg KPV / daily
- **3.2 mg / daily KLOW, for tissue repair**
 - ~ 2 mg GHK-CU / daily
 - ~ 400 mcg BPC-157 / daily
 - ~ 400 mcg TB-500 / daily
 - ~ 400 mcg KPV / daily

With the lower dose for skin and hair repair and regeneration, and the higher dose for more aggressive injury and tissue repair.

Benefits

Robust randomized human trials for systemic SC use of these peptides in combination do not exist. These benefits are based on preclinical studies of GHK-Cu, BPC-157, and TB-500, and the biological mechanisms. These benefits are frequently reported by individual research users.

- **Smoother, healthier-looking skin**, with improvements in hydration, elasticity, and overall tone.
- **Faster recovery from soft-tissue strain**, due to actin dynamics, angiogenesis, and fibroblast migration.
- **Reduced joint stiffness and background inflammation**, often noticed in knees, shoulders, and hips as connective-tissue signaling normalizes.
- **Improved healing of minor injuries**, including small tears, overuse irritation, and delayed-onset soreness, due to combined effects on endothelial repair and extracellular-matrix turnover.
- **Calmer old injury sites**, with many users describing less reactivity or “flare-ups” in previously problematic areas.
- **Better tendon and ligament resilience**, especially during higher-volume training blocks or repetitive-stress activities.

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

- **Enhanced microvascular function**, supporting nutrient delivery and tissue oxygenation in areas under mechanical load.
- **More even skin texture**, including subtle reductions in fine lines and improved firmness over several weeks.
- **Generalized tissue “comfort”**, a broad effect people describe as less tightness, fewer aches, and smoother movement patterns.
- **Reduced chronic inflammation**, reduces pro-inflammatory cytokines, calms mast-cell activation, modulates melanocortin pathways
- **Improved intestinal barrier integrity**, helps tighten epithelial junctions and reduce permeability.

Indications

- Chronic soft-tissue irritation
- Older injuries that flare under load
- High training volume
- Age-related changes in skin quality
- Demanding physical jobs or lifestyles
- Recovering from minor strains or overuse
- Low-grade systemic inflammation
- Running GH-axis peptides

Contraindications

- Copper metabolism disorders
- Liver disease
- Active cancer, elevated cancer risk

Side Effects (Rarely)

- Nausea
- Headache
- Fatigue

Drivers of Diminished Response and Mitigation Strategies

For **receptor desensitization** the blend mixes low receptor-tachyphylaxis components with fragments that do not primarily act via receptors so overall receptor desensitization risk is low to moderate. For **neutralizing antibodies** the presence of synthetic fragments BPC-157 and TB-500 raises the blend risk to moderate because fragment immunogenicity is a documented concern in peptide literature. For **downstream pathway adaptation** combined chronic anti-inflammatory signaling and modulation of repair and angiogenesis pathways increase the chance of intracellular attenuation. For **system level physiological adaptation** cumulative exposure can shift tissue repair set points and reduce effect. **Overall likelihood of waning is moderate.**

KLOW Blend (GHK-CU / BPC-157 / TB-500 / KPV) Available for Research Use Only

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

Mitigation is to **cycle** use. A plausible cycling strategy might be to use KLOW for one month, followed by GHK-Cu for one month, followed by one month without either. With no evidence to support any specific cycling plan practitioners are free to devise their own cycle. A slightly less conservative cycle scheme to **reset downstream pathway adaption and system level physiological adaptation** might be

Month 1: KLOW

Month 2: GHK-Cu + KPV, BPC-157 as needed for acute injury

Month 3: KLOW

Month 4: GHK-Cu + KPV, BPC-157 as needed for acute injury

Month 5: KLOW

Month 6: Holiday

KLOW: Plausible Longevity Benefits

The plausibly pro-longevity rationale for the KLOW blend arises from the convergence of **four distinct but complementary biological programs**: GHK-Cu's gene-level regenerative and anti-inflammatory signaling, TB-500's cytoskeletal and angiogenic repair pathways, BPC-157's gastrointestinal, endothelial, and microvascular stabilization effects, and KPV's antiinflammatory effects. Each peptide **targets a different conserved pathway in age-related decline**, and together they form a mechanistically coherent strategy for supporting tissue integrity, repair capacity, and inflammatory balance, domains that strongly influence healthspan.

GHK-Cu contributes a broad gene-reprogramming signal that shifts cellular behavior **away from pro-inflammatory and pro-oxidative states toward regeneration, extracellular-matrix rebuilding, and antioxidant defense**. Age-related declines in GHK-Cu correlate with reduced collagen synthesis, impaired wound healing, and increased inflammatory tone; restoring this signal may help maintain ECM quality, reduce chronic inflammatory burden, and preserve microcirculatory function. These actions **address structural aging at the level of collagen, elastin, glycosaminoglycans, and metalloprotein regulation, supporting the connective-tissue architecture that underpins organ resilience**.

TB-500 adds a second, orthogonal repair axis by **enhancing actin cytoskeleton dynamics, cell migration, and angiogenesis**. These pathways are central to wound closure, tissue remodeling, and microvascular regeneration, processes that slow markedly with age. TB-500's ability to mobilize progenitor cells, improve endothelial repair, and reduce fibrosis suggests a role in **maintaining tissue plasticity and preventing the progressive stiffening and scarring that accompany aging**. When paired with GHK-Cu's ECM-centric gene modulation, TB-500 helps ensure that tissues not only produce structural proteins but also remodel and integrate them effectively.

BPC-157 complements both signals by **stabilizing the gastrointestinal barrier, protecting microvasculature, and modulating nitric-oxide pathways** that influence endothelial tone and

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

perfusion. Because aging is strongly shaped by chronic low-grade inflammation originating from gut permeability, vascular instability, and impaired angiogenic balance, BPC-157's protective effects on the gut-vascular axis may **reduce systemic inflammatory load and improve nutrient delivery, oxygenation, and waste clearance**. Its anti-ulcer, endothelial-protective, and tendon-repair properties add a third dimension to the blend's regenerative profile, supporting both local and systemic resilience.

KPV adds a fourth longevity-relevant axis by providing a sustained reduction in **chronic inflammatory tone** and preserving **epithelial and mucosal barrier integrity**, both of which are central determinants of biological aging. Mechanistically, KPV suppresses **NFκB, TNFα, IL1β, and IL6**, stabilizes **mast cells**, and strengthens **tight-junction proteins** such as occludin and claudins, actions that reduce **systemic inflammatory load** and limit the continuous immune activation that accelerates tissue degeneration across the lifespan. Because **inflammaging** is driven by persistent cytokine signaling, epithelial permeability, and mast-cell hyperreactivity, KPV's ability to **normalize immune tone** and **maintain barrier stability** may help reduce the background inflammatory pressure that erodes **microvascular function, collagen quality, and mitochondrial efficiency** over time. By lowering inappropriate inflammatory signaling without broad immunosuppression, KPV contributes a **regulatory, low-noise immune environment** that supports the regenerative and microvascular benefits of GHK-Cu, TB-500, and BPC-157, creating a more favorable systemic milieu for **long-term tissue resilience and functional longevity**.

Mechanistically, the combined effect is a **coordinated reduction in chronic inflammatory tone, improved microvascular perfusion, enhanced repair efficiency, and preservation of structural protein homeostasis**. These domains (ECM quality, vascular integrity, inflammatory balance, and regenerative capacity) are central determinants of functional aging. *Although large human trials are lacking and systemic longevity claims remain theoretical*, the mechanistic convergence of these four peptides provides biologically coherent reasons to hypothesize that the KLOW blend may support extended healthspan by sustaining repair capacity, reducing inflammatory and oxidative burden, and preserving tissue architecture and microcirculation in ways tightly linked to resilience and functional longevity.

KLOW and Cancer Risk

There is **no demonstrated evidence** that KLOW (GHK-Cu + BPC-157 + TB-500 + KPV) *causes cancer de novo* and there are no mechanistic arguments that any of the four peptides in KLOW might cause cancer in someone **without an existing malignancy**. However, the legitimate concern is **theoretical support of angiogenesis, cell migration, or micro-tumor growth** if an undetected malignancy already exists. The risk is *theoretical and mechanistic*, not demonstrated in humans or animals.

GHK-Cu overall has an **anti-cancer, anti-tumor profile** with **no evidence of carcinogenesis** and **some evidence of protective effects**.

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

- GHK-Cu has **anti-cancer gene-expression effects** in several studies, including suppression of metastasis-related genes and reactivation of apoptosis pathways.
- It has been described as **oncoprotective** in some contexts, with evidence of inhibiting tumor growth in cell and animal models.
- It is mildly pro-angiogenic, but this has **not** been linked to tumor promotion in vivo.

BPC-157 overall has a **theoretical risk based on angiogenesis**, but **animal data is anti-tumor, anti-metastatic**.

- No human trial data for cancer, animal data is mixed neutral to anti-cancer.
- Mechanistically activates **FAK-paxillin** and **VEGFR2**, pathways used in both healing and cancer metastasis.

Contrary to the theoretical tumor-promoting potential of many angiogenic agents, **BPC-157 has been shown to inhibit uncontrolled cell proliferation, downregulate VEGF expression, and counteract VEGF-driven tumorigenesis, including suppression of Ki-67 and VEGF pathway activation.**

KPV's ability to **lower inflammatory tone** and **maintain barrier stability** is **theoretically anti-tumor** because chronic inflammation, epithelial permeability, and mast-cell hyperreactivity are well established accelerators of tumor progression.

TB-500 is the most **mechanistically concerning for cancer** of the four peptides in KLOW. Most of the data we have is about Thymosin β 4, not TB-500. Because TB-500 is a fragment of Thymosin β 4 we extrapolate TB-500 risk from what we know about Thymosin β 4.

- Thymosin β 4 is **overexpressed in several cancers** and associated with metastasis and poor prognosis.
- TB-500 promotes **angiogenesis, cell migration, and actin polymerization**, all of which tumors can exploit.

No evidence shows TB-500 causes cancer in healthy subjects, but the **biological overlap with metastatic pathways is real**. Expert clinical guidance consistently recommends **avoiding TB-500 in active or recent cancer**.

However, there is evidence that might reduce this mechanistic pro-tumor message: In a myocardial infarction model, **Bock-Marquette et al.** showed that thymosin β 4 promoted cardiomyocyte survival and tissue repair **without inducing dysplasia, abnormal proliferation, or tumor formation**, demonstrating neutral oncologic behavior in a high-proliferation context. In corneal wound-healing studies, **Sosne et al. (2002)** found that thymosin β 4 accelerated epithelial migration and reduced inflammation **with no neoplastic changes or abnormal proliferative lesions** in treated ocular tissue. In a follow-up corneal regeneration study, **Sosne et al. (2004)** reported enhanced epithelial repair driven by thymosin β 4 **without evidence of hyperproliferation or tumor-like growth**, confirming neutral behavior in a second ocular model.

KLOW Blend (GHK-CU / BPC-157 / TB-500 / KPV) Available for Research Use Only

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

In dermal wound-healing experiments, **Philp et al.** demonstrated that thymosin β 4 improved re-epithelialization and reduced inflammatory cytokines **while maintaining normal tissue architecture and showing no tumorigenic activity** in skin. In a cardiac progenitor mobilization study, **Smart et al.** showed that thymosin β 4 activated epicardial progenitor cells and supported cardiomyocyte regeneration **without uncontrolled proliferation or neoplastic transformation**, indicating regenerative but non-oncogenic signaling.

If an individual already has an undiagnosed malignancy or micro-metastatic disease, TB-500's **pro-migration and pro-angiogenic signaling** could *theoretically* support tumor progression. In people without cancer, available data do **not** show carcinogenicity, but the mechanistic overlap with metastatic biology justifies a **cautionary stance**, especially for those with a history of cancer, active malignancy, or elevated cancer risk.

Taking the four peptides in KLOW, taken together, we have some *in vivo* data and a good mechanistic understanding which is reassuring. However, **if a preexisting tumor already exists**, the information is mixed, **KLOW may be simultaneously protective, slowing tumor growth, and angiogenic, accelerating tumor growth**. The overall risk profile of KLOW in someone without symptomatic cancer is **low but not zero**.

Conservative Alternatives Around Cancer Risks

For someone without elevated cancer risk, but with a desire to be cautious, three alternative patterns are available

- Using the separate peptide GHK-Cu, which tends to have an anti-tumor profile, is the most conservative alternative to KLOW blend. It delivers many of the skin and hair benefits and some of the tissue remodeling benefits of KLOW, albeit without as much anti-inflammatory, injury and soft tissue support. At the researcher's discretion short courses of BPC-157 and / or TB-500 (or KLOW blend) could be used for specific acute injury recovery.
- Combine GHK-Cu daily with BPC-157 and / or KPV daily or less frequently. This provides additional soft tissue support.
- Cycling KLOW and GHK-Cu is another legitimate, conservative approach which provides additional soft tissue support while minimizing exposure to BPC-157 and TB-500. One plausible cycling strategy is to alternate vials of KLOW and GHK-Cu. KPV, with no cancer risk, could be continued alongside GHK-Cu.

Consider zinc supplementation alongside GHK-Cu

People often **consider zinc supplementation alongside GHK-Cu** because zinc helps maintain balanced metal homeostasis in the body. GHK-Cu delivers copper to enzymes involved in repair, and zinc plays a complementary role by supporting antioxidant defenses, stabilizing cellular signaling, improved immune response, and helping regulate metalloproteins that manage copper-zinc balance. When overall zinc status is low, the body may have more difficulty buffering shifts in copper availability, so **maintaining adequate zinc intake can help keep these**

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

metal-dependent pathways stable while GHK-Cu is being used. The zinc requirement is not large, probably 20 mg daily is sufficient to support the continuous maximum dose of GHK-Cu.

Biological Mechanism

GHK-Cu is described in research as a naturally occurring **copper-binding tripeptide** that functions as a repair and remodeling signal across multiple tissues. Biologically, it binds copper(II) and delivers it to enzymes involved in **collagen cross-linking, antioxidant defense** (such as superoxide dismutase), and mitochondrial energy production, while simultaneously activating gene programs linked to **extracellular-matrix rebuilding, angiogenesis, and wound healing**. Physiologically, GHK-Cu reduces inflammatory signaling through suppression of **NF-κB, TNF-α, and IL-6**, increases the expression of structural proteins like collagen, elastin, and glycosaminoglycans, and promotes fibroblast and keratinocyte migration during tissue repair. It also improves microcirculation by supporting capillary growth and endothelial function and has been shown to modulate over 4,000 human genes toward a regenerative, anti-inflammatory profile. This combination of copper transport, gene-level repair signaling, and inflammation control is why GHK-Cu appears across research on skin quality, connective-tissue healing, and general tissue regeneration.

BPC-157 is described in research as a **cytoprotective, angiogenic, and anti-inflammatory** peptide that operates through several converging biological pathways. It stabilizes the gut-vascular axis by up-regulating **VEGF, FGF-2**, and other growth-repair signals while simultaneously suppressing **TNF-α, IL-6, and NF-κB-driven inflammation**. At the cellular level, BPC-157 enhances **fibroblast migration, collagen organization, and endothelial repair**, which supports tendon, ligament, and soft-tissue healing. It also interacts with the nitric-oxide system to normalize microvascular tone and protect against oxidative stress. In animal and early mechanistic studies, BPC-157 preserves tight-junction integrity in the gut, accelerates angiogenesis in injured tissue, modulates dopamine and serotonin pathways, and promotes rapid healing across muscle, nerve, and gastrointestinal models. Together, these effects position BPC-157 as a broad-spectrum repair peptide that coordinates inflammation control, vascular restoration, and extracellular-matrix rebuilding.

TB-500 is described in research as a **regenerative, actin-modulating peptide** derived from the N-terminal region of thymosin β₄, and its biological effects center on restoring cellular structure, improving tissue repair, and reducing inflammation. It enhances **actin polymerization**, which is essential for cell migration, angiogenesis, and wound closure, and it supports endothelial and fibroblast activity in injured tissue. TB-500 also down-regulates **NF-κB, TNF-α**, and other inflammatory mediators, helping shift damaged tissue from an inflammatory state into a repair-oriented one. In animal and mechanistic studies, it improves **angiogenesis**, accelerates **muscle and tendon healing**, protects cardiac tissue after injury, and supports nerve regeneration by stabilizing cytoskeletal dynamics. Together, these actions position TB-500 as a

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

broad tissue-repair signal that coordinates structural rebuilding, vascular restoration, and inflammation control across multiple organ systems.

KPV exerts its effects through a combination of **melanocortin-receptor signaling** and **direct suppression of inflammatory pathways**, producing a coordinated reduction in cytokine activity and improvement in epithelial barrier integrity. As a C-terminal fragment of α -MSH, it selectively activates **MC1R** on immune cells, keratinocytes, and epithelial tissues, triggering a rise in intracellular cAMP that inhibits **NF- κ B** translocation and reduces transcription of pro-inflammatory mediators such as TNF- α , IL-1 β , IL-6, COX-2, and iNOS. This same signaling dampens mast-cell degranulation and stabilizes histamine pathways, lowering release of histamine, tryptase, and leukotrienes. Through these mechanisms, KPV shifts macrophages toward an M2-like phenotype, reduces neutrophil recruitment, and normalizes immune tone without suppressing antimicrobial defenses.

At the tissue level, KPV improves **epithelial and mucosal barrier function** by upregulating tight-junction proteins like occludin and claudins, reducing epithelial apoptosis, and enhancing keratinocyte and enterocyte migration during wound repair. These effects **decrease intestinal permeability** and **limit antigen translocation**, which in turn **reduces downstream immune activation**. By lowering peripheral cytokines and limiting LPS-driven signaling, KPV indirectly reduces neuroinflammatory input to the CNS, contributing to clearer cognition and **reduced inflammatory fatigue**. Together, these mechanisms create a profile that is anti-inflammatory, immune-regulatory, and barrier-restorative, making KPV particularly relevant in gut, skin, and systemic inflammatory contexts.

Conclusions

Because of TB-500's angiogenesis and theoretical risk of cancer activation some researchers might choose to cycle KLOW, or even to eschew KLOW in favor of GHK-Cu with or without BPC-157 and KPV.

References

- **Pickart L, Margolina A.** Regenerative and protective actions of the GHK-Cu peptide in the light of the new gene data. J Biomater Sci Polym Ed. 2012;23(8):927-940. doi:10.1163/092050611X581516
- **Maquart FX, Bellon G, Chaqour B, et al.** Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. Cell Mol Biol (Noisy-le-grand). 1993;39(7):625-631.
- **Siméon A, Wegrowski Y, Bontemps Y, Maquart FX.** Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. J Invest Dermatol. 2000;115(6):962-968. doi:10.1046/j.1523-1747.2000.00174.x

(1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

- **Sikiric P, Seiwerth S, Rucman R, et al.** Stable gastric pentadecapeptide BPC-157: novel therapy in gastrointestinal tract. *Curr Pharm Des.* 2018;24(18):1990-2000. doi:10.2174/1381612824666180613115106
- **Chang CH, Tsai WC, Hsu YC, et al.** The effect of pentadecapeptide BPC-157 on tendon healing in a rat model. *J Orthop Res.* 2011;29(6):887-893. doi:10.1002/jor.21336
- **Kang EA, Han YM, An JM, et al.** BPC-157 accelerates the healing of acetic acid-induced colitis in rats via modulating nitric oxide synthesis. *Gut Liver.* 2011;5(3):361-366. doi:10.5009/gnl.2011.5.3.36.
- **Goldstein AL, Hannappel E, Sosne G, Kleinman HK.** Thymosin β 4: a multi-functional regenerative peptide. *Ann N Y Acad Sci.* 2007;1112:109-116. doi:10.1196/annals.1415.033
- **Sosne G, Qiu P, Christopherson PL, Wheeler MK.** Thymosin β 4 suppression of corneal NF- κ B: a potential anti-inflammatory pathway. *Exp Eye Res.* 2004;79(4):553-560. doi:10.1016/j.exer.2004.06.025
- **Malinda KM, Sidhu GS, Mani H, et al.** Thymosin β 4 accelerates wound healing. *J Cell Sci.* 1997;110(Pt 5):569-578.
- **Bock-Marquette I, Saxena A, White MD, Dimaio JM, Srivastava D.** Thymosin β 4 mediates cell survival and repair in the injured myocardium. *Nature.* 2004;432(7016):466-472.
- **Sosne G, Qiu P, Christopherson PL, Wheeler MK.** Thymosin β 4 promotes corneal wound healing. *Invest Ophthalmol Vis Sci.* 2002;43(7):2237-2241.
- **Sosne G, Kleinman HK, Ousley GW.** Thymosin β 4 stimulates corneal epithelial cell migration. *Exp Eye Res.* 2004;78(3):403-409.
- **Philp D, Badamchian M, Scheremeta B, et al.** Thymosin β 4 promotes wound repair and reduces inflammation. *J Invest Dermatol.* 2004;123(5):1085-1095.
- **Smart N, Risebro CA, Melville AA, et al.** Thymosin β 4 induces adult epicardial progenitor mobilization and cardiomyocyte regeneration. *Circulation.* 2007;115(14):1725-1735.