Retatrutide

SC once weekly: 2 mg, titrate to 4 / 6 / 9 / 12 mg

Retatrutide produces substantial weight loss, improves glycemic control, and reduces total, visceral, and liver fat in randomized human studies. [2–5,14,29,31] Trials also demonstrate improvement in obesity-related knee-osteoarthritis pain and obstructive sleep apnea. [9,29,30] It is an investigational, long-acting single peptide that activates GLP-1, GIP, and glucagon receptors. [1] Incretin signaling supports appetite reduction and glucose-dependent insulin secretion; glucagon-receptor activity promotes fatty-acid oxidation and likely contributes to increased energy expenditure. [1,6] Common adverse effects are characterized through approximately 80 weeks, with gastrointestinal intolerance and sensory symptoms requiring attention; cardiovascular-event reduction remains unresolved. [2,10,29,31]

Safety and tolerability: Human observations extend to approximately 80 weeks; gastrointestinal intolerance and dysesthesia are documented. [10,29,31]

Efficacy: Effective for weight loss and glycemic improvement in randomized human trials. [2,14,29,31]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Retatrutide has been evaluated in randomized human trials lasting up to 80 weeks. Gastrointestinal adverse effects predominate; dysesthesia and hypotension also occur. [29,31] In TRIUMPH-4, adverse-event discontinuation reached 18.2% with 12 mg versus 4.0% with placebo. [9] Hepatic observations in the 48-week liver-fat substudy were reassuring, supporting plausible continued hepatic tolerability, while dehydration-related kidney injury remains a specific risk. [4,15] Dose-related heart-rate increases and inconclusive cardiovascular-event estimates require separate cardiovascular assessment. [2,10] Rare or delayed thyroid, pancreatic, reproductive, and neurologic risks cannot be assigned the same confidence as common short-term tolerability findings. [12,13,28]

Retatrutide produces large, dose-dependent reductions in body weight. Mean loss with 12 mg was 24.2% at 48 weeks in the Phase 2 obesity trial. [2] Sponsor-reported efficacy-estimand losses were 28.7% at 68 weeks in TRIUMPH-4 and 22.6% at 80 weeks in TRIUMPH-3. [9,10] In TRIUMPH-1, the treatment-regimen estimate, which includes the effects of treatment discontinuation, was 25.0% with 12 mg versus 3.9% with placebo at 80 weeks. [29] These estimates describe different populations and approaches to intercurrent events and should not be treated as head-to-head comparisons.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Retatrutide

SC once weekly: 2 mg, titrate to 4 / 6 / 9 / 12 mg

Retatrutide improves glycemic control and multiple cardiometabolic markers. Human trials show reductions in hemoglobin A1c, triglycerides, non-HDL cholesterol, systolic blood pressure, and metabolic markers associated with insulin resistance. [2,3,6,7,14] Significant reductions in high-sensitivity C-reactive protein and interleukin-6 were observed in the obesity cohort without diabetes, but not in the separate Phase 2 type 2 diabetes cohort. [7]

Retatrutide substantially reduces visceral adipose tissue. MRI-based human data show dose-related reductions in visceral fat together with reductions in abdominal subcutaneous fat, liver fat, body weight, and waist circumference. [4]

Dosing Patterns

In Phase 3 trials, retatrutide is administered subcutaneously once weekly, beginning at 2 mg and increasing every four weeks until the assigned maintenance dose is reached. Studied maintenance doses are 4 mg, 9 mg, and 12 mg. The escalation sequence is 2 mg to 4 mg for the 4 mg target; 2 mg to 4 mg to 6 mg to 9 mg for the 9 mg target; and 2 mg to 4 mg to 6 mg to 9 mg to 12 mg for the 12 mg target. Protocols allow dose reduction for persistent intolerance or excessive weight loss. These are trial regimens. [17]

Gastrointestinal adverse effects are dose-related. Phase 2 findings indicate that beginning at 2 mg rather than 4 mg can improve tolerability. [2] Four-week escalation intervals provide time to assess tolerability and are consistent with the long-acting PK profile. [1,17]

The studied dose is

- **2 mg to 4 mg to 6 mg to 9 mg to 12 mg for the 12 mg target, once weekly**

Speculative community dose and pattern modifications

- **Start at 1 mg**
- **Increase in smaller increments than those used in trials**
- **Divide the weekly amount into two injections**
- **Reduce the dose and continuing on a maintenance dose once reaching goal**

These are all speculative community practices intended to improve tolerability. They differ from the once-weekly trial schedules. [2,17]

Benefits

- **Appetite reduction:** Retatrutide reduces appetite and likely reduces energy intake. [1,2] Greater appetite suppression than semaglutide or tirzepatide remains speculative; descriptions of reduced hunger, cravings, and food noise are subjective community observations.
- **Visceral-fat reduction:** Retatrutide substantially reduces visceral adipose tissue in humans. [4] Its glucagon-receptor activity likely contributes through increased fatty-acid oxidation and

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

energy expenditure, a mechanistic inference from animal experiments and human metabolomic findings. [1,6]

- **Energy expenditure:** Retatrutide likely increases energy expenditure based on animal evidence, supported by human biomarker findings consistent with increased fatty-acid oxidation. [1,6] The human metabolomic analysis did not directly measure energy expenditure; its exploratory mediation estimates describe associations rather than a quantified thermogenic effect. [6]
- **Metabolic flexibility:** Improved metabolic flexibility is plausible based on human biomarker changes associated with increased fatty-acid oxidation and reduced insulin resistance. [6]
- **Glycemic control:** Retatrutide improves hemoglobin A1c and fasting glucose. [3,14,31] In TRANSCEND-T2D-1, 12 mg reduced A1c by 1.94 percentage points versus 0.81 with placebo at 40 weeks in participants managed with diet and exercise. [14]
- **Reduced progression to type 2 diabetes:** Retatrutide is likely to reduce progression from prediabetes to type 2 diabetes, inferred from its substantial effects on weight and glycemic risk markers. [2,6] Tirzepatide's randomized diabetes-prevention results provide indirect support through shared incretin biology. [18]
- **Insulin sensitivity:** Retatrutide improves fasting-insulin and metabolic biomarker profiles consistent with increased insulin sensitivity. [4,6]
- **Early weight loss:** Retatrutide produces substantial early weight loss that continues with ongoing treatment. [2,29] Faster early weight loss than semaglutide or tirzepatide remains speculative; community descriptions are subjective comparisons.
- **Physical function and quality of life:** Retatrutide improves knee-osteoarthritis pain and physical function in adults with obesity. [9,29] Broader improvements in weight-related mobility, physical function, and quality of life are likely, with indirect support from tirzepatide quality-of-life findings. [23]
- **Obstructive sleep apnea:** Retatrutide reduced apnea–hypopnea events in two randomized 80-week trials enrolling adults with overweight or obesity and moderate-to-severe obstructive sleep apnea, with or without type 2 diabetes. At 12 mg, hybrid-estimand changes were −31.7 versus −9.9 events/hour without diabetes and −25.4 versus −10.3 with diabetes; intention-to-treat analyses were also favorable. [30] This is direct retatrutide evidence, extending earlier findings with tirzepatide. [19]
- **Liver-fat reduction:** Retatrutide produces large, dose-related reductions in liver fat. In a randomized MRI substudy, mean relative reduction at 24 weeks ranged from 42.9% with 1 mg to 82.4% with 12 mg. Participants had at least 10% liver fat and did not have type 2 diabetes. [4]
- **Metabolic liver disease progression:** Retatrutide plausibly improves MASH and slows progression of metabolic liver disease through reductions in liver fat, body weight, and insulin resistance. [4,6] Histologic MASH improvement with tirzepatide provides indirect support; the retatrutide substudy measured MRI liver fat rather than biopsy-defined fibrosis regression. [4,22]
- **Cardiometabolic markers:** Retatrutide reduces systolic blood pressure and atherogenic lipid measures. [2,7] Exploratory analyses found reductions in apolipoprotein B and selected lipoprotein particles; significant high-sensitivity C-reactive protein and interleukin-6 reductions occurred in participants with obesity without diabetes. [7]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- **Cardiovascular risk reduction:** Improvements in weight, glucose, blood pressure, and lipoproteins support a plausible cardiovascular-benefit hypothesis. [2,7,14] However, sponsor-reported TRIUMPH-3 event estimates were inconclusive: MACE-5 hazard ratio 0.82 (95% CI 0.55–1.22) and MACE-3 hazard ratio 1.12 (0.64–1.96). These intervals establish neither benefit nor harm. [10] TRIUMPH-Outcomes is evaluating clinical cardiovascular and kidney events. [11] Tirzepatide's noninferiority to dulaglutide is indirect class context, not proof of retatrutide superiority. [20]
- **Kidney protection:** Retatrutide is likely to reduce long-term kidney risk through improvements in glycemic control, blood pressure, and body weight, supported by exploratory albuminuria findings. [14,15] Most Phase 2 participants had normal baseline albumin excretion, so absolute changes were small; eGFR increased in the obesity trial but not the diabetes trial. [15] Tirzepatide's exploratory kidney-event benefit offers indirect support for this inference. [21]
- **Microvascular protection:** Sustained improvements in glycemic control and metabolic risk factors are likely to reduce long-term microvascular risk. [3,14] Tirzepatide was associated with fewer retinal complications in a retrospective cohort, providing indirect rather than causal retatrutide evidence. [27] This longer-term inference coexists with the risk of transient retinopathy worsening during rapid glucose improvement. [28]

Studied Applications

- **Obesity and overweight:** Weight reduction in adults with obesity or overweight and weight-related health conditions. [2,29]
- **Type 2 diabetes:** Glycemic control and weight reduction in adults with type 2 diabetes and overweight or obesity. [3,14,31]
- **Metabolic dysfunction-associated steatotic liver disease:** Reduction of MRI-measured liver fat and associated metabolic abnormalities. [4]
- **Obesity-related conditions:** Reduced knee-osteoarthritis pain and improved physical function, and reduced apnea–hypopnea events in obstructive sleep apnea. [9,29,30]
- **Cardiovascular and kidney outcomes:** Improvements in cardiovascular risk markers and exploratory kidney parameters; dedicated clinical-outcome evaluation is ongoing. [7,11,15]

Potential Applications

- **Alcohol craving and intake:** Reduced alcohol craving and consumption with retatrutide are plausible through shared GLP-1 effects on reward-related behavior. Self-reported reductions with semaglutide and tirzepatide provide indirect observational support. [24]
- **Nicotine craving:** Reduced nicotine craving with retatrutide is plausible through shared GLP-1 receptor activity. A small semaglutide trial reduced craving, while its primary laboratory smoking outcomes were not significantly different. [25]
- **Neuroprotective and cognitive effects:** Neuroprotective effects from retatrutide are plausible through shared incretin biology, while clinically meaningful cognitive benefits remain speculative. In high-fat-diet mice, tirzepatide reduced microglial activation, oxidative

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

stress, and NLRP3-associated inflammation through SIRT3 signaling , findings likely to translate for retatrutide. [26]

Cautions

- **Medullary thyroid carcinoma and MEN2:** Avoid retatrutide in people with a personal or family history of medullary thyroid carcinoma or Multiple Endocrine Neoplasia syndrome type 2, consistent with trial exclusions and incretin precautions. [17,28] Rodent C-cell findings with related drugs do not establish a causal human thyroid-cancer risk. [28]
- **Pancreatic and gallbladder disease:** A history of pancreatitis or gallbladder disease warrants caution. Pancreatitis and gallbladder disease are relevant incretin precautions; cholecystitis also occurred in the retatrutide liver-fat substudy. [4,28]
- **Severe gastrointestinal motility disorders:** Severe gastroparesis or other major motility disorders warrant caution. Retatrutide delays gastric emptying, most strongly after initial exposure, with attenuation during repeated dosing. [16] Gastrointestinal intolerance is common in its trials. [2,31]
- **Volume depletion and kidney injury:** Persistent vomiting or diarrhea can lead to dehydration and kidney injury. Retatrutide studies recorded dehydration, hypotension, and isolated acute kidney injury. Hydration and renal assessment matter particularly during escalation or persistent gastrointestinal symptoms. [15,17]
- **Insulin and sulfonylureas:** Combining potent glucose lowering with insulin or sulfonylureas can increase hypoglycemia risk; medication adjustment requires qualified clinical guidance. This precaution is extrapolated from related incretin therapy. [28] The absence of severe hypoglycemia in diet-and-exercise-only TRANSCEND-T2D-1 does not resolve combination-treatment risk. [14]
- **Diabetic retinopathy:** Rapid glucose improvement can temporarily worsen retinopathy. Existing disease or new visual symptoms warrant monitoring, even when longer-term metabolic control is expected to help. [28]
- **Oral hormonal contraceptives:** Reduced contraceptive absorption and effectiveness with retatrutide are plausible through delayed gastric emptying. [16] Nonoral or additional barrier contraception is a precaution extrapolated from tirzepatide; its four-week rule is not a validated retatrutide-specific interval. [28]
- **Anesthesia and deep sedation:** Delayed gastric emptying plausibly increases retained-stomach-content and aspiration risk. [16,28] Inform the procedural team about retatrutide use so management can be individualized.

Side Effects

- **Gastrointestinal effects:** Nausea, diarrhea, constipation, vomiting, and reduced appetite are documented in retatrutide trials. [2,9,31] Dyspepsia and other upper-abdominal

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

symptoms can also occur. [2] Lower initial exposure improved tolerability in Phase 2, and Phase 3 protocols use four-week escalation intervals with provision for dose reduction. [2,17]

- **Fatigue and hypotension:** Fatigue was reported in the Phase 2 obesity trial. [2] In TRIUMPH-2, hypotension occurred in approximately 1%, 5%, and 6% at 4, 9, and 12 mg versus less than 1% with placebo. [31]
- **Hair shedding:** Weight-loss-associated hair shedding is plausible during substantial weight reduction with retatrutide. Hair loss associated with weight reduction is reported for tirzepatide, providing indirect support; telogen effluvium is a possible explanation. [28]
- **Dysesthesia:** Altered skin sensations have occurred in Phase 3 trials. In TRIUMPH-4, rates were 8.8% at 9 mg and 20.9% at 12 mg versus 0.7% with placebo; most events were mild. [9] In TRIUMPH-2, rates were approximately 4%–7% across retatrutide doses versus 1% with placebo. [31]
- **Resting heart rate:** Dose-dependent increases occurred in Phase 2, peaked around week 24, and subsequently declined. [2] This pattern should be assessed separately from clinical cardiovascular-event outcomes. [10]

Pregnancy and Breastfeeding

Retatrutide should be avoided during pregnancy, while attempting to conceive, and while breastfeeding. Trials excluded pregnancy and breastfeeding, and protocols require discontinuation if pregnancy occurs. [17] Reproductive-risk assessment relies on indirect evidence from related incretin drugs. [12,13]

Observational evidence involving other GLP-1-based drugs provides limited reassurance about inadvertent exposure around conception or early pregnancy. It does not establish safe continued retatrutide exposure. [12,13] Low milk transfer after a single tirzepatide dose is likewise not a retatrutide lactation-safety result. [28]

A retatrutide-specific preconception washout interval has not been established. Pregnancy planning should include discontinuation sufficiently in advance under qualified clinical guidance, accounting for its long-acting PK. [1,12,13]

Retatrutide vs Tirzepatide

Tirzepatide activates GIP and GLP-1 receptors. Retatrutide additionally activates the glucagon receptor, creating a distinct metabolic profile. [1,8]

Retatrutide likely produces greater glucagon-driven stimulation of fatty-acid oxidation and energy expenditure than tirzepatide, a mechanistic inference supported by receptor biology, animal experiments, and human metabolomic associations. [1,6] Tirzepatide reduced appetite and increased fat oxidation in a human study without altering measured metabolic adaptation. [8]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Retatrutide

SC once weekly: 2 mg, titrate to 4 / 6 / 9 / 12 mg

Mechanistic hypothesis: Retatrutide is likely to produce greater human weight loss, visceral-fat reduction, liver-fat reduction, and endurance benefits than tirzepatide through its additional glucagon-receptor activity. [1,4,6,8] Comparative cardiovascular benefit remains unresolved given the inconclusive TRIUMPH-3 event estimates. [10,20] Community descriptions of faster fat loss, easier fasting, and a different subjective energy profile are speculative observations.

Retatrutide to Support Body Recomposition

Retatrutide produces large reductions in total fat mass. In the Phase 2 DXA substudy, lean mass also fell, with a proportion similar to that seen with other weight-loss treatments; lean mass is not identical to skeletal muscle. [5] Glucagon-receptor activity and increased fatty-acid oxidation provide a plausible mechanism for preferential fat loss, while a specific muscle-sparing effect remains speculative. [1,5,6] Adequate nutrition and resistance training are reasonable supporting measures: in an independent weight-loss trial in older adults, resistance-containing exercise preserved more lean mass than aerobic training alone. [32]

Biological Mechanism

Retatrutide activates GLP-1, GIP, and glucagon receptors, coordinating effects on appetite regulation, glucose-dependent insulin secretion, and lipid metabolism. [1] In animal experiments, glucagon-receptor signaling contributed to increased energy expenditure in addition to incretin-associated intake reduction. [1] Human metabolomic findings are consistent with increased fatty-acid oxidation and improved insulin resistance, supporting likely translation of this mechanism. [6] Gastric-emptying delay is greatest early in treatment and attenuates with repeated exposure, so persistent weight loss need not imply a sustained maximal gastric-emptying effect. [16] Together, these pathways provide a mechanistic basis for the observed weight and glycemic outcomes. [2,14,29,31]

References

[1] Coskun T, Urva S, Roell WC, et al. LY3437943, a novel triple glucagon, GIP, and GLP-1 receptor agonist for glycemic control and weight loss: From discovery to clinical proof of concept. *Cell Metab*. 2022;34(9):1234-1247.e9. doi:10.1016/j.cmet.2022.07.013

[2] Jastreboff AM, Kaplan LM, Frías JP, et al. Triple-Hormone-Receptor Agonist Retatrutide for Obesity - A Phase 2 Trial. *N Engl J Med*. 2023;389(6):514-526. doi:10.1056/nejmoa2301972

[3] Rosenstock J, Frias J, Jastreboff AM, et al. Retatrutide, a GIP, GLP-1 and glucagon receptor agonist, for people with type 2 diabetes: a randomised, double-blind, placebo and active-controlled, parallel-group, phase 2 trial conducted in the USA. *Lancet*. 2023;402(10401):529-544. doi:10.1016/s0140-6736(23)01053-x

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[4] Sanyal AJ, Kaplan LM, Frias JP, et al. Triple hormone receptor agonist retatrutide for metabolic dysfunction-associated steatotic liver disease: a randomized phase 2a trial. *Nat Med*. 2024;30(7):2037-2048. doi:10.1038/s41591-024-03018-2

[5] Coskun T, Wu Q, Schloot NC, et al. Effects of retatrutide on body composition in people with type 2 diabetes: a substudy of a phase 2, double-blind, parallel-group, placebo-controlled, randomised trial. *Lancet Diabetes Endocrinol*. 2025;13(8):674-684. doi:10.1016/s2213-8587(25)00092-0

[6] Pearson MJ, Willency JA, Lin Y, et al. Retatrutide and lipid and metabolite profiles in participants with obesity with or without type 2 diabetes. *J Clin Endocrinol Metab*. 2026;111(11):3217-3228. doi:10.1210/clinem/dgag201

[7] Ruotolo G, Harris C, Lin Y, et al. Retatrutide-Associated Improvements in Cardiovascular Risk Biomarkers in Adults With Obesity With or Without Type 2 Diabetes. *Diabetes Obes Metab*. 2026;28(11):10183-10194. doi:10.1111/dom.71200

[8] Ravussin E, Sanchez-Delgado G, Martin CK, et al. Tirzepatide did not impact metabolic adaptation in people with obesity, but increased fat oxidation. *Cell Metab*. 2025;37(5):1060-1074.e4. doi:10.1016/j.cmet.2025.03.011

[9] Eli Lilly and Company. Lilly's triple agonist, retatrutide, delivered weight loss of up to an average of 71.2 lbs along with substantial relief from osteoarthritis pain in first successful Phase 3 trial. Published December 11, 2025. Accessed October 25, 2026. https://investor.lilly.com/news-releases/news-release-details/lillys-triple-agonist-retatrutide-delivered-weight-loss-average

[10] Eli Lilly and Company. Lilly's triple agonist, retatrutide, successful in two additional Phase 3 obesity trials, delivering significant improvements in weight and A1C. Published July 23, 2026. Accessed October 25, 2026. https://lilly.gcs-web.com/news-releases/news-release-details/lillys-triple-agonist-retatrutide-successful-two-additional

[11] ClinicalTrials.gov. The Effect of Retatrutide Once Weekly on Cardiovascular Outcomes and Kidney Outcomes in Adults Living With Obesity (TRIUMPH-Outcomes). NCT06383390. Updated October 8, 2026. Accessed October 25, 2026. https://clinicaltrials.gov/study/NCT06383390

[12] Ozbek L, Shah E, Al-Shiab R, et al. Safety of GLP-1 and Dual GLP-1/GIP Receptor Agonists in Preconception, Pregnancy, and Lactation: A Systematic Review of Maternal, Fetal, and Neonatal Outcomes. *Diabetes Obes Metab*. 2026;28(6):4503-4528. doi:10.1111/dom.70699

[13] Maslin K, Shawe J, Blowers S, et al. Incretin-Based Medications in Women and Reproduction: A Systematic Scoping Review and Consensus Guidelines for Clinical Practice. *Obes Rev*. Published online July 29, 2026. e70203. doi:10.1111/obr.70203

[14] Bajaj HS, Welch M, Shah P, et al. Efficacy and safety of retatrutide, a GIP, GLP-1, and glucagon receptor agonist, in people with type 2 diabetes and inadequate glycaemic control with diet and exercise (TRANSCEND-T2D-1): a double-blind, randomised, phase 3 trial. *Lancet*. 2026;407(10546):2402-2413. doi:10.1016/s0140-6736(26)00967-0

[15] Heerspink HJL, Lu Z, Du Y, et al. The Effect of Retatrutide on Kidney Parameters in Participants With Type 2 Diabetes Mellitus and/or Obesity. *Kidney Int Rep*. 2025;10(6):1980-1992. doi:10.1016/j.ekir.2025.03.049

[16] Urva S, O'Farrell L, Du Y, et al. The novel GIP, GLP-1 and glucagon receptor agonist retatrutide delays gastric emptying. *Diabetes Obes Metab*. 2023;25(9):2784-2788. doi:10.1111/dom.15167

[17] Giblin K, Kaplan LM, Somers VK, et al. Retatrutide for the treatment of obesity, obstructive sleep apnea and knee osteoarthritis: Rationale and design of the TRIUMPH registrational clinical trials. *Diabetes Obes Metab*. 2026;28(1):83-93. doi:10.1111/dom.70209

[18] Jastreboff AM, le Roux CW, Stefanski A, et al. Tirzepatide for Obesity Treatment and Diabetes Prevention. *N Engl J Med*. 2025;392(10):958-971. doi:10.1056/nejmoa2410819

[19] Malhotra A, Grunstein RR, Fietze I, et al. Tirzepatide for the Treatment of Obstructive Sleep Apnea and Obesity. *N Engl J Med*. 2024;391(13):1193-1205. doi:10.1056/nejmoa2404881

[20] Nicholls SJ, Pavo I, Bhatt DL, et al. Cardiovascular Outcomes with Tirzepatide versus Dulaglutide in Type 2 Diabetes. *N Engl J Med*. 2025;393(24):2409-2420. doi:10.1056/nejmoa2505928

[21] Zoungas S, D'Alessio D, Pavo I, et al. A comparison of the effects of tirzepatide and dulaglutide on major kidney events in people with type 2 diabetes: pre-specified exploratory analyses of the SURPASS-CVOT trial. *Lancet Diabetes Endocrinol*. 2026;14(7):544-557. doi:10.1016/s2213-8587(26)00032-x

[22] Loomba R, Hartman ML, Lawitz EJ, et al. Tirzepatide for Metabolic Dysfunction-Associated Steatohepatitis with Liver Fibrosis. *N Engl J Med*. 2024;391(4):299-310. doi:10.1056/nejmoa2401943

[23] Gudzune KA, Stefanski A, Cao D, et al. Association between weight reduction achieved with tirzepatide and quality of life in adults with obesity: Results from the SURMOUNT-1 study. *Diabetes Obes Metab*. 2025;27(2):539-550. doi:10.1111/dom.16046

[24] Quddos F, Hubshman Z, Tegge A, et al. Semaglutide and Tirzepatide reduce alcohol consumption in individuals with obesity. *Sci Rep*. 2023;13(1):20998. doi:10.1038/s41598-023-48267-2

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[25] Hendershot CS, Bremmer MP, Paladino MB, et al. Once-Weekly Semaglutide in Adults With Daily Cigarette Use: A Randomized Clinical Trial. *JAMA Netw Open*. 2026;9(5):e2614898. doi:10.1001/jamanetworkopen.2026.14898

[26] Ma J, Liu Y, Hu J, et al. Tirzepatide administration improves cognitive impairment in HFD mice by regulating the SIRT3-NLRP3 axis. *Endocrine*. 2025;87(2):486-497. doi:10.1007/s12020-024-04013-w

[27] Shah J, Razavi P, Festok M, et al. Tirzepatide and Reduced Risk of Diabetic Retinopathy and Related Complications: A Multicenter US Cohort Study. *Ophthalmology*. 2026;133(6):728-732. doi:10.1016/j.ophtha.2026.01.013

[28] Eli Lilly and Company. Zepbound (tirzepatide) injection: prescribing information. Revised August 2026. US Food and Drug Administration. Accessed October 25, 2026. https://www.accessdata.fda.gov/drugsatfda_docs/label/2026/217806s041lbl.pdf

[29] Jastreboff AM, Kaplan LM, Davies MJ, et al. Retatrutide, a Triple Hormone Receptor Agonist, for Treatment of Obesity. *N Engl J Med*. Published online September 29, 2026. doi:10.1056/nejmoa2604169

[30] Grunstein RR, Somers VK, Azarbarzin A, et al. Retatrutide Once Weekly for Treatment of Obstructive Sleep Apnea. *N Engl J Med*. Published online October 21, 2026. doi:10.1056/nejmoa2611477

[31] Bellido V, le Roux CW, Ekinci EI, et al. Retatrutide in adults with obesity and type 2 diabetes (TRIUMPH-2): a double-blind, parallel-group, randomised, placebo-controlled, phase 3 trial. *Lancet*. 2026;408(10563):1455-1472. doi:10.1016/s0140-6736(26)01861-1

[32] Villareal DT, Aguirre L, Gurney AB, et al. Aerobic or Resistance Exercise, or Both, in Dieting Obese Older Adults. *N Engl J Med*. 2017;376(20):1943–1955. doi:10.1056/NEJMoa1616338