Tesamorelin + Ipamorelin For Research Use Only

Speculative: SC Tesa 0.5–2 mg + Ipa 100–250 mcg, 3–7 days/week

Tesamorelin activates the growth hormone-releasing hormone (GHRH) receptor, whereas ipamorelin activates the ghrelin receptor system. Visceral fat reduction, lower liver fat, improved body composition, and selected cognitive outcomes are supported by tesamorelin trials in specific populations. [1–8,12,17] Recovery and fitness benefits from adding ipamorelin are plausible translational applications. Increased endogenous growth hormone (GH) output and downstream insulin-like growth factor 1 (IGF-1) are therefore likely with combined stimulation. [10,18,20,28] A stronger or smoother response and improved tolerability remain speculative combination-specific effects.

💉 Tesamorelin safety is characterized through 52 weeks in selected adults, with recognized glucose, fluid-retention, and immune risks. Acceptable short-term tolerability of the combination is plausible by extrapolation; serious and delayed outcomes require separate assessment. [3–5,19,25–27]

⭐ ⭐ ⭐ Combined GH stimulation is likely by mechanistic and class-based inference; additional recovery, fitness, and body-recomposition benefits are plausible. [10,18,21,22,28]

⭐ ⭐ ⭐ Visceral fat reduction is established for tesamorelin. A corresponding benefit from the combination is plausible, depending on effective tesamorelin exposure. [2–5]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Dosage & Patterns

The pivotal phase 3 trials used the historical tesamorelin formulation at 2 mg subcutaneously once daily for 26 weeks, with extensions to 52 weeks. [2–5]

The following speculative experimental regimens are retained from community proposals, rather than validated combination doses or conversions from the approved formulations:

- **Tesamorelin 2 mg with ipamorelin 250 mcg subcutaneously**, speculative experimental regimen.
- **Tesamorelin 0.5–1 mg with ipamorelin 100–250 mcg subcutaneously**, speculative experimental lower-dose approach intended to improve tolerability.
- **Daily, five days per week, or every other day** are speculative community schedules. A **fasted schedule** is also proposed; it is not a requirement in the approved tesamorelin dosing instructions. [25,26]
- **12 weeks ON / 6 weeks OFF, or 12–24 weeks ON / 4–12 weeks OFF**, are common community proposals. These schedules remain speculative. Tesamorelin trials generally used continuous daily treatment through 26 or 52 weeks. [3–5] Clinical response, IGF-1,

glucose control, and adverse effects provide a more defensible basis for reassessing exposure than a fixed calendar cycle. [25,26]

Benefits

The following findings concern tesamorelin alone unless otherwise stated. Its human evidence includes VAT, liver fat, lipids, lean mass, muscle composition, and selected cognitive outcomes. [1–8,12,17] Ipamorelin may augment the secretory response through complementary receptor stimulation; extension to clinical benefits is a translational inference. [18,20,28]

- **Visceral fat reduction**: Across two phase 3 trials involving 806 adults with HIV-associated abdominal fat accumulation, tesamorelin produced a placebo-adjusted 15.4% VAT reduction at 26 weeks. Abdominal subcutaneous fat was largely preserved; continued treatment maintained VAT and waist improvements through 52 weeks. [5]
- **Improved lipid profile**: The pooled phase 3 analysis found lower triglycerides and a lower total-cholesterol-to-HDL ratio at 26 weeks. These are metabolic outcomes rather than demonstrated cardiovascular-event prevention. [5]
- **Reduced liver fat**: In a 12-month trial of HIV-associated fatty liver disease, 61 participants were enrolled and 60 received treatment. Tesamorelin reduced hepatic fat fraction by 4.1 percentage points relative to placebo, corresponding to a 37% relative reduction. [7] A secondary biopsy signal suggested less fibrosis progression. [7,15]
- **Improved executive function**: A 20-week trial randomized 152 older adults, including people with mild cognitive impairment (MCI), to tesamorelin 1 mg daily or placebo. Overall cognition and executive function improved; verbal memory showed a trend (P=.08), with no significant treatment-by-diagnosis interaction. The findings do not establish an MCI-specific delayed-recall benefit. [17]
- **Increased lean body mass**: A 12-week dose-ranging study in adults with HIV-associated abdominal fat accumulation found increased lean mass with tesamorelin 1 and 2 mg daily relative to placebo. [1] An earlier trial of native GHRH also increased lean mass, but it was a different intervention. [9]
- **Improved muscle composition**: A responder-selected analysis compared 193 tesamorelin recipients with 148 placebo recipients and found improvements in trunk-muscle density and area over 26 weeks. These imaging findings concern selected VAT responders rather than demonstrated gains in strength or physical function. [12] Related analyses found increased fat density in responders and modest changes in inflammatory or fibrinolytic markers associated with VAT reduction. [13,14]

Potential Applications

- **Excess visceral fat**: [8,16]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- **Fatty liver**: [6,7,15]
- **Cognition, recovery, fitness, and body recomposition**: Additional benefit from complementary GH-axis stimulation is plausible. [10,17,18, 21,22,28]

Cautions

- **Malignancy**: Active malignancy contraindicates tesamorelin. A previous malignancy should be inactive and treatment complete; recurrence requires discontinuation. [25,26]
- **Pregnancy and lactation**: Pregnancy is a contraindication. Human milk exposure and infant effects are inadequately characterized; the labeled HIV population is advised not to breastfeed. Reproductive safety of the combination is not established. [25,26]
- **Glucose regulation**: Assess glucose before and during exposure. In the tesamorelin registration trials, HbA1c of at least 6.5% developed in 5% of treated participants versus 1% with placebo by week 26. Patients with diabetes require monitoring, including for retinopathy. [25] Neutral mean glucose results in smaller trials do not eliminate this individual risk. [7,11]
- **Elevated IGF-1**: Monitor age-adjusted IGF-1 and consider discontinuation for persistent values above +3 SDS, especially when clinical benefit is limited. [25,26]
- Other contraindications and cautions: Disruption of the hypothalamic-pituitary axis is a contraindication. Unexplained masses warrant assessment. Acute critical illness requires particular caution based on GH-class experience. Glucocorticoid replacement and drugs affected by GH-related changes in metabolism may require reassessment. [25,26]

Side Effects

- **Fluid retention**: Peripheral edema can occur; the fluid burden may be important in cardiovascular disease. Long-term cardiovascular-event safety remains unresolved. [17,25]
- **Musculoskeletal and peripheral nerve symptoms**: Arthralgia, myalgia, and carpal-tunnel symptoms are recognized effects of tesamorelin. [25,26]
- Sleep: Deeper sleep and vivid dreams are anecdotal, speculative effects. In the tesamorelin cognition trial, measured sleep quality did not improve relative to placebo. [17]
- Other observed effects: Headache, nausea, and injection-site reactions have been reported. [19,25]

Cancer Risk

Tesamorelin raises IGF-1, sometimes above age-adjusted reference ranges. [25] A large observational study associated higher IGF-1 with greater overall cancer incidence but divergent associations across cancer sites; it does not establish drug causation. [24] Promotion of an occult or recurrent malignancy remains a biologically plausible risk of sustained GH/IGF-1 stimulation. Short trials cannot resolve long-latency carcinogenic risk.

Drivers of Diminished Response and Mitigation Strategies

Negative feedback and receptor or intracellular adaptation are plausible contributors to variable response; their clinical importance depends on exposure and biological context. [10,29,30] In

vitro neutralizing antibodies were detected during tesamorelin treatment, but comparable mean VAT and IGF-1 responses argue against treating antibody-mediated loss as a demonstrated common outcome. [25] Sustained efficacy through 52 weeks also conflicts with a blanket prediction of a high likelihood of clinically meaningful waning. [3–5]

Controlled tesamorelin trials maintained visceral adipose tissue (VAT) reductions during continuous treatment for up to 52 weeks. [3–5] Anti-tesamorelin antibodies were common, but mean VAT and IGF-1 responses were comparable in participants with and without antibodies, including those with in vitro neutralizing activity. [25] These findings argue against routinely assuming that treatment breaks are needed to preserve efficacy.

Periodic OFF periods during indefinite or repeated use remain a conservative, mechanistically plausible way to reduce cumulative GH/IGF-1 exposure. That rationale must be weighed against the observed return of visceral fat after tesamorelin withdrawal. [3,4]

Monitoring

Persistent IGF-1 elevation above +3 standard deviation scores (SDS), worsening glucose control, or clinically important edema, arthralgia, or carpal-tunnel symptoms warrant reassessment of GH-axis stimulation. [25,26] Reconsidering exposure after laboratory abnormalities and symptoms have resolved is a conservative clinical inference, not an established restart protocol for the combination.

Biological Mechanism

Tesamorelin activates pituitary GHRH receptors, whereas ipamorelin acts through the GH-secretagogue/ghrelin receptor system. [18,20,25] Human challenges with native GHRH and GHRP-2 produced a larger combined GH response than either alone, supporting a class-based rationale for complementary stimulation, although that study did not demonstrate superadditive synergy. [28] Increased output with tesamorelin plus ipamorelin is likely by inference. Retention of endogenous feedback is plausible, but does not guarantee normal IGF-1 exposure. [10,25]

Receptor Binding and Intracellular Signaling

GHRH-receptor signaling primarily engages Gs, adenylyl cyclase, cAMP, and protein kinase A. Ghrelin-receptor signaling involves Gq/11, phospholipase C, inositol phosphates, and calcium mobilization. The pathways overlap: in porcine somatotropes, ghrelin-dependent GH release required both PLC/PKC and cAMP/PKA signaling and extracellular calcium influx. Combined ghrelin/GHRH exposure increased cAMP additively without increasing GH secretion beyond either stimulus in that preparation. Thus, complementary signaling supports a plausible additive effect rather than obligatory synergy in every system. [29]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Pituitary Dynamics and GH Pulsatility

Tesamorelin increased basal secretion and GH pulse area after two weeks in healthy men. Sampling began about eight hours after the final injection, supporting the interpretation that tesamorelin probably enhances multiple successive endogenous pulses through biological effects that outlast its short plasma half-life. Whether this reflects each administration, adaptations to repeated dosing, or both remains unresolved. [10] Intravenous ipamorelin generated a discrete GH response. [18] A larger combined secretory response is likely by class-based inference; a longer or smoother pulse pattern is speculative. [28]

Peripheral Growth-Hormone and IGF-1 Effects

GH-receptor signaling increases hepatic IGF-1 production and engages JAK2 and STAT pathways, with cell-context-dependent signaling through Shc and downstream mitogenic pathways. [25,30] Increased lipolysis, lipid oxidation, and altered very-low-density-lipoprotein metabolism are plausible contributors to the observed fat and lipid changes. [5,15] Tesamorelin-associated liver-biopsy changes included increased oxidative-phosphorylation gene signatures. [15] IGF-1 stimulated anabolic protein metabolism in cultured human muscle cells and mitochondrial biogenesis and turnover during experimental myoblast differentiation. These findings support plausible protein-synthesis and mitochondrial mechanisms for the combination. [31,32]

Conclusions

Tesamorelin has clinically supported effects on VAT and selected metabolic endpoints in the studied populations; cognition and muscle-imaging findings require their narrower study context. [1–8,12,17] Adding ipamorelin is likely to increase GH stimulation through complementary receptor activity, while additional recovery, fitness, and body-composition benefits remain plausible translational inferences. [18,20–22,28] A smoother response, better tolerability, and benefits from scheduled cycling remain speculative. Safety judgments should remain endpoint-specific. [24–27]

References

Tesamorelin clinical trials and human studies

[1] Falutz J, Allas S, Kotler D, et al. A placebo-controlled, dose-ranging study of a growth hormone releasing factor in HIV-infected patients with abdominal fat accumulation. *AIDS*. 2005;19(12):1279-1287. doi:10.1097/01.aids.0000180099.35146.30

[2] Falutz J, Allas S, Blot K, et al. Metabolic effects of a growth hormone-releasing factor in patients with HIV. *N Engl J Med*. 2007;357(23):2359-2370. doi:10.1056/nejmoa072375

[3] Falutz J, Allas S, Mamputu JC, et al. Long-term safety and effects of tesamorelin, a growth hormone-releasing factor analogue, in HIV patients with abdominal fat accumulation. *AIDS*. 2008;22(14):1719-1728. doi:10.1097/qad.0b013e32830a5058

[4] Falutz J, Potvin D, Mamputu JC, et al. Effects of tesamorelin, a growth hormone-releasing factor, in HIV-infected patients with abdominal fat accumulation: a randomized placebo-controlled trial with a safety extension. *J Acquir Immune Defic Syndr.* 2010;53(3):311-322. doi:10.1097/qai.0b013e3181cbdaff

[5] Falutz J, Mamputu JC, Potvin D, et al. Effects of tesamorelin (TH9507), a growth hormone-releasing factor analog, in human immunodeficiency virus-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind placebo-controlled phase 3 trials with safety extension data. *J Clin Endocrinol Metab.* 2010;95(9):4291-4304. doi:10.1210/jc.2010-0490

[6] Stanley TL, Feldpausch MN, Oh J, et al. Effect of tesamorelin on visceral fat and liver fat in HIV-infected patients with abdominal fat accumulation: a randomized clinical trial. *JAMA*. 2014;312(4):380-389. doi:10.1001/jama.2014.8334

[7] Stanley TL, Fourman LT, Feldpausch MN, et al. Effects of tesamorelin on non-alcoholic fatty liver disease in HIV: a randomised, double-blind, multicentre trial. *Lancet HIV.* 2019;6(12):e821-e830. doi:10.1016/s2352-3018(19)30338-8

[8] Makimura H, Feldpausch MN, Rope AM, et al. Metabolic effects of a growth hormone-releasing factor in obese subjects with reduced growth hormone secretion: a randomized controlled trial. *J Clin Endocrinol Metab.* 2012;97(12):4769-4779. doi:10.1210/jc.2012-2794

[9] Koutkia P, Canavan B, Breu J, Torriani M, Kissko J, Grinspoon S. Growth hormone-releasing hormone in HIV-infected men with lipodystrophy: a randomized controlled trial. *JAMA*. 2004;292(2):210-218. doi:10.1001/jama.292.2.210

[10] Stanley TL, Chen CY, Branch KL, Makimura H, Grinspoon SK. Effects of a growth hormone-releasing hormone analog on endogenous GH pulsatility and insulin sensitivity in healthy men. *J Clin Endocrinol Metab.* 2011;96(1):150-158. doi:10.1210/jc.2010-1587

[11] Clemmons DR, Miller S, Mamputu JC. Safety and metabolic effects of tesamorelin, a growth hormone-releasing factor analogue, in patients with type 2 diabetes: A randomized, placebo-controlled trial. *PLoS One.* 2017;12(6):e0179538. doi:10.1371/journal.pone.0179538

[12] Adrian S, Scherzinger A, Sanyal A, et al. The Growth Hormone Releasing Hormone Analogue, Tesamorelin, Decreases Muscle Fat and Increases Muscle Area in Adults with HIV. *J Frailty Aging.* 2019;8(3):154-159. doi:10.14283/jfa.2018.45

[13] Lake JE, La K, Erlandson KM, et al. Tesamorelin improves fat quality independent of changes in fat quantity. *AIDS.* 2021;35(9):1395-1402. doi:10.1097/qad.0000000000002897

[14] Stanley TL, Falutz J, Mamputu JC, Soulban G, Potvin D, Grinspoon SK. Effects of tesamorelin on inflammatory markers in HIV patients with excess abdominal fat: relationship with visceral adipose reduction. *AIDS.* 2011;25(10):1281-1288. doi:10.1097/qad.0b013e328347f3f1

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[15] Fourman LT, Billingsley JM, Agyapong G, et al. Effects of tesamorelin on hepatic transcriptomic signatures in HIV-associated NAFLD. *JCI Insight*. 2020;5(16):e140134. doi:10.1172/jci.insight.140134

[16] Russo SC, Ockene MW, Arpante AK, et al. Efficacy and safety of tesamorelin in people with HIV on integrase inhibitors. *AIDS*. 2024;38(12):1758-1764. doi:10.1097/qad.0000000000003965

[17] Baker LD, Barsness SM, Borson S, et al. Effects of growth hormone–releasing hormone on cognitive function in adults with mild cognitive impairment and healthy older adults: results of a controlled trial. *Arch Neurol*. 2012;69(11):1420-1429. doi:10.1001/archneurol.2012.1970

Ipamorelin human evidence

[18] Gobburu JV, Agersø H, Jusko WJ, Ynddal L. Pharmacokinetic-pharmacodynamic modeling of ipamorelin, a growth hormone releasing peptide, in human volunteers. *Pharm Res*. 1999;16(9):1412-1416. doi:10.1023/a:1018955126402

[19] Beck DE, Sweeney WB, McCarter MD; Ipamorelin 201 Study Group. Prospective, randomized, controlled, proof-of-concept study of the Ghrelin mimetic ipamorelin for the management of postoperative ileus in bowel resection patients. *Int J Colorectal Dis*. 2014;29(12):1527-1534. doi:10.1007/s00384-014-2030-8

Ipamorelin animal and mechanistic evidence

[20] Raun K, Hansen BS, Johansen NL, et al. Ipamorelin, the first selective growth hormone secretagogue. *Eur J Endocrinol*. 1998;139(5):552-561. doi:10.1530/eje.0.1390552

[21] Svensson J, Lall S, Dickson SL, et al. The GH secretagogues ipamorelin and GH-releasing peptide-6 increase bone mineral content in adult female rats. *J Endocrinol*. 2000;165(3):569-577. doi:10.1677/joe.0.1650569

[22] Andersen NB, Malmlöf K, Johansen PB, Andreassen TT, Ørtoft G, Oxlund H. The growth hormone secretagogue ipamorelin counteracts glucocorticoid-induced decrease in bone formation of adult rats. *Growth Horm IGF Res*. 2001;11(5):266-272. doi:10.1054/ghir.2001.0239

[23] Venkova K, Mann W, Nelson R, Greenwood-Van Meerveld B. Efficacy of ipamorelin, a novel ghrelin mimetic, in a rodent model of postoperative ileus. *J Pharmacol Exp Ther*. 2009;329(3):1110-1116. doi:10.1124/jpet.108.149211

Related IGF-1 epidemiology

[24] Qian F, Huo D. Circulating Insulin-Like Growth Factor-1 and Risk of Total and 19 Site-Specific Cancers: Cohort Study Analyses from the UK Biobank. *Cancer Epidemiol Biomarkers Prev*. 2020;29(11):2332-2342. doi:10.1158/1055-9965.epi-20-0743

Additional prescribing and mechanistic sources

[25] Theratechnologies Inc. EGRIFTA WR (tesamorelin) prescribing information. Revised March 2025. DailyMed. Accessed October 29, 2026. Source

[26] Theratechnologies Inc. EGRIFTA SV (tesamorelin) prescribing information. Revised March 2024. DailyMed. Accessed October 29, 2026. Source

[27] US Food and Drug Administration. Pharmacy Compounding Advisory Committee briefing document: ipamorelin acetate and ipamorelin free base. October 29, 2024. Accessed October 29, 2026. Source

[28] Tiulpakov AN, Brook CG, Pringle PJ, Peterkova VA, Volevodz NN, Bowers CY. GH responses to intravenous bolus infusions of GH releasing hormone and GH releasing peptide 2 separately and in combination in adult volunteers. *Clin Endocrinol (Oxf).* 1995;43(3):347-350. doi:10.1111/j.1365-2265.1995.tb02042.x

[29] Malagón MM, Luque RM, Ruiz-Guerrero E, et al. Intracellular signaling mechanisms mediating ghrelin-stimulated growth hormone release in somatotropes. *Endocrinology.* 2003;144(12):5372-5380. doi:10.1210/en.2003-0723

[30] Han Y, Leaman DW, Watling D, et al. Participation of JAK and STAT proteins in growth hormone-induced signaling. *J Biol Chem.* 1996;271(10):5947-5952. doi:10.1074/jbc.271.10.5947

[31] Guan X, Yan Q, Wang D, Du G, Zhou J. IGF-1 Signaling Regulates Mitochondrial Remodeling during Myogenic Differentiation. *Nutrients.* 2022;14(6):1249. doi:10.3390/nu14061249

[32] Crown AL, He XL, Holly JM, Lightman SL, Stewart CE. Characterisation of the IGF system in a primary adult human skeletal muscle cell model, and comparison of the effects of insulin and IGF-I on protein metabolism. *J Endocrinol.* 2000;167(3):403-415. doi:10.1677/joe.0.1670403