Prostamax (KEDP) Available for Research Use Only

(200 mcg–1 mg daily SC for 10–20 days, repeated once or twice annually)

Prostamax (KEDP) is an experimental synthetic tetrapeptide investigated within the Khavinson short-peptide bioregulator research program. [3] Prostamax has demonstrated anti-inflammatory, antifibrotic, and antihyperplastic effects in specific rat prostate models, with plausible translation to urinary symptom relief. [1–3] It is distinct from the animal-prostate extract Prostatilen. [3,8] Cultured human lymphocytes show changes in chromatin organization and ribosomal-gene activity. [4–6] Direct DNA-interaction experiments support a proposed gene-regulatory mechanism, while prostate-tissue culture findings suggest context-dependent reparative effects. [11,12]

💉 Plausibly safe; Plausible organ tolerability; human repeat-course and cancer safety unresolved. [3,6]

⭐ Speculative human symptom relief. Demonstrated selected prostate effects in rat models [1–3]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

The patent reports no significant blood-count or biochemical changes in its prostatitis patients and no pathological changes in general condition, selected laboratory measures, or examined organs during rat IM toxicity studies lasting 90 days and six months. [3] These observations support plausible organ tolerability over months, with low confidence in translation to humans because reporting is limited and the studies are patent examples.

The strongest direct efficacy evidence is preclinical. The two disease-model papers involve overlapping investigators. Their consistent direction supports a biological rationale, but differences in model, limited independent replication, and incomplete reporting keep human symptom benefit in the plausible category. [1,2]

How Evidence Scope Was Applied in This Review

Evidence was evaluated here with an adjustment for source quality. Much of the Prostamax evidence comes from a single regional source, including small or methodologically limited studies that have not been independently replicated. When a claim depended primarily on this evidence, its evidence classification was reduced to reflect the additional uncertainty while still recognizing the evidence as informative.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Prostamax (KEDP) Available for Research Use Only

(200 mcg–1 mg daily SC for 10–20 days, repeated once or twice annually)

Prostamax vs Prostatilen

Prostatilen is an animal-prostate extract containing a mixture of peptides. Russian product information lists chronic prostatitis and postoperative prostate conditions among its uses. **Prostamax** is the defined synthetic KEDP sequence; **equivalence to the extract or its complete biological activity has not been demonstrated.** [3,8]

Human-cell experiments on Prostamax are mechanistic studies. Patent clinical examples are preliminary observations. [3–6]

Russian registration and prescribing information for Prostatilen apply to that product. They do not establish clinical approval or a standard-of-care role for Prostamax. [8] A multicomponent extract and a single synthetic peptide may differ in targets, exposure, efficacy, and safety. Evidence cannot be transferred solely because both are described as prostate bioregulators. Prostatilen has product-specific IM and rectal formulations and dosing instructions. Those milligram doses and schedules cannot be converted into an equivalent Prostamax regimen. [8]

Dosage & Patterns

No validated human SC regimen was identified. Human SC PK remains uncharacterized.

The patent describes human IM administration spanning approximately 0.68 mcg–6.8 mg daily for a 150-lb adult, for 10–40 days. This broad reported exposure range is not a recommended regimen; dose-specific outcomes are not provided. [3] The 2014 hyperplasia study has an unresolved dose-unit concern, so its dose is excluded from human extrapolation. [2]

A rodent study for prostate inflammation used 220 mcg HED daily for 15 days.

The speculative, experimental community pattern discussed is

- **200 mcg–1 mg daily SC for 10–20 days, repeated once or twice annually**

Combining Prostamax with Epitalon or other bioregulator peptides is a speculative community practice.

Benefits

- **Reduced prostate inflammation** in animals. The aseptic prostatitis experiment found less inflammatory tissue change after treatment; the patent also describes improvement in a bacterial prostatitis model. These findings support plausible human anti-inflammatory benefit, rather than an inference of antibacterial activity. [1,3]
- **Plausible improvement in urinary symptoms and pelvic discomfort**. The patent describes 35 treated chronic-prostatitis patients and 19 treated prostate-adenoma patients, with separate comparison groups of 14 and 17. It reports improved symptoms and uroflow

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

measures. Randomization and blinding are not described, and outcome tables emphasize before–after changes; these are preliminary human observations. [3]

- **Reduced inflammation-associated fibrosis** in animals. Collagen accumulation fell in the sutured-prostate rat model. The authors also proposed epithelial protection, but the epithelial-area comparison with untreated prostatitis was not marked statistically significant in the reported table. [1]
- **Reduced experimentally induced prostate enlargement** in animals. In 40 rats assigned to four groups, KEDP reduced lateral-prostate mass by 24% and volume by 40% versus the sulpiride control. This induced model is distinct from naturally developing human BPH. [2]
- Chromatin modulation in cultured human cells. Experiments report altered chromatin condensation, thermal behavior, and ribosomal-gene activity. These are laboratory endpoints supporting a plausible regulatory mechanism. [4–6]

Applications

- Chronic prostatitis, particularly inflammation-associated tissue injury. [1,3] Clinical evaluation must distinguish bacterial disease and other causes of pelvic symptoms. [9]
- Benign prostatic hyperplasia and associated urinary symptoms. [2,3]
- Preventive prostate healthspan support, based on chromatin and tissue-repair hypotheses. [4–7,11,12]

Cautions

- **Known or suspected prostate cancer**: tumor promotion is a theoretical concern when altering cell regulation. The available chromatin findings cannot determine the direction of cancer risk. [4–7,11,12]
- **Unexplained or worsening urinary symptoms:** Evaluate the cause before attributing symptoms to benign enlargement or attempting experimental treatment. [9]
- **Urgent symptoms:** Inability to urinate, fever with painful urinary symptoms, blood in urine, or severe lower-abdominal pain require prompt medical care. [9]
- Concurrent medicines and hormones: document them during assessment so symptom changes can be interpreted. Prostamax-specific interactions with testosterone, DHT, and standard prostate medicines remain uncharacterized.

Diminished Response and Interpretation

Receptor downregulation, neutralizing antibodies, and compensatory adaptation remain uncharacterized for Prostamax. Speculatively, altered cellular state or downstream adaptation could change responsiveness during repeated exposure.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Biological Mechanism

The underlying molecular mechanisms of peptide bioregulators such as Prostamax remain poorly characterized. A proposed mechanism is interaction with DNA and chromatin that changes access to transcriptional machinery, rather than a defined receptor-agonist pathway. Prostamax altered DNA melting behavior in cell-free experiments, and cultured lymphocytes showed chromatin changes. The causal sequence connecting these observations to prostate effects remains incompletely characterized. [3–7,12]

The scientific literature more often describes peptide bioregulators through their observed downstream cellular and physiological effects rather than through well-established mechanisms explaining how those effects originate.

Prostamax altered chromatin organization in cultured human blood cells, including cells from older donors. Reported effects include decondensation of densely packed chromatin and altered ribosomal-gene activity. [4,6] Microcalorimetry found shifts in chromatin melting behavior interpreted as partial structural relaxation. [5] DNA melting experiments provide additional support for direct physicochemical interaction; docking of individual amino acids in that work does not establish sequence-specific KEDP targeting of particular genes. [12]

Prostate responses appear context-dependent. Organotypic cultures from young and aged rats showed growth stimulation with Prostamax, supporting a reparative hypothesis. [11] Reduced epithelial expansion in induced hyperplasia and the proposed preservation of epithelium during inflammatory injury describe different experimental contexts. Neither uniform growth stimulation nor uniform growth inhibition adequately describes these findings. [1,2]

Prostamax's antifibrotic rationale rests on reduced collagen accumulation in experimental prostatitis. [1] Plausibly, less inflammation and congestion could improve local tissue function, secretion, and bladder emptying. The symptom and uroflow observations in the patent are preliminary clinical support for the urinary endpoints. [3]

Chromatin findings in lymphocytes show that laboratory effects are not confined to prostate cells. Broader immune regulation or systemic healthspan effects are speculative extrapolations from these observations. [4–7]

Cell regulation can have different consequences in healthy, inflamed, hyperplastic, and malignant tissue. One cultured-lymphocyte study reported increased sister-chromatid exchange after Prostamax exposure. This is a DNA-exchange endpoint, not proof of beneficial DNA repair or of clinical carcinogenicity. It warrants attention in genomic-safety assessment. Current findings support neither established cancer protection nor established cancer promotion. [4–7]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Conclusions

Prostamax has preclinical support for selected prostate effects, preliminary patent-reported human observations, and a plausible chromatin-related mechanism. Animal tolerability findings support cautious inference about selected organ endpoints over months. Human SC dosing, repeated-course safety, and oncologic outcomes remain unresolved. Prostamax remains an investigational peptide with a biological rationale. [1–7,11,12]

References

[1] Borovskaya TG, Pakhomova AV, Vychuzhanina AV, et al. Experimental studying of the drug efficiency Prostamax in the therapy of chronic aseptic prostatitis and its complications. Modern Research in Inflammation. 2013;2(3):54–58. doi:10.4236/mri.2013.23007

[2] Borovskaya TG, Fomina TI, Shchemerova JA, et al. Experimental Study of Efficiency of Tertapeptide Lysil-Glutamyl-Aspartyl-Proline Using the Model of Benign Prostatic Hyperplasia. Modern Research in Inflammation. 2014;3:108–112. doi:10.4236/mri.2014.33013

[3] Khavinson VKh, Malinin VV, Grigoriev EI, inventors. Tetrapeptide regulating prostate functions and its compositions and uses. European patent EP1353939B1. Published August 25, 2004. Source

[4] Khavinson VKh, Lezhava TA, Malinin VV. Effects of short peptides on lymphocyte chromatin in senile subjects. Bull Exp Biol Med. 2004;137(1):78-81. doi:10.1023/b:bebm.0000024393.40560.05

[5] Meskhi T, Khachidze D, Barbakadze Sh, et al. The influence of the peptide bioregulator prostamax on heterochromatin of human lymphocytes in situ. Biofizika. 2004;49(6):1091-1093. Russian. Source

[6] Dzhokhadze TA, Buadze TZh, Gaĭozishvili MN, Baratashvili NA, Lezhava TA. Deheterochromatinization of the chromatin in old age induced by oligopeptide bioregulator (Lys-Glu-Asp-Pro). Georgian Med News. 2012;(212):76-82. Russian. Source

[7] Khavinson VK, Popovich IG, Linkova NS, Mironova ES, Ilina AR. Peptide Regulation of Gene Expression: A Systematic Review. Molecules. 2021;26(22):7053. doi:10.3390/molecules26227053

[8] Cytomed. Prostatilen: Russian prescribing information and patient leaflets. Accessed October 25, 2026. Source

[9] National Institute of Diabetes and Digestive and Kidney Diseases. Prostatitis: inflammation of the prostate. Accessed October 25, 2026. Source

[10] US Food and Drug Administration. Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers. Published July 2005. Source

[11] Zakutskiĭ AN, Chalisova NI, Ryzhak GA, Aniskina AI, Filippov SV, Zeziulin PN. The tissue-specific effect of synthetic peptides-biologic regulators in organotypic tissues culture in young and old rats. Adv Gerontol. 2006;19:93-96. Russian. Source

[12] Solovyev AY, Tarnovskaya SI, Chernova IA, Shataeva LK, Skorik YA. The interaction of amino acids, peptides, and proteins with DNA. Int J Biol Macromol. 2015;78:39-45. doi:10.1016/j.ijbiomac.2015.03.054

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076