Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

Metformin 500 mg orally daily; tadalafil 5 mg orally daily

Metformin and tadalafil have complementary metabolic and vascular actions that make combined experimental healthspan benefits plausible. [3–6,13,36] Metformin lowers glucose and influences intestinal metabolism, cellular energy sensing, and maintenance pathways. [3,4,40] Tadalafil inhibits phosphodiesterase type 5 (PDE5), sustaining cyclic guanosine monophosphate (cGMP) signaling downstream of nitric oxide (NO). [25] Endothelial improvements have been observed in selected human studies, and increased cerebral blood flow was observed with 20 mg daily in patients with cerebral small-vessel disease. [23,28–30] Metabolic, vascular, and neuroprotective extrapolations as plausible. [9,11,14,19,32]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Both drugs have substantial clinical safety experience.

Why Signaling Changes Matter for Healthspan

Many interventions that extend healthspan in model organisms converge on a small set of conserved aging pathways: **reduced insulin/IGF signaling, activation of AMPK/sirtuins, suppression of mTORC1, enhanced autophagy, improved mitochondrial function, reduced chronic inflammation, and preserved proteostasis**. [3–6]

If a therapy reliably **activates or restores** these pathways in tissues relevant to aging (metabolic organs, vasculature, brain), it provides **biologically plausible** grounds to hypothesize improved healthspan, especially when multiple complementary pathways are affected. [5,6]

Metformin: Links to Healthspan

Primary cellular effects

Mitochondrial and redox signaling: Metformin can modulate mitochondrial complex I and cellular redox balance, sometimes increasing the AMP-to-ATP ratio and activating energy-stress signaling; the mechanisms depend on tissue and exposure. [3,4] Human metabolomic observations and genetic experiments in male mice support an intestinal complex I mechanism that increases glucose utilization. [40] Adaptive stress responses

and cellular maintenance relevant to age-related resilience remain biologically plausible consequences. [5,6]

AMPK and mTOR signaling: Metformin can activate AMP-activated protein kinase and reduce mTORC1 signaling in relevant metabolic contexts, supporting glucose uptake, fatty-acid oxidation, and cellular maintenance. These effects could plausibly improve metabolic resilience and slow aspects of age-related functional decline. [3,4,33,34]

Gut-mediated effects: Metformin can increase intestinal glucose utilization, influence GLP-1 signaling and bile-acid metabolism, and alter the gut microbiome. [3,4,38,40] Enrichment of Akkermansia and several short-chain-fatty-acid-producing taxa was associated with metformin use in an observational study; a separate randomized trial demonstrated microbiome changes and improved glucose tolerance after transfer of treated donors' microbiota to mice. [38,42] These findings support plausible contributions to glucose regulation and broader metabolic and inflammatory effects. [38,42]

Downstream maintenance programs

Autophagy and proteostasis: By influencing AMPK and mTORC1 signaling, metformin could plausibly support autophagy and the clearance of damaged proteins and organelles. Greater autophagic and proteostatic activity supports improved healthspan in multiple experimental organisms. [3,33–35]

Glucose and insulin regulation: Metformin reduces hepatic glucose production and improves glucose regulation. In people with insulin resistance, prediabetes, or diabetes, this can reduce chronic hyperinsulinemia. Lower insulin exposure could plausibly support healthspan, while broader suppression of insulin-like growth factor signaling should not be assumed. [1,4,10]

Inflammatory regulation: Metformin reduced inflammatory signaling in experimental systems, including hippocampal cytokines and microglial activation in aged male mice. [34,39] A broader reduction in chronic low-grade inflammaging and its contribution to age-related disease is plausible. [5,6]

Senescence-associated signaling: Metformin suppressed components of the senescence-associated secretory phenotype through IKK/NF-κB signaling in cultured-cell experiments. [39] A broader reduction in human tissue dysfunction through these pathways is plausible. [5,6,39]

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

Metformin 500 mg orally daily; tadalafil 5 mg orally daily

Why these map to healthspan

AMPK activation, mTOR suppression, and enhanced autophagy are canonical cellular-maintenance signals in model organisms. Metformin's ability to engage these programs provides a mechanistic bridge from drug action to cellular processes that influence healthspan. [3,5,33–35]

Tadalafil: Links to Healthspan

Primary vascular / signaling effects

Endothelial signaling: Tadalafil inhibits PDE5-mediated cGMP degradation, sustaining signaling downstream of NO and promoting smooth-muscle relaxation. [25] Endothelial or vascular-marker improvements have been observed in selected human populations. [23,29,30] These effects could plausibly protect vascular-dependent tissues from aspects of age-related decline. [13,36]

Microvascular perfusion: Tadalafil can affect tissue perfusion and oxygen delivery, with responses depending on the vascular bed and disease context. [18,20,28] In a small acute study of postmenopausal women, muscle capillary recruitment increased without a corresponding increase in total forearm blood flow or glucose uptake. [20] Cerebral blood-flow findings in small-vessel disease and cardiac findings in diabetic mice support plausible, context-dependent support for cardiac, renal, skeletal-muscle, and cerebral function; preservation of multi-organ function remains a translational hypothesis. [18,20,28,37]

Downstream maintenance and protective programs

Mitochondrial protection: Eight weeks of tadalafil treatment improved cardiac mitochondrial respiration and reduced reactive-oxygen-species production in diabetic db/db mice, with changes in NO/SIRT1/PGC-1α signaling. [37] Autophagy effects are context-dependent: tadalafil reduced excessive muscle autophagy in a mouse burn-injury model. [43] Broader modulation of autophagy and mitophagy is a mechanistic hypothesis, and comparable systemic cytoprotective effects in humans are plausible. [37,43]

Vascular inflammation and platelet signaling: Human studies report changes in endothelial and related biomarkers. [23] A small platelet study found reduced P-selectin in the subgroup without cardiovascular risk factors, rather than uniformly across participants. [21] Observational outcome studies support a possible reduction in cardiovascular events. [13,36]

Neurovascular support: Tadalafil increased cerebral blood flow in a three-month trial in patients with cerebral small-vessel disease. [28] Through perfusion and cGMP-related signaling, support for synaptic function and protection from neurodegenerative processes is plausible; metabolite-clearance effects remain a mechanistic hypothesis. [7,8,17] CREB and BDNF findings across PDE5-inhibitor animal studies should be attributed to the compound and model tested. [7,17]

Lymphatic and interstitial flow: PDE5 inhibition could plausibly modulate lymphatic smooth-muscle tone and facilitate drainage in some contexts. Rat thoracic-duct experiments showed that cGMP/PKG mediates flow-dependent relaxation, which should not be equated with increased active lymphatic pumping. [26] Reports of sildenafil treatment for lymphatic malformations offer related-compound evidence. [27] Improved interstitial-fluid clearance and immune-cell trafficking with tadalafil remain speculative extrapolations. [26,27]

Why these map to healthspan

Vascular function contributes to oxygen and nutrient delivery and to susceptibility to ischemic organ injury. [18,20,28] Sustained NO–cGMP signaling therefore offers a plausible route to preserving function across vascular-dependent tissues, with observational cardiovascular findings providing additional support for that hypothesis. [13,36]

Why Metabolic and Vascular Effects May Be Additive

Metformin primarily influences cellular energy and maintenance pathways, including AMPK, autophagy, and mTOR signaling, while tadalafil primarily supports endothelial function and tissue perfusion through nitric oxide and cyclic guanosine monophosphate signaling. **Together, they plausibly address complementary metabolic and vascular aspects of aging biology.** [3,4,25,28]

Improved perfusion could support nutrient delivery, immune surveillance, and tissue repair, while improved cellular metabolic resilience could reduce downstream vascular stress. **These complementary mechanisms make additive healthspan effects plausible.** [3,5,20,28]

Practical Interpretation

Both drugs influence conserved pathways that are causally linked to healthspan in experimental models, including metabolic maintenance, autophagy, inflammatory

regulation, and vascular integrity. Their complementary effects create a biologically coherent and plausible rationale for improved human healthspan. [3–6,33–37]

Current mechanistic, preclinical, biomarker, and observational evidence supports a plausible combined healthspan benefit. [3–6,13,36] Confidence depends on convergence of relevant findings, comparable exposures, and clinically meaningful outcomes; favorable biomarkers alone do not establish an additive clinical benefit. [9,11,14,19,32]

Sex-Specific Differences in PDE5 Inhibitor Data

Most large observational PDE5-inhibitor datasets involve men; 99.4% of participants in one cardiovascular meta-analysis were male. [36] Relevant vascular and microcirculatory responses in women are biologically likely and have been observed in an acute study of postmenopausal women. [20] Broader cardiovascular and healthspan benefits in women remain possible. Tolerability may differ: in the 20 mg daily ETLAS-2 trial, treatment discontinuation was more frequent among women than men. [19]

Metformin's Exercise-Interference Debate

A twelve-week randomized exercise study in 53 older adults at risk for diabetes found that metformin attenuated improvements in cardiorespiratory fitness, insulin sensitivity, and skeletal-muscle mitochondrial respiration. Mitochondrial protein synthesis was not reduced. [11] The tested regimen was generally 2,000 mg daily, reduced to 1,500 mg for some participants. Altered mitochondrial function is a plausible mechanism, but the findings do not establish the same response to 500 mg daily or to every training and metabolic context. [11]

Dosing

The plausible, experimental daily metformin regimen for healthspan discussed here is

- **Metformin: 500 mg orally daily**

The referenced immediate-release label starts diabetes treatment at 500 mg twice daily or 850 mg once daily, with titration. [24] The 500 mg daily healthspan regimen is a conservative experimental extrapolation from metabolic and mechanistic evidence. [3–6,10]

The plausible, experimental daily tadalafil regimen for healthspan discussed here is

- **Tadalafil: 5 mg orally daily**

Metformin and Tadalafil: The Simplest Oral Rx Healthspan Stack

Metformin 500 mg orally daily; tadalafil 5 mg orally daily

This dose is approved for BPH and has been studied for daily treatment of erectile dysfunction. [22,25] Its proposed healthspan use extrapolates from vascular and mechanistic findings. [23,30,36]

Benefits

- **Metabolic regulation:** Metformin improves glucose regulation and can reduce chronic hyperinsulinemia in people with insulin resistance, prediabetes, or diabetes. [1,4,10]
- **Autophagy and stress responses:** Metformin enhanced autophagy-related processes in diabetic mouse hearts and aged mouse hippocampus. [33,34] Broader support for human cellular maintenance is plausible. [3–6]
- **Proteostasis:** Metformin activated chaperone-mediated autophagy through TAK1–IKK–Hsc70 signaling in experimental systems and improved pathology in an Alzheimer-model mouse study. [35] This complements AMPK/mTOR-related maintenance mechanisms and supports a plausible human proteostasis benefit. [3,33–35]
- **Cardiovascular outcomes:** Tadalafil exposure was associated with fewer major cardiovascular events and lower all-cause mortality in men with erectile dysfunction. [13] Convergent PDE5-inhibitor observational findings support a possible causal cardiovascular benefit, while residual confounding and differences in prescribing populations limit stronger inference. [13,36]
- **Cerebral and cognitive support:** In the ETLAS-2 MRI substudy, 20 mg tadalafil daily for three months increased whole-brain cerebral blood flow; 42 arterial-spin-labeling datasets were analyzable. [28] The cognitive substudy, involving 60 participants, found no overall benefit and one isolated slowing of reaction time, which the authors considered potentially spurious. [32] Neuroprotective effects remain plausible from animal findings, while human cognitive benefit is speculative. [7,8,17,32]

Cautions

- **Renal or hepatic impairment and metabolic acidosis:** Metformin is contraindicated at eGFR below 30 mL/min/1.73 m^2 or with metabolic acidosis; initiation at eGFR 30–45 is not recommended. [24] Daily tadalafil is not recommended with creatinine clearance below 30 mL/min or dialysis. Hepatic disease also changes prescribing suitability. [24,25]
- **Nitrates or guanylate-cyclase stimulators:** Tadalafil is contraindicated with nitrates and with agents such as riociguat because of hypotension risk. [25]

- **Cardiovascular and medication concerns:** Hypotension, unstable cardiovascular disease, alpha-blockers, antihypertensives, substantial alcohol exposure, and potent CYP3A4 inhibitors require prescribing review. [25] Acute hypoxic illness, dehydration, surgery, and certain contrast procedures can require interruption of metformin. [24]
- **Pregnancy and breastfeeding:** The proposed healthspan combination should be distinguished from indication-specific medical treatment. Metformin and tadalafil have different reproductive and lactation considerations; decisions should follow the relevant prescribing information and clinical indication. [24,25]

Side effects

- **Metformin:** Gastrointestinal symptoms are common during initiation. Vitamin B12 depletion is a cumulative concern, supported by long-term Diabetes Prevention Program follow-up. [41] Lactic acidosis is a serious risk in predisposed patients. [24]
- **Tadalafil:** Common effects include headache, dyspepsia, back pain, myalgia, congestion, and flushing. Prolonged erection and sudden visual or hearing changes require prompt medical assessment. [25]

Conclusions

Metformin and tadalafil are established prescription drugs with complementary actions relevant to metabolic maintenance and vascular function. [3,4,25] Experimental mechanisms and selected human biomarker and observational findings make a combined healthspan benefit plausible. [5,6,13,23,28,36] Dose, population, competing findings, and endpoint-specific tolerability determine the strength and applicability of that inference. [9,11,14,19,22,32,41]

References

[1] UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet*. 1998;352(9131):854–865. doi:10.1016/S0140-6736(98)07037-8

[2] Martin-Montalvo A, Mercken EM, Mitchell SJ, et al. Metformin improves healthspan and lifespan in mice. *Nat Commun*. 2013;4:2192. doi:10.1038/ncomms3192

[3] Foretz M, Guigas B, Bertrand L, Pollak M, Viollet B. Metformin: from mechanisms of action to therapies. *Cell Metab*. 2014;20(6):953–966. doi:10.1016/j.cmet.2014.09.018

[4] LaMoia TE, Shulman GI. Cellular and Molecular Mechanisms of Metformin Action. *Endocr Rev*. 2021;42(1):77–96. doi:10.1210/endrev/bnaa023

[5] Barzilai N, Crandall JP, Kritchevsky SB, Espeland MA. Metformin as a Tool to Target Aging. *Cell Metab*. 2016;23(6):1060–1065. doi:10.1016/j.cmet.2016.05.011

[6] Mohammed I, Hollenberg MD, Ding H, Triggle CR. A Critical Review of the Evidence That Metformin Is a Putative Anti-Aging Drug That Enhances Healthspan and Extends Lifespan. *Front Endocrinol (Lausanne)*. 2021;12:718942. doi:10.3389/fendo.2021.718942

[7] Puzzo D, Staniszewski A, Deng SX, et al. Phosphodiesterase 5 inhibition improves synaptic function, memory, and amyloid-beta load in an Alzheimer's disease mouse model. *J Neurosci*. 2009;29(25):8075–8086. doi:10.1523/jneurosci.0864-09.2009

[8] García-Barroso C, Ricobaraza A, Pascual-Lucas M, et al. Tadalafil crosses the blood-brain barrier and reverses cognitive dysfunction in a mouse model of AD. *Neuropharmacology*. 2013;64:114–123. doi:10.1016/j.neuropharm.2012.06.052

[9] Strong R, Miller RA, Antebi A, et al. Longer lifespan in male mice treated with a weakly estrogenic agonist, an antioxidant, an α-glucosidase inhibitor or a Nrf2-inducer. *Aging Cell*. 2016;15(5):872–884. doi:10.1111/acel.12496

[10] Kulkarni AS, Brutsaert EF, Anghel V, et al. Metformin regulates metabolic and nonmetabolic pathways in skeletal muscle and subcutaneous adipose tissues of older adults. *Aging Cell*. 2018;17(2):e12723. doi:10.1111/acel.12723

[11] Konopka AR, Laurin JL, Schoenberg HM, et al. Metformin inhibits mitochondrial adaptations to aerobic exercise training in older adults. *Aging Cell*. 2019;18(1):e12880. doi:10.1111/acel.12880

[12] American Federation for Aging Research. TAME: Targeting Aging with Metformin. Accessed October 23, 2026. AFAR trial overview

[13] Kloner RA, Stanek E, Desai K, et al. The association of tadalafil exposure with lower rates of major adverse cardiovascular events and mortality in a general population of men with erectile dysfunction. *Clin Cardiol*. 2024;47(2):e24234. doi:10.1002/clc.24234

[14] Gronich N, Stein N, Saliba W. Phosphodiesterase-5 Inhibitors and Dementia Risk: A Real-World Study. *Neuroepidemiology*. 2025;59(3):193–202. doi:10.1159/000540057

[15] Desai RJ, Mahesri M, Lee SB, et al. No association between initiation of phosphodiesterase-5 inhibitors and risk of incident Alzheimer's disease and related

dementia: results from the Drug Repurposing for Effective Alzheimer's Medicines study. *Brain Commun*. 2022;4(5):fcac247. doi:10.1093/braincomms/fcac247

[16] Abouelmagd ME, Abdelmeseh M, Elrosasy A, et al. Phosphodiesterase-5 inhibitors use and the risk of alzheimer's disease: a systematic review and meta-analysis. *Neurol Sci*. 2024;45(11):5261–5270. doi:10.1007/s10072-024-07583-9

[17] El-Bakly W, Wagdy O, Sobhy A, et al. The efficacy and underlying mechanism of phosphodiesterase- 5 inhibitors in preventing cognitive impairment and Alzheimer pathology: A systematic review of animal studies. *Behav Brain Res*. 2019;372:112004. doi:10.1016/j.bbr.2019.112004

[18] Ölmestig J, Marlet IR, Hansen RH, et al. Tadalafil may improve cerebral perfusion in small-vessel occlusion stroke-a pilot study. *Brain Commun*. 2020;2(1):fcaa020. doi:10.1093/braincomms/fcaa020

[19] Ölmestig J, Mortensen KN, Thomas MB, et al. Tadalafil Treatment in Patients With Cerebral Small Vessel Disease: The ETLAS-2 Randomized Clinical Trial. *Stroke*. 2025;56(10):2846–2857. doi:10.1161/strokeaha.125.051602

[20] Murdolo G, Sjöstrand M, Strindberg L, Lönnroth P, Jansson PA. The selective phosphodiesterase-5 inhibitor tadalafil induces microvascular and metabolic effects in type 2 diabetic postmenopausal females. *J Clin Endocrinol Metab*. 2013;98(1):245–254. doi:10.1210/jc.2012-1830

[21] De Bon E, Bonanni G, Saggiorato G, Bassi P, Cella G. Effects of tadalafil on platelets and endothelium in patients with erectile dysfunction and cardiovascular risk factors: a pilot study. *Angiology*. 2010;61(6):602–606. doi:10.1177/0003319710362977

[22] Porst H, Rajfer J, Casabé A, et al. Long-term safety and efficacy of tadalafil 5 mg dosed once daily in men with erectile dysfunction. *J Sex Med*. 2008;5(9):2160–2169. doi:10.1111/j.1743-6109.2008.00935.x

[23] Abomandour A, El-Shoura EAM, Fathi BA, Elewa HA, Abd-Eldayem AM, Zalat ZA. Effects of tadalafil on endothelial, angiogenic, metabolic, and microRNA biomarker profiles in sexually active men with BPH and erectile dysfunction: a randomized controlled trial. *Naunyn Schmiedebergs Arch Pharmacol*. Published online August 1, 2026. doi:10.1007/s00210-026-05771-2

[24] US National Library of Medicine. Metformin hydrochloride tablets: prescribing information. DailyMed. Accessed October 23, 2026. DailyMed prescribing information

[25] US National Library of Medicine. Tadalafil tablets: prescribing information. DailyMed. Accessed October 23, 2026. DailyMed prescribing information

[26] Gasheva OY, Gashev AA, Zawieja DC. Cyclic guanosine monophosphate and the dependent protein kinase regulate lymphatic contractility in rat thoracic duct. *J Physiol*. 2013;591(18):4549–4565. doi:10.1113/jphysiol.2013.258681

[27] Swetman GL, Berk DR, Vasanawala SS, Feinstein JA, Lane AT, Bruckner AL. Sildenafil for severe lymphatic malformations. *N Engl J Med*. 2012;366(4):384–386. doi:10.1056/nejmc1112482

[28] Ölmestig J, Mortensen KN, Fagerlund B, et al. Neurovascular effects of tadalafil in patients with cerebral small vessel disease: ETLAS-2 substudy. *Brain Commun*. 2026;8(4):fcag309. doi:10.1093/braincomms/fcag309

[29] Rosano GM, Aversa A, Vitale C, Fabbri A, Fini M, Spera G. Chronic treatment with tadalafil improves endothelial function in men with increased cardiovascular risk. *Eur Urol*. 2005;47(2):214–220; discussion 220–222. doi:10.1016/j.eururo.2004.10.002

[30] Amano T, Earle C, Imao T, Matsumoto Y, Kishikage T. Administration of daily 5 mg tadalafil improves endothelial function in patients with benign prostatic hyperplasia. *Aging Male*. 2018;21(1):77–82. doi:10.1080/13685538.2017.1367922

[31] Fang J, Zhang P, Zhou Y, et al. Endophenotype-based in silico network medicine discovery combined with insurance record data mining identifies sildenafil as a candidate drug for Alzheimer's disease. *Nat Aging*. 2021;1(12):1175–1188. doi:10.1038/s43587-021-00138-z

[32] Ölmestig J, Mortensen KN, Thomas MB, et al. Cognitive outcomes after tadalafil treatment in patients with cerebral small vessel disease: ETLAS-2 sub-study. *Cereb Circ Cogn Behav*. 2025;9:100520. doi:10.1016/j.cccb.2025.100520

[33] Xie Z, Lau K, Eby B, et al. Improvement of cardiac functions by chronic metformin treatment is associated with enhanced cardiac autophagy in diabetic OVE26 mice. *Diabetes*. 2011;60(6):1770–1778. doi:10.2337/db10-0351

[34] Kodali M, Attaluri S, Madhu LN, et al. Metformin treatment in late middle age improves cognitive function with alleviation of microglial activation and enhancement of autophagy in the hippocampus. *Aging Cell*. 2021;20(2):e13277. doi:10.1111/acel.13277

[35] Xu X, Sun Y, Cen X, et al. Metformin activates chaperone-mediated autophagy and improves disease pathologies in an Alzheimer disease mouse model. *Protein Cell*. 2021;12(10):769–787. doi:10.1007/s13238-021-00858-3

[36] Soulaidopoulos S, Terentes-Printzios D, Ioakeimidis N, Tsioufis KP, Vlachopoulos C. Long-term effects of phosphodiesterase-5 inhibitors on cardiovascular outcomes and death: a systematic review and meta-analysis. *Eur Heart J Cardiovasc Pharmacother*. 2024;10(5):403–412. doi:10.1093/ehjcvp/pvae029

[37] Koka S, Aluri HS, Xi L, Lesnefsky EJ, Kukreja RC. Chronic inhibition of phosphodiesterase 5 with tadalafil attenuates mitochondrial dysfunction in type 2 diabetic hearts: potential role of NO/SIRT1/PGC-1α signaling. *Am J Physiol Heart Circ Physiol*. 2014;306(11):H1558–H1568. doi:10.1152/ajpheart.00865.2013

[38] Wu H, Esteve E, Tremaroli V, et al. Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. *Nat Med*. 2017;23(7):850–858. doi:10.1038/nm.4345

[39] Moiseeva O, Deschênes-Simard X, St-Germain E, et al. Metformin inhibits the senescence-associated secretory phenotype by interfering with IKK/NF-κB activation. *Aging Cell*. 2013;12(3):489–498. doi:10.1111/acel.12075

[40] Sebo ZL, Chakrabarty RP, Grant RA, et al. Metformin inhibits mitochondrial complex I in intestinal epithelium to promote glycaemic control. *Nat Metab*. 2026;8(6):1291–1304. doi:10.1038/s42255-026-01530-y

[41] Aroda VR, Edelstein SL, Goldberg RB, et al. Long-term Metformin Use and Vitamin B12 Deficiency in the Diabetes Prevention Program Outcomes Study. *J Clin Endocrinol Metab*. 2016;101(4):1754–1761. doi:10.1210/jc.2015-3754

[42] de la Cuesta-Zuluaga J, Mueller NT, Corrales-Agudelo V, et al. Metformin Is Associated With Higher Relative Abundance of Mucin-Degrading Akkermansia muciniphila and Several Short-Chain Fatty Acid-Producing Microbiota in the Gut. *Diabetes Care*. 2017;40(1):54–62. doi:10.2337/dc16-1324

[43] Hosokawa S, Koseki H, Nagashima M, et al. Title efficacy of phosphodiesterase 5 inhibitor on distant burn-induced muscle autophagy, microcirculation, and survival rate. *Am J Physiol Endocrinol Metab*. 2013;304(9):E922–E933. doi:10.1152/ajpendo.00078.2013