GHK-Cu

1 mg daily for cosmetic goals, 2 mg daily for tissue remodeling

GHK-Cu is an endogenous copper-binding tripeptide. [15,16] GHK-Cu is proposed for skin and hair support, tissue repair, and healthspan applications, with local matrix effects demonstrated in animal wounds and systemic anti-inflammatory effects demonstrated in specific mouse models. [3–6,22,26] It can influence collagen and glycosaminoglycan production, redox signaling, and tissue remodeling. [2–6] Broader human healthspan benefits remain a plausible translational hypothesis. [23,24,28]

💉 💉 Short-term hepatic and renal tolerability is supported by reassuring tissue histology in one 28-day mouse silicosis experiment. [26] Longer-term organ tolerability is plausible as a conditional inference. [19,21]

⭐ ⭐ Local extracellular-matrix accumulation was demonstrated in rat wound chambers. [3,4] Systemic skin and hair benefits from subcutaneous (SC) administration are likely extrapolations from local and mechanistic findings. [1,17]

⭐ Whole-body healthspan benefits are a plausible mechanistic hypothesis based on tissue-repair, redox, and model-organism findings. [6,23,24,28]

C Angiogenic: Active or suspected malignancy warrants precaution because GHK-related proangiogenic and proliferative signaling could plausibly interact with tumor biology. [11,12]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

At day 28, the mouse silicosis study reported reassuring liver, kidney, and spleen histology after repeated IP GHK-Cu. [26] Favorable long-term cardiovascular, neurological, endocrine, reproductive, and carcinogenic safety remains speculative for each of these endpoints. [6,22,26]

Dose & Patterns

Human dose for SC GHK-Cu remains speculative. The community discussed SC regimens are [17,19]

- **1 mg daily for cosmetic skin and hair goals.**
- **2 mg daily for soft-tissue repair and remodeling goals.**

The following studied animal doses represented much larger human equivalent doses.

Animal Regimen	Human Equivalent Dose (150 lbs) [18]
Mouse IP — acute lung injury [22]	5.5 or 55 mg daily for 3 days
Mouse IP — smoke-related lung injury [6]	1.1, 11, or 110 mg every 48 hours for 12 weeks

Mouse IP — silica-related fibrosis [26]	11 or 110 mg twice daily, days 1–27
Mouse oral — colitis [27]	

Benefits

- **Local wound healing:** GHK-Cu increased connective-tissue accumulation in rat wound chambers. [3,4] A topical diabetic-ulcer trial also reported improved plantar-ulcer closure alongside standardized wound care. [8] Support for wound repair after human SC dosing is a plausible extrapolation across these routes and settings.
- **Local collagen synthesis:** GHK-Cu increased collagen synthesis in fibroblast cultures and collagen accumulation in rat wound chambers. [2,4] A corresponding local matrix response is biologically likely where comparable tissue exposure is achieved; a systemic cosmetic response remains an extrapolation.
- **Extracellular-matrix remodeling:** GHK-Cu modulated collagen, glycosaminoglycans, decorin, and biglycan in experimental wound tissue. [3,4] In a rat ACL-reconstruction model, intra-articular treatment improved selected laxity and stiffness outcomes at six weeks, but not ultimate load, gait, or histological scores; benefits were not maintained at twelve weeks. [7]
- **Systemic tissue repair:** Plausibly supports repair signaling beyond the administration site, supported by reduced lung injury or fibrosis after intraperitoneal (IP) treatment in mouse models. [5,6,22,26]
- **Skin quality:** Plausibly supports firmness, texture, and fine-line improvement after SC administration, extrapolating from topical cosmetic and cell findings. [1,17]
- **Scar appearance:** Plausibly supports more organized matrix remodeling and improved scar texture, a mechanistic extrapolation from wound-matrix effects. [3,4]
- **Hair growth support:** Plausibly supports follicular repair signaling and healthier hair growth after SC administration, as a translational hypothesis. [1]
- **Inflammatory regulation:** Reduced inflammatory readouts in mouse pulmonary-injury models. [5,6,22,26] Modulation of chronic age-related inflammation after human SC administration is plausible.
- **Antioxidant protection:** Improved selected oxidative-stress and antioxidant readouts in mouse lung and macrophage experiments. [6,22,26] Broader systemic antioxidant support in humans is plausible.

Drivers of Diminished Response and Mitigation Strategies

GHK-Cu has several proposed molecular interactions rather than an established receptor-agonist mechanism. [15,16,26] Downstream adaptation is still possible.

Both, continuous administration and cycling are speculative patterns. Sustained response is separate from the short treatment schedules used in injury models. [6,22,26]

GHK-Cu's Plausible Healthspan Rationale

GHK-Cu has plausible relevance to healthspan because preclinical research links it to matrix maintenance, tissue repair, and inflammatory and oxidative-stress signaling. [3–6,22,26] Plasma GHK concentrations of approximately 200 ng/mL in young adults fall to 80 ng/mL in older adults. [1,9] A connection between lower GHK availability and reduced repair capacity remains a plausible hypothesis. In an observational silicosis cohort, lower plasma GHK was associated with poorer lung function, a disease association distinct from an effect of administering GHK-Cu. [26]

Cell-based gene-expression studies and computational comparisons suggest that GHK can influence networks associated with matrix maintenance, inflammatory regulation, antioxidant defenses, DNA repair, and cellular stress responses. [23,24] A connectivity map analysis identified a GHK signature opposing emphysema-associated expression patterns, and GHK improved collagen-gel remodeling by COPD-derived lung fibroblasts in vitro. [23] These findings support a plausible transcriptional mechanism for tissue resilience; extending that mechanism to SC GHK-Cu across organs remains a translational inference.

In vitro and local wound studies show effects on collagen, glycosaminoglycans, and repair-related cells. [2–4,12] GHK-Cu also reduced pathological collagen deposition in mouse bleomycin and silica models, supporting context-dependent matrix regulation rather than indiscriminate collagen production. [5,26] Through these actions, SC GHK-Cu could plausibly support structural integrity and repair capacity in skin, connective tissue, and other organs.

Systemically administered GHK-Cu reduced inflammatory and oxidative-stress readouts in mouse lung-injury experiments, demonstrating biological effects beyond the injection site. [6,22,26] IP pretreatment before an acute inflammatory challenge and repeated IP treatment during cigarette-smoke exposure represent distinct experimental contexts. [6,22] An oral dose also improved disease and barrier-related readouts in a mouse colitis model. [27] Together, these findings make modulation of chronic inflammatory and oxidative burden plausible after suitable systemic exposure.

In vitro studies provide a possible connection between GHK-related signaling and regenerative-cell activity. GHK experiments increased VEGF and other trophic-factor secretion from human mesenchymal stromal cells, with integrin involvement. [11] Copper-free GHK increased basal-cell proliferation and stem-associated markers in engineered skin. [12] These findings support a plausible progenitor-cell and paracrine-repair hypothesis for GHK-Cu.

Taken together, matrix remodeling, redox regulation, and regenerative-cell signaling provide a biologically coherent rationale for plausible healthspan benefits from SC GHK-Cu.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[3,6,11,12,23,26] A separate C. elegans study reported improved movement, stress resistance, and mitochondrial measures with DAF-16/SKN-1 involvement, adding model-organism support for this hypothesis. [28]

Cautions

- **Reactions to copper-containing products** [19]
- **Wilson disease or other impaired copper handling** [21]
- **Active or suspected malignancy** [11,12]

Reported Side Effects

- **Injection-site pain, redness, swelling, irritation** are described anecdotally. Combining KPV and GHK-Cu in the same syringe is discussed anecdotally to prevent injection site reactions.

Biological Mechanism

GHK is glycyl-L-histidyl-L-lysine; GHK-Cu denotes its copper(II) complex. Copper coordination depends on pH, ligands, and solution conditions, and ligand exchange with other copper-binding molecules is possible. [15,16] These properties provide several possible routes to biological activity rather than a receptor-agonist mechanism.

GHK-Cu increased collagen synthesis in fibroblast cultures and altered matrix accumulation in experimental wounds. [2–4] In silica-exposed macrophage experiments, pull-down and surface-plasmon-resonance assays supported interaction with peroxiredoxin 6 (PRDX6); PRDX6 silencing weakened antioxidant and mitochondrial effects. [26] This supports a PRDX6-dependent mechanism in that system, while inflammatory and Nrf2-related findings in other lung models add convergent pathway evidence. [5,6,22]

GHK-Cu plausibly influences copper availability to copper-dependent biological processes through coordination and ligand exchange. [15,16] The behavior of a particular preparation depends on its composition, including the balance between complexed and labile copper. [25]

Exploratory analyses of GHK-treated cell expression profiles reported changes in thousands of transcripts and proposed DNA-repair and stress-response effects. [24] The COPD study provides a more focused example linking a computational signature to fibroblast behavior. [23] These findings support a plausible transcriptional basis for regeneration

Zinc Supplementation

SC GHK-Cu introduces copper directly into systemic circulation, potentially altering copper–zinc homeostasis. Intracellular copper binds strongly to metallothioneins, cysteine-rich proteins involved in metal sequestration and regulation. Copper has a higher affinity for metallothionein than zinc and can displace bound zinc, potentially altering its intracellular distribution and availability for zinc-dependent enzymes, transcription factors, and signaling pathways.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[20,21,29] Sustained copper exposure could therefore produce functional zinc insufficiency through altered metal sequestration. Unlike the well-established reduction in copper absorption caused by excessive oral zinc, clinically significant zinc depletion from subcutaneous GHK-Cu remains a mechanistically plausible but unproven effect in humans. [13,14,20,21]

Adult zinc requirements are 8 mg daily for nonpregnant, nonlactating women and 11 mg daily for men; pregnancy and lactation have different requirements. The adult tolerable upper intake level is 40 mg daily from all sources, except medically supervised therapeutic use. Sustained excess zinc can impair copper absorption and contribute to hematological, neurological, or immune problems. [20,21]

Monitoring

Baseline and periodic serum copper, ceruloplasmin, and liver biomarkers may be considered as an experimental precaution, particularly during prolonged exposure or with abnormal copper handling. Copper and ceruloplasmin are influenced by inflammation and other physiological factors, so isolated results require clinical interpretation. [21]

References

[1] Pickart L, Margolina A. Regenerative and Protective Actions of the GHK-Cu Peptide in the Light of the New Gene Data. *Int J Mol Sci*. 2018;19(7):1987. doi:10.3390/ijms19071987

[2] Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. *FEBS Lett*. 1988;238(2):343–346. doi:10.1016/0014-5793(88)80509-x

[3] Siméon A, Wegrowski Y, Bontemps Y, Maquart FX. Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu(2+). *J Invest Dermatol*. 2000;115(6):962–968. doi:10.1046/j.1523-1747.2000.00166.x

[4] Maquart FX, Bellon G, Chaqour B, et al. In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+ in rat experimental wounds. *J Clin Invest*. 1993;92(5):2368–2376. doi:10.1172/jci116842

[5] Ma WH, Li M, Ma HF, et al. Protective effects of GHK-Cu in bleomycin-induced pulmonary fibrosis via anti-oxidative stress and anti-inflammation pathways. *Life Sci*. 2020;241:117139. doi:10.1016/j.lfs.2019.117139

[6] Zhang Q, Yan L, Lu J, Zhou X. Glycyl-L-histidyl-L-lysine-Cu2+ attenuates cigarette smoke-induced pulmonary emphysema and inflammation by reducing oxidative stress pathway. *Front Mol Biosci*. 2022;9:925700. doi:10.3389/fmolb.2022.925700

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[7] Fu SC, Cheuk YC, Chiu WY, Yung SH, Rolf CG, Chan KM. Tripeptide-copper complex GHK-Cu (II) transiently improved healing outcome in a rat model of ACL reconstruction. *J Orthop Res*. 2015;33(7):1024–1033. doi:10.1002/jor.22831

[8] Mulder GD, Patt LM, Sanders L, et al. Enhanced healing of ulcers in patients with diabetes by topical treatment with glycyl-l-histidyl-l-lysine copper. *Wound Repair Regen*. 1994;2(4):259–269. doi:10.1046/j.1524-475x.1994.20406.x

[9] Pickart L, Vasquez-Soltero JM, Margolina A. The human tripeptide GHK-Cu in prevention of oxidative stress and degenerative conditions of aging: implications for cognitive health. *Oxid Med Cell Longev*. 2012;2012:324832. doi:10.1155/2012/324832

[10] Dou Y, Lee A, Zhu L, Morton J, Ladiges W. The potential of GHK as an anti-aging peptide. *Aging Pathobiol Ther*. 2020;2(1):58–61. doi:10.31491/apt.2020.03.014

[11] Jose S, Hughbanks ML, Binder BY, Ingavle GC, Leach JK. Enhanced trophic factor secretion by mesenchymal stem/stromal cells with Glycine-Histidine-Lysine (GHK)-modified alginate hydrogels. *Acta Biomater*. 2014;10(5):1955–1964. doi:10.1016/j.actbio.2014.01.020

[12] Choi HR, Kang YA, Ryoo SJ, et al. Stem cell recovering effect of copper-free GHK in skin. *J Pept Sci*. 2012;18(11):685–690. doi:10.1002/psc.2455

[13] Yuzbasiyan-Gurkan V, Grider A, Nostrant T, Cousins RJ, Brewer GJ. Treatment of Wilson's disease with zinc: X. Intestinal metallothionein induction. *J Lab Clin Med*. 1992;120(3):380–386. PubMed

[14] Sturniolo GC, Mestriner C, Irato P, Albergoni V, Longo G, D'Incà R. Zinc therapy increases duodenal concentrations of metallothionein and iron in Wilson's disease patients. *Am J Gastroenterol*. 1999;94(2):334–338. doi:10.1111/j.1572-0241.1999.851_w.x

[15] Lau SJ, Sarkar B. The interaction of copper(II) and glycyl-L-histidyl-L-lysine, a growth-modulating tripeptide from plasma. *Biochem J*. 1981;199(3):649–656. doi:10.1042/bj1990649

[16] Conato C, Gavioli R, Guerrini R, et al. Copper complexes of glycyl-histidyl-lysine and two of its synthetic analogues: chemical behaviour and biological activity. *Biochim Biophys Acta*. 2001;1526(2):199–210. doi:10.1016/s0304-4165(01)00127-1

[17] Mokhtar J, Mohamed B, Haddad J, et al. The Regenerative Potential of Glycyl-L-Histidyl-L-Lysine–Copper in Aesthetic Medicine. *Aesthet Surg J*. Published online August 20, 2026. doi:10.1093/asj/sjag169

[18] US Food and Drug Administration. *Guidance for Industry: Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers*. July 2005. Accessed October 21, 2026. FDA guidance

[19] US Food and Drug Administration. Certain Bulk Drug Substances for Use in Compounding that May Present Significant Safety Risks. Accessed October 21, 2026. FDA

[20] National Institutes of Health, Office of Dietary Supplements. Zinc: Fact Sheet for Health Professionals. Accessed October 21, 2026. NIH ODS

[21] National Institutes of Health, Office of Dietary Supplements. Copper: Fact Sheet for Health Professionals. Accessed October 21, 2026. NIH ODS

[22] Park JR, Lee H, Kim SI, Yang SR. The tri-peptide GHK-Cu complex ameliorates lipopolysaccharide-induced acute lung injury in mice. *Oncotarget*. 2016;7(36):58405–58417. doi:10.18632/oncotarget.11168

[23] Campbell JD, McDonough JE, Zeskind JE, et al. A gene expression signature of emphysema-related lung destruction and its reversal by the tripeptide GHK. *Genome Med*. 2012;4(8):67. doi:10.1186/gm367

[24] Pickart L, Vasquez-Soltero JM, Margolina A. GHK and DNA: resetting the human genome to health. *Biomed Res Int*. 2014;2014:151479. doi:10.1155/2014/151479

[25] Mateescu DM, Gavrilescu DM, Mincioaga RI, et al. GHK-Cu as a Bioactive Metallopeptide and Drug-Delivery Cargo: Coordination Chemistry, Formulation Science, Therapeutic Evidence, and a Translational Roadmap. *Pharmaceutics*. 2026;18(9):1077. doi:10.3390/pharmaceutics18091077

[26] Bian Y, Deng M, Liu J, et al. The glycyl-l-histidyl-l-lysine-Cu2+ tripeptide complex attenuates lung inflammation and fibrosis in silicosis by targeting peroxiredoxin 6. *Redox Biol*. 2024;75:103237. doi:10.1016/j.redox.2024.103237

[27] Mao S, Huang J, Li J, et al. Exploring the beneficial effects of GHK-Cu on an experimental model of colitis and the underlying mechanisms. *Front Pharmacol*. 2025;16:1551843. doi:10.3389/fphar.2025.1551843

[28] Wen H, Zhao K, Luo X, et al. The GHK-Cu delays aging in Caenorhabditis elegans via coordinated regulation of mitochondrial function and activation of DAF-16/SKN-1 pathways. *Biogerontology*. 2026;27(3):100. doi:10.1007/s10522-026-10444-x

[29] Mehlenbacher MR, Elsiesy R, Lakha R, et al. Metal binding and interdomain thermodynamics of mammalian metallothionein-3: enthalpically favoured Cu+ supplants entropically favoured Zn2+ to form Cu4+ clusters under physiological conditions. Chem Sci. 2022;13(18):5289-5304. doi:10.1039/D2SC00676F.