

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

ARA-290 (Cibinetide) is an 11-amino-acid peptide derived from erythropoietin (EPO) that selectively activates the innate-repair receptor to produce tissue-protective, anti-inflammatory, and neuroregenerative signaling without stimulating erythropoiesis. It is used experimentally for small-fiber neuropathy and other nerve-injury syndromes.

  Probably safe (some human evidence)

  Probably effective (some positive human signals)

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 6-Aug-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/ara-290.pdf>



The evidence for ARA-290 long term human safety is limited because there are **no peer-reviewed long-term safety trials**, However, **all Phase II trials showed favorable safety profile**, with **no reported erythropoietic effects** and **generally good tolerability**. Across published trials, **no clinically meaningful increases in hemoglobin, hematocrit, or reticulocyte counts** were observed, supporting that ARA-290 **does not stimulate erythropoiesis** at therapeutic doses, a critical safety distinction from full-length EPO. The most commonly reported adverse events (AE) in trials were **mild and transient**, e.g., **headache, fatigue, and occasional injection-site reactions**, with low discontinuation rates attributable to AEs. Serious adverse events directly attributed to ARA-290 have not been reported in the Phase II datasets to date.

The evidence for ARA-290 efficacy is good, with clinical reports documenting **objective and symptomatic improvements**.

Dose Used In Trials

Phase II/IIb programs tested **daily SC administration of 2 – 4 mg/day with 2mg daily producing improvement in small-fiber neuropathy symptoms, reduced pain scores, improved autonomic function, increased corneal nerve fiber density, and anti-inflammatory effects**. Human Trials at **4 mg/day producing stronger objective nerve-repair signal (corneal nerve fiber area and regenerating skin fibers)**.

ARA-290 has a **short plasma half-life** but triggers sustained tissue responses when receptor engagement exceeds nanomolar affinity.

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

- **2 - 4 mg/day for acute repair** is the plausible dose for ARA-290, based on Phase II/IIb programs

Longevity / Healthspan Dose

In vitro research shows the **Innate Repair Receptor** has **high affinity** and responds to brief, low-level pulses. Even **very small extracellular concentrations of ARA-290 initiated the repair cascade**. Across multiple cell-based assays, IRR engagement and downstream signaling, including JAK2/STAT3 activation, NF-κB suppression, and macrophage M1→M2 polarization, occur in the **low-nanomolar range**, with measurable effects often appearing at **sub-nanomolar concentrations**.

There is no detailed published PK data, but making some standard assumptions the likely C-max ranges for various subq doses of ARA-290 are

- **0.5 mg → ~0.18 – 2.39 nM**
- **1 mg → ~0.35 – 4.78 nM**
- **2.0 mg → ~0.70 – 9.56 nM**

Based on the *in vitro* work:

- **Plausible receptor engagement** was found at “**sub- to low-nM,**” so plausibly **~0.1–10 nM**.
- **Plausible functional activation window** was found at “**~0.1 nM to a few tens of nM**” for endpoints like angiogenesis, anti-inflammatory gene programs, so plausibly **~1–30 nM**.

The target tissue type should be taken into consideration when making dose decisions. For example, **disc endplate and nucleus pulposus** are relatively poorly perfused; **interstitial concentrations** there will be lower than plasma and may require **2–10×** plasma overshoot to reach robust functional levels.

This information, taken together, suggests the following plausible maintenance / longevity doses

- **0.5 mg subq overlaps the lower functional window; likely sufficient to *transiently* engage receptor signaling in well-perfused tissues and may trigger downstream signaling in some cell types.** For poorly perfused targets, this may be marginal after accounting for tissue access loss.
- **1 mg sits squarely within the lower–mid functional activation window; likely to produce a stronger and more reliable receptor engagement** signal systemically and in many tissues.
- **2.0 mg** provides a larger margin

Mechanistically, **all three doses plausibly reach the repair-cascade activation window**, with larger doses offering more certainty to account for incorrect assumptions in our mechanistically extrapolated PK data, and a stronger buffer against variability in tissue penetration and.

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

The following table shows six kinds of tissue that might plausibly benefit from ARA-290 in the context of healthspan, their sensitivity / resistance to plasma concentration, and an extrapolation of the optimal application frequency based on the tissues's turnover, ongoing low-grade stress, etc.

Tissue	Sensitivity to ARA-290 plasma concentration	Likely need for exposure frequency
Small-fiber peripheral nerves (sensory/autonomic)	High ; IRR upregulation on stressed small fibers; effects seen at sub- to low-nM in vitro.	Daily or every-other-day ; repeated pulses support ongoing repair and symptom control.
Microvascular endothelium	High-moderate ; endothelial cytoprotection and anti-inflammatory signaling activated at low-nM .	Every-other-day to weekly ; frequent pulses help maintain endothelial resilience but may tolerate spacing.
Immune cells / tissue macrophages	Moderate ; macrophage M1→M2 shifts and cytokine suppression occur in sub- to low-nM ranges.	Every-other-day ; periodic engagement reduces chronic inflammation while limiting desensitization.
Cardiac myocardium	Moderate ; cardioprotective signaling observed in preclinical models at low-nM exposures.	Every-other-day to weekly ; intermittent pulses likely sufficient for stress-resilience effects.
Central nervous system glia (microglia)	Moderate-low ; anti-inflammatory effects reported at low-nM but CNS access is limited.	Weekly or pulsed courses ; less frequent dosing may suffice due to slower CNS kinetics and tissue access limits.
Pancreatic islets and metabolic tissues	Low-moderate ; preliminary signals for β -cell protection at low-nM but evidence is limited.	Every-other-day to weekly ; conservative, intermittent exposure until stronger clinical data exist.

However, daily high dose ARA-290 for healthspan may not be practical: it is more likely to result in receptor desensitization, and it may be expensive. Taking into consideration that some tissues which benefit most from frequent dosing activate at lower plasma saturation levels, while other

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

tissues which need higher plasma concentration might not need less frequent dosing, the following is a plausible longevity dosing strategy.

- **4mg Monday, 2 mg Wednesday, 1 mg Friday for healthspan**

Mechanistically extrapolating this M/W/F 4/2/1 dose strategy to the six tissue types result in the following healthspan benefits while requiring just 28mg for four weeks.

Tissue	Likely Activation	Plausible healthspan impact of 4–2–1 mg weekly dose
Small-fiber peripheral nerves	M/W/F	Good nerve protection, moderate regeneration. This pattern could meaningfully slow small-fiber loss, reduce neuropathic symptoms, and support autonomic stability.
Microvascular endothelium	M/W/F	Solid microvascular resilience. Likely improves endothelial health, capillary perfusion, and wound-healing capacity, supporting organ function and tissue repair with age.
Immune cells / macrophages	M/W	Effective inflammaging modulation. This schedule should lower chronic inflammatory burden, good for healthspan via reduced systemic inflammation.
Cardiac myocardium	M/W	Only moderate cardio protection. May enhance stress resilience and limits low-grade cardiac inflammation over time; plausibly helpful for preserving cardiac reserve in an aging context.
CNS glia / microglia	M	Plausible modest neuroinflammatory control. May reduce microglial overactivation and support cognitive health, especially over long horizons.
Pancreatic islets / metabolic tissues	M	Supportive metabolic resilience. Likely modest improvements in stress tolerance and metabolic stability.

Here's a **single, clean, non-redundant bullet list** that merges both sets of ARA-290 benefits while removing overlap and combining similar concepts.

Benefits

- **Reduction in neuropathic pain and sensory symptoms.**
- **Objective small-fiber nerve regeneration.**

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

- **Protection and stabilization of small-fiber nerves**, helping slow nerve loss and support autonomic function over time.
- **Strong anti-inflammatory and tissue-protective signaling**, including suppression of pro-inflammatory cytokines, anti-apoptotic effects, and mitochondrial protection.
- **Plausible neuroinflammatory reduction**, supporting microglial stability and long-term cognitive health.
- **Improved microvascular resilience**, with better endothelial health, capillary perfusion, and wound-healing capacity.
- **Moderate cardioprotection**, enhancing stress resilience and reducing low-grade cardiac inflammation.
- **Supportive metabolic resilience**, with preliminary human evidence for improved glycemic and lipid markers and mechanistic support for better metabolic stress tolerance.

Indications

- **Small-fiber neuropathy (sarcoidosis-associated)** — *Human clinical data*.
- **Diabetic small-fiber neuropathy / neuropathic symptoms** — *Human clinical data (small cohorts)*.
- **Experimental use for other nerve-injury or inflammatory neuropathies** — *Theoretical / animal evidence supports exploration*.

Contraindications

- **Concurrent use of agents that require erythropoietic stimulation monitoring** — *Theoretical / safety precaution* (although ARA-290 is nonerythropoietic, caution is advised in complex hematologic disease).
- **Uncontrolled cardiovascular disease where novel vasomodulatory agents are risky** — *Theoretical / precaution from trial exclusion criteria*.

Side Effects

- **Transient systemic symptoms (fatigue, headache)** — *Human clinical data*: mild, self-limited events reported in trials.
- **No erythropoiesis or hematocrit elevation reported in trials** — *Human clinical data* (important safety distinction).

ARA-290 and Healthspan

ARA-290's relevance to aging biology comes from its ability to **reduce chronic inflammation**, **protect small-fiber nerves**, and **preserve microvascular function**, three domains that decline predictably with age. IRR expression is induced only on stressed or inflamed cells, meaning **ARA-290 acts precisely where damage is occurring**. This targeted activation leads to NF-κB

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

suppression, reduced pro-inflammatory cytokines, and a shift of macrophages toward the M2 pro-repair phenotype, supporting healthier tissue environments over time.

In human studies, ARA-290 has demonstrated improvements in small-fiber neuropathy, autonomic function, and corneal nerve fiber density, suggesting a capacity to preserve or restore neuronal integrity. Because **small-fiber degeneration is strongly associated with aging, metabolic dysfunction, and systemic inflammation, these findings form a core part of the healthspan rationale.** Additional research shows ARA-290 can improve metabolic markers, protect pancreatic β -cells, and modulate microglial activation, linking it to broader domains of metabolic resilience and neuroinflammation control.

Animal studies further support ARA-290's role in aging biology: **chronic IRR activation reduces cardiac inflammation, improves mitochondrial stress tolerance, enhances autophagy flux, and decreases lipofuscin accumulation,** all of which are central hallmarks of aging. These findings suggest that periodic activation of the IRR pathway may help maintain tissue homeostasis and reduce age-related functional decline.

ARA-290 is plausible for healthspan extension because it provides targeted anti-inflammatory signaling, nerve protection, microvascular support, and metabolic resilience. Its mechanism aligns with modern longevity frameworks focused on reducing chronic inflammatory load, preserving neuronal function, and maintaining tissue repair capacity across the lifespan.

Drivers of Diminished Response and Mitigation Strategies

For **receptor desensitization** the risk is low-moderate because multiple papers describe ARA-290 as having

- **Rapid receptor engagement**
- **Short plasma half-life**
- **Long-lasting downstream anti-inflammatory and cytoprotective signaling**

This is the **opposite of a system prone to receptor desensitization.** The **risk for neutralizing antibodies is low-moderate** because published clinical studies of ARA-290 report acceptable safety and do not highlight neutralizing antibody formation. Chronic activation of JAK2, STAT5 and related survival and anti-inflammatory pathways can induce compensatory changes in gene expression and signaling thresholds, a phenomenon described in cytokine receptor biology, so **downstream pathway adaptation through intracellular attenuation is plausible with long term use.** Homeostatic mechanisms in the immune and vascular systems can counterbalance sustained reparative signaling with any continuous healthspan application, making **system level physiological adaptation** a concern.

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

Mechanistically, cycling ARA-290 is plausibly necessary to mitigate waning. Specific cycling protocols and decisions should be viewed as experimental. A plausible cycle suggestion avoiding plateauing effects and resulting in strong response when restarting might be

- **1 month off every 6–9 months**

Monitoring

In addition to standard baseline monitoring for CBC and metabolic panels, **monitoring hematology to confirm absence of erythropoiesis** is recommended.

Applicability for Subjects with Cushing's Disease (Speculative)

ARA-290's **anti-inflammatory, pro-survival, and microvascular-protective biology** makes it a plausible candidate to address specific Cushing-related complications (small-fiber neuropathy, microvascular injury), but **no direct clinical evidence** supports its use in Cushing disease. ARA-290 targets the innate-repair receptor to trigger **anti-inflammatory, anti-apoptotic, and microvascular-protective signaling**; these actions **plausibly address several pathologies seen in Cushing disease**. Important gaps (immune dysregulation, metabolic milieu, and long-term safety) mean benefit is speculative and unproven.

Cushing disease produces chronic hypercortisolism with systemic inflammation, immune dysfunction, microvascular injury, small-fiber neuropathy, muscle wasting, and impaired tissue repair. ARA-290 selectively activates the **innate-repair receptor (EPOR/CD131)** to induce **JAK/STAT and PI3K/Akt pro-survival signaling, suppress NF-κB-driven inflammation, and improve microvascular perfusion**, mechanisms that have reduced nerve inflammation and promoted small-fiber regeneration in human neuropathy trials and in animal models.

Potentially relevant Cushing-related targets ARA-290 addresses

- **Small-fiber neuropathy and neuropathic pain:** ARA-290 improved corneal nerve fiber density and pain in sarcoidosis/diabetic neuropathy trials, suggesting it can **promote small-fiber repair and reduce neuroinflammation**, which is relevant to Cushing-associated neuropathic symptoms.
- **Microvascular dysfunction and tissue perfusion:** By **stabilizing endothelium and improving microcirculation**, ARA-290 could mitigate ischemic contributors to tissue damage and impaired wound healing seen with hypercortisolism.
- **Inflammation and immune dysregulation:** ARA-290's **anti-inflammatory signaling** may counteract the paradoxical systemic inflammation and innate immune reprogramming reported in Cushing patients.

Important reasons ARA-290 might *not* work or could be limited for Cushing's

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

- **Different disease drivers:** Cushing disease's dominant driver is **chronic glucocorticoid excess**, which causes broad endocrine, metabolic, and catabolic effects (bone loss, muscle atrophy, hyperglycemia) that ARA-290's neurovascular focus does not directly reverse.
- **Immune complexity:** Cushing patients show **concurrent systemic inflammation and cellular immunosuppression**; modulating one axis (IRR signaling) may not correct the complex immune phenotype.

Practical considerations and safety

- **Evidence stage:** ARA-290 has Phase II data in neuropathy but **no Phase III confirmation** and is investigational.
- **Monitoring and interactions:** In Cushing patients (cardiometabolic risk, infection susceptibility), any experimental therapy requires specialist oversight; potential interactions with immunomodulation or vascular status must be considered.

Biological Mechanism

ARA-290 is an **EPO-derived 11-amino-acid peptide** engineered to selectively engage the **innate repair receptor (IRR)** — a heteromeric complex of **EPOR and CD131 (βcR)** — expressed on stressed **neurons, glia, endothelial, and immune cells**. Activation of the IRR triggers **JAK2/STAT and PI3K/Akt pro-survival signaling, suppresses NF-κB-mediated proinflammatory cytokine production, and stabilizes mitochondrial function**, collectively producing **anti-apoptotic, anti-inflammatory, microvascular-protective, and pro-regenerative effects**. Crucially, ARA-290 **does not activate the homodimeric erythropoietic EPO receptor**, so it avoids red-cell stimulation and associated thrombotic risks observed with full-length EPO. Preclinical in vitro and animal models show **reduced neuronal apoptosis, improved nerve conduction and regeneration, and restored microvascular perfusion**, which align with the objective corneal nerve fiber improvements and symptom relief seen in small human trials.

References

- Brines M, Patel NS, Villa P, et al. **Nonerythropoietic, tissue-protective peptides derived from the tertiary structure of erythropoietin**. Proc Natl Acad Sci U S A. 2008;105(31):10925–10930. doi:10.1073/pnas.0805594105.
- Ueba H, Brines M, Yamin M, et al. **[Helix B surface peptide effects in cardiac/other models]**. Proc Natl Acad Sci U S A. 2010;107:14357–14362.
- Brines M, Cerami A. **The innate repair receptor: a new therapeutic target in tissue protection**. Mol Med. 2012;18:486–496.
- Heij L, Niesters M, Swartjes M, et al. **Safety and efficacy of ARA-290 in sarcoidosis patients with symptoms of small fiber neuropathy: a randomized, double-blind pilot study**. Mol Med. 2012;18:1380–1387.

ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/1mg)

- Dahan A, Dunne AN, Swartjes M, et al. **ARA-290 improves symptoms in patients with sarcoidosis-associated small nerve fiber loss and increases corneal nerve fiber density.** Mol Med. 2013;19:XXXX–XXXX.
- Brines M, Dunne AN, van Velzen M, et al. **ARA-290, a nonerythropoietic peptide engineered from erythropoietin, improves metabolic control and neuropathic symptoms in patients with type 2 diabetes.** Mol Med. 2015;21:XXXX–XXXX.