

# ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

ARA-290 (cibinetide) is an 11-amino-acid peptide modeled on erythropoietin (EPO) that selectively activates the innate-repair receptor to produce tissue-protective, anti-inflammatory, and neuroregenerative signaling without stimulating erythropoiesis. It has been studied mainly for small-fiber neuropathy; tissue-protective mechanisms are described under Biological Mechanism.



**Limited short-term** human safety data; long-term safety unknown.



**Probably effective** (some positive human efficacy signals).

*Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.*

*This document was updated 23-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/ara-290.pdf>*



**Human safety evidence is limited to small, relatively short studies, predominantly 28-day neuropathy trials and a nine-participant, 12-week eye study. These cannot establish the safety of chronic use.** Serious adverse events directly attributed to ARA-290 have not been reported in the Phase II datasets to date.

Evidence for efficacy is promising, preliminary and outcome-specific. Small randomized studies reported symptom and nerve-imaging improvements. (Heij et al., 2012; Dahan et al., 2013; Brines et al., 2015; Culver et al., 2017).

## Dose Used In Trials

Human neuropathy studies used **2 mg intravenously (IV) three times weekly for four weeks and 4 mg subcutaneously (SC) daily for 28 days. A dose-ranging sarcoidosis trial tested 1, 4, and 8 mg SC daily for 28 days. Trials found improvements in small-fiber neuropathy symptoms, reduced pain scores, improved autonomic function, increased corneal nerve fiber density, and anti-inflammatory effects. Human Trials at 4 mg/day produced stronger objective nerve-repair signal (corneal nerve fiber area and regenerating skin fibers). (Heij et al., 2012; Dahan et al., 2013; Culver et al., 2017).**

Human pharmacokinetic measurements after 4 mg SC showed a **peak plasma concentration of approximately 3 ng/mL (2.4 nM) and a terminal half-life of approximately 20 minutes. (Brines et al., 2015).**

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

- In the 64-participant dose-ranging trial, only 4 mg/day significantly improved the primary corneal nerve fiber area endpoint versus placebo. The 1 and 8 mg groups did not. This does not establish a universal optimal dose or a simple higher-dose/greater-benefit relationship (Culver et al., 2017).

### Longevity / Healthspan Dose

*In vitro* research shows the **Innate Repair Receptor** has **high affinity** and responds to brief, low-level pulses. Even **very small extracellular concentrations of ARA-290 initiated the repair cascade**. Across multiple cell-based assays, IRR engagement and downstream signaling, including JAK2/STAT3 activation, NFκB suppression, and macrophage M1→M2 polarization, occur in the **low-nanomolar range**, with measurable effects often appearing at **sub-nanomolar concentrations**.

Published human pharmacokinetic data are limited but include measured exposure after 4 mg SC and 2 mg IV. (Dahan et al., 2013).

Based on *in vitro* work:

- **Plausible receptor engagement** was found at “**sub- to low-nM.**”
- **Plausible functional activation window** was found at “**~0.1 nM to a few tens of nM**” for endpoints like angiogenesis, anti-inflammatory gene programs.

The target tissue type should be taken into consideration when making dose decisions. For example, **disc endplate and nucleus pulposus** are relatively poorly perfused; extrapolating from non-ARA-290 PK, **interstitial concentrations** there will be lower than plasma and may require **2–10×** plasma overshoot to reach robust functional levels.

**Lower intermittent doses might engage IRR signaling**, but human PK/PD data do not establish a minimum effective dose for healthspan or chronic tissue protection. Extrapolating from mechanistic biology, plausible maintenance / longevity doses

- **0.5 mg SC overlaps the lower functional window; likely sufficient to transiently engage receptor signaling in well-perfused tissues and may trigger downstream signaling in some cell types.** For poorly perfused targets, this may be marginal after accounting for tissue access loss.
- **1 mg sits squarely within the lower–mid functional activation window; likely to produce a stronger and more reliable receptor engagement** signal systemically and in many tissues.
- **2.0 mg** provides a larger margin

Mechanistically, **all three doses plausibly reach the repair-cascade activation window**, with larger doses offering more certainty to account for incorrect assumptions in our mechanistically extrapolated PK data, and a stronger buffer against variability in tissue penetration.

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

The following table summarizes the evidence relevant to six proposed tissue targets. Human concentration thresholds and preventive dosing frequencies have not been established for any of tissue that might plausibly benefit from ARA-290 in the context of healthspan, their sensitivity / resistance to plasma concentration, and an extrapolation of the optimal application frequency based on the tissues' turnover, ongoing low-grade stress, etc. This extrapolation is speculative.

| <b>Tissue</b>                                            | <b>Sensitivity to ARA-290 plasma concentration</b>                                                            | <b>Likely need for exposure frequency</b>                                                                              |
|----------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| <b>Small-fiber peripheral nerves (sensory/autonomic)</b> | <b>High;</b> IRR upregulation on stressed small fibers; effects seen at <b>sub- to low nM</b> in vitro.       | <b>Daily or every-other-day;</b> repeated pulses support ongoing repair and symptom control.                           |
| <b>Microvascular endothelium</b>                         | <b>High-moderate;</b> endothelial cytoprotection and anti-inflammatory signaling activated at <b>low nM</b> . | <b>Every-other-day to weekly;</b> frequent pulses help maintain endothelial resilience but may tolerate spacing.       |
| <b>Immune cells / tissue macrophages</b>                 | <b>Moderate;</b> macrophage M1→M2 shifts and cytokine suppression occur in <b>sub- to low nM</b> ranges.      | <b>Every-other-day;</b> periodic engagement reduces chronic inflammation while limiting desensitization.               |
| <b>Cardiac myocardium</b>                                | <b>Moderate;</b> cardioprotective signaling observed in preclinical models at <b>low nM</b> exposures.        | <b>Every-other-day to weekly;</b> intermittent pulses likely sufficient for stress-resilience effects.                 |
| <b>Central nervous system glia (microglia)</b>           | <b>Moderate-low;</b> anti-inflammatory effects reported at low nM but CNS access is limited.                  | <b>Weekly or pulsed courses;</b> less frequent dosing may suffice due to slower CNS kinetics and tissue access limits. |
| <b>Pancreatic islets and metabolic tissues</b>           | <b>Low-moderate;</b> preliminary signals for $\beta$ -cell protection at low nM but evidence is limited.      | <b>Every-other-day to weekly;</b> conservative, intermittent exposure until stronger clinical data exist.              |

However, daily high dose ARA-290 for healthspan may not be practical: it is plausibly more likely to result in receptor desensitization, and it may be expensive. Taking into consideration that some tissues which benefit most from frequent dosing activate at lower plasma saturation levels, while several other tissues which need higher plasma concentration need less frequent dosing, the following is a hypothetical longevity dosing strategy.

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

- **4mg Monday, 2 mg Wednesday, 2 mg Friday for healthspan** - *Hypothetical regimen; not studied clinically or preclinically as a healthspan protocol*

Speculatively, extrapolating this M/W/F 4/2/1 dose strategy to the six tissue types result in the following healthspan benefits while requiring just 28 mg for four weeks.

| Tissue                                | Likely Activation | Plausible healthspan impact of 4–2–2 mg weekly dose                                                                                                                                              |
|---------------------------------------|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Small-fiber peripheral nerves         | M/W/F             | <b>Good nerve protection, moderate regeneration.</b> This pattern could meaningfully slow small-fiber loss, reduce neuropathic symptoms, and support autonomic stability.                        |
| Microvascular endothelium             | M/W/F             | <b>Solid microvascular resilience.</b> May improve endothelial health, capillary perfusion, and wound-healing capacity, supporting organ function and tissue repair with age.                    |
| Immune cells / macrophages            | M/W/F             | <b>Effective inflammaging modulation.</b> This schedule may lower chronic inflammatory burden, good for healthspan via reduced systemic inflammation.                                            |
| Cardiac myocardium                    | M/W/F             | <b>Only moderate cardio protection.</b> May enhance stress resilience and limits low-grade cardiac inflammation over time; plausibly helpful for preserving cardiac reserve in an aging context. |
| CNS glia / microglia                  | M                 | <b>Plausible modest neuroinflammatory control.</b> May reduce microglial overactivation and support cognitive health, especially over long horizons.                                             |
| Pancreatic islets / metabolic tissues | M                 | <b>Supportive metabolic resilience.</b> Plausible modest improvements in stress tolerance and metabolic stability.                                                                               |

### Benefits

- **Reduction in neuropathic pain and sensory symptoms.**
- **Objective small-fiber nerve regeneration.**
- **Protection and stabilization of small-fiber nerves**, helping slow nerve loss and support autonomic function over time.
- **Strong anti-inflammatory and tissue-protective signaling**, including suppression of pro-inflammatory cytokines, anti-apoptotic effects, and mitochondrial protection.
- **Plausible neuroinflammatory reduction**, supporting microglial stability and long-term cognitive health.
- **Improved microvascular resilience**, with better endothelial health, capillary perfusion, and wound-healing capacity.

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

- **Moderate cardioprotection**, enhancing stress resilience and reducing low-grade cardiac inflammation.
- **Supportive metabolic resilience**, with preliminary human evidence for improved glycemic and lipid markers.

### Indications

- **Small-fiber neuropathy (sarcoidosis-associated)** — *Human clinical data*.
- **Diabetic small-fiber neuropathy / neuropathic symptoms** — *Human clinical data (small cohorts)*.
- **Experimental use for other nerve-injury or inflammatory neuropathies** — *Theoretical / animal evidence supports exploration*.

### Precautions

- **Concurrent use of agents that require erythropoietic stimulation monitoring** — *Theoretical / safety precaution* (although ARA-290 is nonerythropoietic, caution is advised in complex hematologic disease).
- **Uncontrolled cardiovascular disease where novel vasomodulatory agents are risky** — *Theoretical / precaution from trial exclusion criteria*.

### Side Effects

- **No erythropoiesis**. Small trials generally found no clinically significant hematologic changes or erythropoietic signal at studied doses.
- **Mild events occurred in both active and placebo groups**; the evidence does not establish headache, fatigue, or injection-site reactions as a consistent drug-attributable pattern (Heij et al., 2012; Dahan et al., 2013; Brines et al., 2015).

### ARA-290 and Healthspan

The healthspan rationale is mechanistic: tissue repair and inflammatory responses are relevant to aging biology comes from its ability to **reduce chronic inflammation, protect small-fiber nerves, and preserve microvascular function**, three domains that decline predictably with age. IRR expression is preferentially induced/upregulated in stressed or inflamed cells, meaning **ARA-290 acts where damage is occurring**. This targeted activation leads to NF-κB suppression, reduced pro-inflammatory cytokines, and a shift of macrophages toward the M2 pro-repair phenotype, supporting healthier tissue environments over time.

In human studies, ARA-290 has demonstrated improvements in small-fiber neuropathy, autonomic function, and corneal nerve fiber density, suggesting a capacity to preserve or restore neuronal integrity. Because **small-fiber degeneration is strongly associated with aging, metabolic dysfunction, and systemic inflammation, these findings form a core part of the healthspan rationale**. Additional research shows ARA-290 can improve metabolic markers,

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

protect pancreatic  $\beta$ -cells, and modulate microglial activation, linking it to broader domains of metabolic resilience and neuroinflammation control.

Animal studies further support ARA-290's role in aging biology: **chronic IRR activation is discussed for improvements in cardiac inflammation, and mitochondrial stress tolerance**, all of which are central hallmarks of aging. These findings suggest that periodic activation of the IRR pathway may help maintain tissue homeostasis and reduce age-related functional decline. **The study implicated monocyte responses rather than demonstrating general rejuvenation (Al-Onaizi et al., 2022).**

**ARA-290 is plausible for healthspan extension because it provides targeted anti-inflammatory signaling, nerve protection, microvascular support, and metabolic resilience.** Its mechanism aligns with modern longevity frameworks focused on reducing chronic inflammatory load, preserving neuronal function, and maintaining tissue repair capacity across the lifespan.

### Drivers of Diminished Response and Mitigation Strategies

**Receptor desensitization with repeated ARA-290 exposure has not been adequately studied; its likelihood is unknown, but mechanistically low-moderate** because multiple papers describe ARA-290 as having

- **Rapid receptor engagement**
- **Short plasma half-life**
- **Long-lasting downstream anti-inflammatory and cytoprotective signaling**

This is the **opposite of a system prone to receptor desensitization**. The **risk for Antibody formation** has been assessed: short studies reported no detected anti-ARA-290/cibinetide antibodies (Brines et al., 2015; Lois et al., 2020). These observations do not exclude **neutralizing antibodies during prolonged use**. **Downstream adaptation is a biological possibility, but not a demonstrated.**

### Applicability for Subjects with Cushing's Disease (Speculative)

ARA-290's **anti-inflammatory, pro-survival, and microvascular-protective biology** makes it a plausible candidate to address specific Cushing-related complications (small-fiber neuropathy, microvascular injury), but **no direct clinical evidence** supports its use in Cushing disease. ARA-290 targets the innate-repair receptor to trigger **anti-inflammatory, anti-apoptotic, and microvascular-protective signaling**; these actions **plausibly address several pathologies seen in Cushing disease**. Important gaps (immune dysregulation, metabolic milieu, and long-term safety) mean benefit is speculative and unproven.

**Cushing disease can produce neuropathic symptoms, peripheral nerve abnormalities, muscle wasting, and impaired tissue repair.** ARA-290 selectively activates the **innate-repair**

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

receptor (EPOR/CD131) to induce JAK/STAT and PI3K/Akt pro-survival signaling, suppress NFκB-driven inflammation, and improve microvascular perfusion, mechanisms that have reduced nerve inflammation and promoted small-fiber regeneration in human neuropathy trials and in animal models.

### Potentially Relevant Cushing-related targets ARA-290 addresses

- **Small-fiber neuropathy and neuropathic pain:** ARA-290 improved corneal nerve fiber density and pain in sarcoidosis/diabetic neuropathy trials, suggesting it can **promote small-fiber repair and reduce neuroinflammation**, which is relevant to Cushing-associated neuropathic symptoms.
- Microvascular dysfunction and impaired repair: experimental tissue-protection findings provide a hypothesis, but no direct evidence of benefit in hypercortisolism.
- **Inflammation and immune dysregulation:** ARA-290's **anti-inflammatory signaling** may counteract the paradoxical systemic inflammation and innate immune reprogramming reported in Cushing patients.

### Important extrapolation may fail

- **Different disease drivers:** Cushing disease's dominant driver is **chronic glucocorticoid excess**, which causes broad endocrine, metabolic, and catabolic effects (bone loss, muscle atrophy, hyperglycemia) that ARA-290's neurovascular focus does not directly reverse.
- **Immune complexity:** Cushing patients show **concurrent systemic inflammation and cellular immunosuppression**; modulating one axis (IRR signaling) may not correct the complex immune phenotype.

### Practical considerations and safety

- **Evidence stage:** neuropathy phase II findings do not establish efficacy or safety in Cushing's disease.
- **Monitoring and interactions:** disease-specific cardiometabolic, thrombotic, and infection risks warrant endocrinology oversight. Interactions with Cushing's therapies are not established (Fleseriu et al., 2021).

### Biological Mechanism

ARA-290 was designed to reproduce tissue-protective effects of EPO without stimulating its erythropoietic pathway. Its proposed target, the **innate repair receptor (IRR)**, is described as an EPOR/CD131 complex whose expression can increase with injury or inflammation. Preclinical work supports cytoprotective **and anti-inflammatory signaling, including JAK/STAT and PI3K/Akt pathways; this model does not establish receptor occupancy or clinical effects in every human tissue (Brines et al., 2008; Brines and Cerami, 2012).**

Cell and animal studies report changes in **inflammatory responses, cell survival, and nerve repair, but do not prove the clinical benefits or chronic dosing assumptions discussed**

## ARA-290 (Cibinetide) Available for Research Use Only

(Repair 2 - 4 mg / day, Longevity M/W/F 4/2/2 mg)

above. Safety observations belong to the human studies and cannot be inferred solely from receptor selectivity (Swartjes et al., 2011; Liu et al., 2014).

### References

- Brines M, Patel NS, Villa P, et al. Nonerythropoietic, tissue-protective peptides derived from the tertiary structure of erythropoietin. *Proc Natl Acad Sci U S A*. 2008;105(31):10925–10930. doi:10.1073/pnas.0805594105. [PubMed](#)
- Brines M, Cerami A. The receptor that tames the innate immune response. *Mol Med*. 2012;18:486–496. doi:10.2119/molmed.2011.00414. [PubMed](#)
- Heij L, Niesters M, Swartjes M, et al. Safety and efficacy of ARA 290 in sarcoidosis patients with symptoms of small fiber neuropathy: a randomized, double-blind pilot study. *Mol Med*. 2012;18:1430–1436. doi:10.2119/molmed.2012.00332. [PubMed Central \(PMC\)](#)
- Dahan A, Dunne A, Swartjes M, et al. ARA 290 improves symptoms in patients with sarcoidosis-associated small nerve fiber loss and increases corneal nerve fiber density. *Mol Med*. 2013;19:334–345. doi:10.2119/molmed.2013.00122. [PubMed](#)
- Brines M, Dunne AN, van Velzen M, et al. ARA 290, a nonerythropoietic peptide engineered from erythropoietin, improves metabolic control and neuropathic symptoms in patients with type 2 diabetes. *Mol Med*. 2015;20:658–666. doi:10.2119/molmed.2014.00215. [PubMed](#)
- Culver DA, Dahan A, Bajorunas D, et al. Cibinetide improves corneal nerve fiber abundance in patients with sarcoidosis-associated small nerve fiber loss and neuropathic pain. *Invest Ophthalmol Vis Sci*. 2017;58(6):BIO52–BIO60. doi:10.1167/iovs.16-21291. [PubMed](#)
- Swartjes M, Morariu A, Niesters M, et al. ARA 290, a peptide derived from the tertiary structure of erythropoietin, produces long-term relief of neuropathic pain: an experimental study in rats and  $\beta$ -common receptor knockout mice. *Anesthesiology*. 2011;115:1084–1092. doi:10.1097/ALN.0b013e31822fcefcd. [PubMed Central \(PMC\)](#)
- Muller C, Yassin K, Li LS, et al. ARA290 improves insulin release and glucose tolerance in type 2 diabetic Goto-Kakizaki rats. *Mol Med*. 2016;21(1):969–978. Published online 29-Dec-2015. doi:10.2119/molmed.2015.00267. [PubMed Central \(PMC\)](#)
- Liu Y, Luo B, Han F, et al. Erythropoietin-derived nonerythropoietic peptide ameliorates experimental autoimmune neuritis by inflammation suppression and tissue protection. *PLoS One*. 2014;9(3):e90942. doi:10.1371/journal.pone.0090942.
- Lois N, Gardner E, McFarland M, et al. A phase 2 clinical trial on the use of cibinetide for the treatment of diabetic macular edema. *J Clin Med*. 2020;9(7):2225. doi:10.3390/jcm9072225.
- Al-Onaizi MA, Thériault P, Lecordier S, et al. Early monocyte modulation by the non-erythropoietic peptide ARA 290 decelerates AD-like pathology progression. *Brain Behav Immun*. 2022;99:363–382. doi:10.1016/j.bbi.2021.07.016.
- Fleseriu M, Auchus R, Bancos I, et al. Consensus on diagnosis and management of Cushing's disease: a guideline update. *Lancet Diabetes Endocrinol*. 2021;9(12):847–875. doi:10.1016/S2213-8587(21)00235-7.