# GHK-Cu Topical Serum

## 2% GHK-Cu in 2% high- low-molecular-weight hyaluronic acid an aqueous cosmetic base

**Topical GHK-Cu is used for improving photoaged skin, eyebrow growth, and local tissue repair, with possible cosmetic benefits and mixed clinical wound-healing results. [1,13,17,24–26] GHK-Cu is a copper complex of the naturally occurring tripeptide glycyl-L-histidyl-L-lysine (Gly-His-Lys). [5,6] Copper binding and exchange contribute to its biological activity. [5,6] Fibroblast and wound experiments demonstrate effects on collagen, glycosaminoglycans, and matrix-remodeling enzymes. [2–4,21,24] Anti-inflammatory and antioxidant effects in preclinical systems provide a plausible additional rationale for skin applications. [22]**

🩹🩹🩹 Favorable local tolerability in small, short-term cosmetic studies. [13,23,24]

⭐ ⭐ Likely improvement in photoaged facial skin with formulation-specific human findings. [14,17,24]

⭐ ⭐ Possible eyebrow growth benefit after 12 weeks; scalp application remains an extrapolation. [13]

*Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.*

## Formulation

A proposed experimental formulation is 2% GHK-Cu raw powder in an aqueous cosmetic base containing 2% total mixed high- and low-molecular-weight hyaluronic acid. This is a formulation proposal, not an optimized composition established by the collagen-IV experiments. [10] The active content of the raw material and the meaning of the percentage must be specified; as advertised percentage of a diluted supplier blend is not the same as a percentage of active GHK-Cu. [11,23]

## Use Patterns

The possible cosmetic routine might be

- **Several drops 2% topical serum once or twice daily, to face or scalp**

Apply a thin, even layer to clean, dry skin, avoiding direct eye and mucosal contact. The small eyebrow trial used twice-daily application for 12 weeks. Scalp use is an experimental extrapolation from eyebrow and repair biology. [13,27]


*This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076*

# GHK-Cu Topical Serum

**2% GHK-Cu in 2% high- low-molecular-weight hyaluronic acid an aqueous cosmetic base**

## Benefits

- **Photoaged skin appearance:** Improvement in visible texture and overall appearance is possible. An older conference report described cosmetic benefits. [14] An eight-week trial evaluated a specific GHK-Cu nanocarrier serum. [24]
- **Fine lines and wrinkles:** An eight-week study reported greater wrinkle improvement with a GHK-Cu nanocarrier serum than with control serum. In a post-CO2-laser trial GHK-Cu did not improve objective wrinkle, erythema, or skin-quality measures over the control regimen at 12 weeks, although satisfaction was higher. [17]
- **Skin firmness and density:** Improvements in thickness, density, firmness, and laxity are possible. [9,14]
- **Uneven pigmentation appearance:** Improvement in mottled pigmentation is possible. [9,14]
- **Hair growth:** A randomized, double-blind study found modest increases in eyebrow hair count and diameter with nominal 2% GHK-Cu serum at 12 weeks. [13]
- **Collagen synthesis support:** GHK-Cu increases collagen synthesis in fibroblast cultures and collagen accumulation in rat experimental wounds. [2,3,24] A small human thigh-skin pilot found increased procollagen. [15] A local collagen response is plausible with effective dermal delivery. [10,11,24]
- **Matrix remodeling support:** GHK-Cu modulates glycosaminoglycans, proteoglycans, matrix metalloproteinases (MMPs), and their tissue inhibitors (TIMPs) in wound and fibroblast models. [4,21,24] Coordinated remodeling in human skin is plausible; the findings include increases as well as decreases in individual enzymes and inhibitors. [21,24]
- **Scar appearance support:** Improvement in scar texture and appearance is plausible through effects on collagen production, matrix remodeling, inflammation, and wound repair. [2–4,21,22]
- **Basement membrane support:** Support of the dermal-epidermal junction is plausible from collagen-IV findings in fibroblast and ex vivo skin experiments, including GHK-Cu combined with low-molecular-weight hyaluronic acid. [10]
- **Inflammatory signal regulation:** GHK-Cu reduced NF-κB/p38 signaling and inflammatory cytokines in stimulated macrophages and a mouse lung-injury model; application to cosmetic skin treatment is a translational inference. [22]
- **Antioxidant defense support:** Local antioxidant support is plausible from reduced reactive oxygen species and restoration of superoxide dismutase activity in the same preclinical inflammatory models. [22] Copper binding may also modify copper availability and reactivity. [5,6]

## Cautions

- **Copper allergy**. [24]
- **Active dermatitis**. [18]


*This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076*

- **Use on chronic ulcers, burns, infected wounds, or other significant injuries** belongs in supervised medical wound care. Cosmetic formulation proposals should not be treated as ulcer-treatment regimens. [1,25,26]
- **Application over a known or suspected skin cancer** warrants clinical assessment. [6,12,21,27]
- **Wilson disease or another disorder of copper handling** warrants clinician guidance. [5,18]

## Biological Mechanism

GHK binds copper with high affinity; several complexes can coexist depending on pH and ligand availability. [5,6] In experimental systems, GHK-Cu affects fibroblast collagen and elastin production, glycosaminoglycans, proteoglycans, and MMP/TIMP expression. [2–4,21,24] Keratinocyte integrin and p63 responses have been demonstrated with copper-GHK. [12,27] Angiogenic support and recruitment of repair processes remain plausible components of its remodeling biology. [3,8] Inflammatory cytokine and oxidative-stress effects have also been observed in macrophage and mouse lung-injury models. [22] These overlapping, context-dependent responses do not reduce to a single established receptor-agonist pathway. [5,6,21,22]

## Formulation Chemistry with Hyaluronic Acid

The proposed 2% GHK-Cu plus 2% total mixed-molecular-weight hyaluronic acid serum combines a mechanistically active complex with a hydrating polymer. Human studies of hyaluronic acid formulations support hydration benefits across molecular weights. [19,20] Higher-molecular-weight material can contribute surface hydration and film properties; lower-molecular-weight material may interact differently with skin. [10,19,20] GHK-Cu is hydrophilic and its stability and delivery depend on the finished formulation. [11,16,18]

In UV-exposed fibroblast and ex vivo skin models, GHK-Cu combined with low-molecular-weight hyaluronic acid enhanced collagen-IV outcomes, with the strongest tested combination at a 1:9 GHK-Cu:HA mass ratio. Collagen-I and -VII responses did not show the same synergy and some combinations were antagonistic. These findings provide a plausible rationale for combining the ingredients, but do not establish the proposed 1:1 active-mass mixture as optimal. [10]

## References

[1] Mulder GD, Patt LM, Sanders L, et al. Enhanced healing of ulcers in patients with diabetes by topical treatment with glycyl-l-histidyl-l-lysine copper. *Wound Repair Regen*. 1994;2(4):259–269. doi:10.1046/j.1524-475x.1994.20406.x

[2] Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. *FEBS Lett*. 1988;238(2):343–346. doi:10.1016/0014-5793(88)80509-x

[3] Maquart FX, Bellon G, Chaqour B, et al. In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+ in rat experimental wounds. *J Clin Invest*. 1993;92(5):2368–2376. doi:10.1172/jci116842

[4] Siméon A, Wegrowski Y, Bontemps Y, Maquart FX. Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu(2+). *J Invest Dermatol*. 2000;115(6):962–968. doi:10.1046/j.1523-1747.2000.00166.x

[5] Lau SJ, Sarkar B. The interaction of copper(II) and glycyl-L-histidyl-L-lysine, a growth-modulating tripeptide from plasma. *Biochem J*. 1981;199(3):649–656. doi:10.1042/bj1990649

[6] Conato C, Gavioli R, Guerrini R, et al. Copper complexes of glycyl-histidyl-lysine and two of its synthetic analogues: chemical behaviour and biological activity. *Biochim Biophys Acta*. 2001;1526(2):199–210. doi:10.1016/s0304-4165(01)00127-1

[7] Gorouhi F, Maibach HI. Role of topical peptides in preventing or treating aged skin. *Int J Cosmet Sci*. 2009;31(5):327–345. doi:10.1111/j.1468-2494.2009.00490.x

[8] Pickart L, Vasquez-Soltero JM, Margolina A. GHK Peptide as a Natural Modulator of Multiple Cellular Pathways in Skin Regeneration. *Biomed Res Int*. 2015;2015:648108. doi:10.1155/2015/648108

[9] Pickart L, Margolina A. Regenerative and Protective Actions of the GHK-Cu Peptide in the Light of the New Gene Data. *Int J Mol Sci*. 2018;19(7):1987. doi:10.3390/ijms19071987

[10] Jiang F, Wu Y, Liu Z, Hong M, Huang Y. Synergy of GHK-Cu and hyaluronic acid on collagen IV upregulation via fibroblast and ex-vivo skin tests. *J Cosmet Dermatol*. 2023;22(9):2598–2604. doi:10.1111/jocd.15763

[11] Badenhorst T, Svirskis D, Wu Z. Physicochemical characterization of native glycyl-l-histidyl-l-lysine tripeptide for wound healing and anti-aging: a preformulation study for dermal delivery. *Pharm Dev Technol*. 2016;21(2):152–160. doi:10.3109/10837450.2014.979944

[12] Choi HR, Kang YA, Ryoo SJ, et al. Stem cell recovering effect of copper-free GHK in skin. *J Pept Sci*. 2012;18(11):685–690. doi:10.1002/psc.2455

[13] Bo SL, Nararatwanchai T, Supornpun N, Poomarapan K, Thaploka S. The efficacy of 2% copper peptide (GHK-Cu) serum for eyebrow hypotrichosis: a randomized, double-blind, vehicle-controlled, split-face comparative study. *Procedia of Multidisciplinary Research*. 2026;4(5):16. Full text

[14] Leyden JJ, Stevens T, Finkey MB, et al. Skin care benefits of copper peptide containing facial cream. Presented at: American Academy of Dermatology Annual Meeting; 2002. Secondary citation in reference [9]. Original proceedings and complete author metadata not verified.

[15] Abdulghani AA, Sherr A, Shirin S, et al. Effects of topical creams containing vitamin C, a copper-binding peptide cream and melatonin compared with tretinoin on the ultrastructure of normal skin. *Dis Manag Clin Outcomes*. 1998;1(4):136–141. doi:10.1016/S1088-3371(98)00011-4

[16] Mazurowska L, Mojski M. Biological activities of selected peptides: skin penetration ability of copper complexes with peptides. *J Cosmet Sci*. 2008;59(1):59–69. PubMed

[17] Miller TR, Wagner JD, Baack BR, Eisbach KJ. Effects of topical copper tripeptide complex on CO2 laser-resurfaced skin. *Arch Facial Plast Surg*. 2006;8(4):252–259. doi:10.1001/archfaci.8.4.252

[18] Hostynek JJ, Dreher F, Maibach HI. Human skin retention and penetration of a copper tripeptide in vitro as function of skin layer towards anti-inflammatory therapy. *Inflamm Res*. 2010;59(11):983–988. doi:10.1007/s00011-010-0214-4

[19] Pavicic T, Gauglitz GG, Lersch P, et al. Efficacy of cream-based novel formulations of hyaluronic acid of different molecular weights in anti-wrinkle treatment. *J Drugs Dermatol*. 2011;10(9):990–1000. PubMed

[20] Muhammad P, Novianto E, Setyorini M, et al. Effectiveness of topical hyaluronic acid of different molecular weights in xerosis cutis treatment in elderly: a double-blind, randomized controlled trial. *Arch Dermatol Res*. 2024;316(6):329. doi:10.1007/s00403-024-03003-2

[21] Siméon A, Monier F, Emonard H, et al. Expression and activation of matrix metalloproteinases in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. *J Invest Dermatol*. 1999;112(6):957–964. doi:10.1046/j.1523-1747.1999.00606.x

[22] Park JR, Lee H, Kim SI, Yang SR. The tri-peptide GHK-Cu complex ameliorates lipopolysaccharide-induced acute lung injury in mice. *Oncotarget*. 2016;7(36):58405–58417. doi:10.18632/oncotarget.11168

[23] Johnson W Jr, Bergfeld WF, Belsito DV, et al. Safety Assessment of Tripeptide-1, Hexapeptide-12, Their Metal Salts and Fatty Acyl Derivatives, and Palmitoyl Tetrapeptide-7 as Used in Cosmetics. *Int J Toxicol*. 2018;37(3 suppl):90S–102S. doi:10.1177/1091581818807863

[24] Badenhorst T, Svirskis D, Merrilees M, Bolke L, Wu Z. Effects of GHK-Cu on MMP and TIMP Expression, Collagen and Elastin Production, and Facial Wrinkle Parameters. *J Aging Sci*. 2016;4(3):166. doi:10.4172/2329-8847.1000166

[25] ProCyte Corporation. *Annual report on Form 10-K for the year ended December 31, 1995*. Filed March 25, 1996. Accessed October 21, 2026. SEC filing

[26] Bishop JB, Phillips LG, Mustoe TA, et al. A prospective randomized evaluator-blinded trial of two potential wound healing agents for the treatment of venous stasis ulcers. *J Vasc Surg*. 1992;16(2):251–257. doi:10.1067/mva.1992.37086

[27] Kang YA, Choi HR, Na JI, et al. Copper-GHK increases integrin expression and p63 positivity by keratinocytes. *Arch Dermatol Res*. 2009;301(4):301–306. doi:10.1007/s00403-009-0942-x


*This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076*