

Thymosin Alpha-1 **For Research Use Only** (1.6 mg SC, once - twice weekly, various patterns)

Thymosin Alpha 1 (Ta1) is being investigated for immune support with sepsis and critical care. Ta1 could plausibly support aspects of immune function relevant to healthy aging and antiviral responses, but weekly use for longevity and as-needed use after infection exposure remain speculative schedules.

 Probably safe (Well tolerated in clinical studies; safety during indefinite use is unknown)

  Likely to produce immunomodulatory effects.

 Plausibly effective for supporting healthspan.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 4-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/thymosin-alpha-1.pdf>



Ta1 is a 28-amino-acid endogenous peptide used clinically as an immune-modulating agent (marketed as Zadaxin) that likely supports T-cell maturation and could plausibly support immune surveillance and is used clinically in some countries for specific indications.

Ta1 has generally shown favorable tolerability in clinically studied regimens. Its safety during indefinite use remains unknown.

Ta1 likely supports aspects of immune function, including antigen presentation and T-cell responses.

Ta1 could plausibly support aspects of immune function affected by aging, including T-cell responses and coordination between innate and adaptive immunity.

Dosage & Patterns

Clinical studies have evaluated several Ta1 doses and schedules, including 1.6 mg per injection. The schedule, tolerability, and effectiveness depend on the condition being treated and the treatment duration.

Clinical studies have evaluated the following dosing schedules:

- **Ta1 at 1.6 mg by subcutaneous injection twice weekly has been studied for chronic hepatitis B. Daily dosing has also been evaluated in some short-course clinical regimens.**

Thymosin Alpha-1 For Research Use Only

(1.6 mg SC, once - twice weekly, various patterns)

- **In a metastatic melanoma trial, Ta1 was adjunctly administered subcutaneously at 1.6, 3.2, or 6.4 mg per injection on days 8–11 and 15–18 of each 28-day cycle.** Treatment continued for up to six cycles unless disease progression or unacceptable toxicity occurred. This was a cancer-treatment protocol, not an established schedule for general immune support. In this melanoma trial, even at these higher doses, adding Ta1 to the cancer-treatment regimen did not produce additional toxicity compared with the control regimen.
- **In the TESTS sepsis trial, Ta1 was administered subcutaneously at 1.6 mg every 12 hours for seven days,** unless treatment ended earlier because of ICU discharge, death, or withdrawal of consent. This regimen did not significantly reduce 28-day mortality compared with placebo. In the TESTS sepsis trial, rates of adverse events and serious adverse events did not differ significantly between the Ta1 and placebo groups.

Ta1 has generally shown favorable tolerability during studied treatment periods. These findings do not establish the safety of indefinite use for general immune support or longevity.

Benefits

- **Ta1 likely supports coordination between innate and adaptive immune responses** through its effects on antigen presentation and T-cell function.
- **Ta1 could plausibly support antiviral immune responses** through its effects on pathogen recognition, antigen presentation, and T-cell function.
- **Ta1 likely supports T-cell function** through its effects on T-cell maturation, activation, and signaling.
- **Ta1 could plausibly help regulate excessive inflammatory signaling** through its effects on immune-cell activity and cytokine responses.
- **Ta1 could plausibly support recovery after illness** by helping coordinate immune responses and regulate excessive inflammatory signaling.
- When excessive inflammatory signaling contributes to mood or cognitive symptoms, Ta1 could plausibly offer indirect benefit through its immunomodulatory effects.
- When prolonged stress contributes to impaired immune function, Ta1 could plausibly provide support through its effects on antigen presentation and T-cell responses.

Potential applications

- Plausible support in selected conditions involving impaired T-cell responses.
- Plausible support for antiviral immune responses in selected viral infections.
- Plausible support for aspects of immune function affected by aging.

Cautions

- **Caution:** Safety during pregnancy and breastfeeding has not been established. Use requires an individualized assessment of potential benefits and risks by the treating clinician.

Thymosin Alpha-1 **For Research Use Only**

(1.6 mg SC, once - twice weekly, various patterns)

- Caution: Combining Ta1 with immune-stimulating cancer therapies could plausibly alter immune responses. Such combinations should be assessed by the treating oncologist.
- Caution: In autoimmune conditions, Ta1 could plausibly alter disease activity through its immunomodulatory effects. Whether those effects are beneficial or harmful may depend on the condition and should be assessed by the treating clinician.
- Contraindication: Ta1 should generally not be used in patients receiving deliberate immunosuppression, such as organ-transplant recipients, unless the treating specialist determines that the potential benefits clearly outweigh the risks.

Side Effects

- Possible fatigue; its relationship to Ta1 may be uncertain when illness or other treatments could also contribute.
- Possible headache; its relationship to Ta1 may be uncertain when illness or other treatments could also contribute.
- Injection-site soreness, redness, or discomfort.

Drivers of Diminished Response and Mitigation Strategies

The risk of receptor desensitization with repeated Ta1 use is uncertain. Adaptation within the immune-signaling pathways it influences is plausible and could reduce responsiveness over time. Whether neutralizing antibodies contribute to diminished responsiveness during repeated Ta1 use is uncertain. Immune homeostatic mechanisms could plausibly attenuate sustained immunomodulatory effects. **The overall likelihood of diminished responsiveness during prolonged Ta1 use is uncertain.**

Intermittent administration or treatment breaks could plausibly limit sustained immune-signaling adaptation, but their effectiveness in preventing diminished responsiveness is unknown.

Biological Mechanism

Ta1 is a thymic-derived immunoregulatory peptide that likely modulates pathogen recognition and T-cell responses and could plausibly help regulate excessive inflammatory signaling. Ta1 likely influences Toll-like-receptor signaling in dendritic cells and other innate immune cells, supporting antigen presentation and subsequent T-cell responses. Ta1 likely supports regulatory immune pathways in some contexts, including those involving regulatory T cells, and could plausibly help limit excessive inflammatory signaling. Together, these mechanisms provide a plausible basis for supporting immune responses while limiting inappropriate or prolonged inflammatory activation, depending on the biological context.

Thymosin Alpha-1 For Research Use Only

(1.6 mg SC, once - twice weekly, various patterns)

Conclusions

Tα1 has generally shown favorable tolerability in clinically studied regimens, but these studies do not establish long-term safety for continuous use. Supporting aspects of immune function affected is plausible.

References

- Garaci E, Pica F, Rasi G, Favalli C. Thymosin alpha 1 in the treatment of cancer: from basic research to clinical application. *Int J Immunopharmacol*. 2000;22(12):1067–1076. doi:10.1016/S0192-0561(00)00075-8.
- Romani L, Bistoni F, Montagnoli C, et al. Thymosin alpha1: an endogenous regulator of inflammation, immunity, and tolerance. *Ann N Y Acad Sci*. 2007;1112:326–338. doi:10.1196/annals.1415.002.
- Liu F, Wang HM, Wang T, Zhang YM, Zhu X. The efficacy of thymosin α1 as immunomodulatory treatment for sepsis: a systematic review of randomized controlled trials. *BMC Infect Dis*. 2016;16:488. doi:10.1186/s12879-016-1823-5.
- Ershler WB, Gravenstein S, Geloo ZS. Thymosin alpha-1 as an adjunct to influenza vaccination in the elderly: rationale and trial summaries. *Ann N Y Acad Sci*. 2007;1112:375–384.
- Costantini C, Bellet MM, Pariano M, et al. A reappraisal of thymosin alpha1 in cancer therapy. *Front Oncol*. 2019;9:873. doi:10.3389/fonc.2019.00873.
- Gu B, Zhou Y, Nie Y, et al. Efficacy of thymosin α1 for sepsis: a systematic review and meta-analysis of randomized controlled trials. *Front Cell Infect Microbiol*. 2025;15:1673959. doi:10.3389/fcimb.2025.1673959.
- Wu J, Pei F, Zhou L, et al. The efficacy and safety of thymosin α1 for sepsis (TESTS): multicentre, double blinded, randomised, placebo controlled, phase 3 trial. *BMJ*. 2025;388:e082583. doi:10.1136/bmj-2024-082583. Trial registration: NCT02867267. Correction published: *BMJ*. 2025;389:r1098. doi:10.1136/bmj.r1098.
- Goldstein AL, Low TL, McAdoo M, et al. Thymosin alpha1: isolation and sequence analysis of an immunologically active thymic polypeptide. *Proc Natl Acad Sci U S A*. 1977;74(2):725–729. doi:10.1073/pnas.74.2.725.
- Romani L, Bistoni F, Gaziano R, et al. Thymosin alpha 1 activates dendritic cells for antifungal Th1 resistance through toll-like receptor signaling. *Blood*. 2004;103(11):4232–4239. doi:10.1182/blood-2003-11-4036.
- Romani L, Bistoni F, Perruccio K, et al. Thymosin alpha1 activates dendritic cell tryptophan catabolism and establishes a regulatory environment for balance of inflammation and tolerance. *Blood*. 2006;108(7):2265–2274. doi:10.1182/blood-2006-02-004762.

Thymosin Alpha-1 For Research Use Only
(1.6 mg SC, once - twice weekly, various patterns)

- Bozza S, Gaziano R, Bonifazi P, et al. Thymosin alpha1 activates the TLR9/MyD88/IRF7-dependent murine cytomegalovirus sensing for induction of anti-viral responses in vivo. *Int Immunol*. 2007;19(11):1261–1270. doi:10.1093/intimm/dxm097.
- Mishra S, Telang G, Sureshababu A, et al. Thymosin α 1 Augments CD8⁺ T-Cell Activation and Reverses Exhaustion In Vitro. *Asian Pac J Cancer Prev*. 2026;27(6):2089–2096. doi:10.31557/APJCP.2026.27.6.2089.
- Chadwick D, Pido-Lopez J, Pires A, et al. A pilot study of the safety and efficacy of thymosin α 1 in augmenting immune reconstitution in HIV-infected patients with low CD4 counts taking highly active antiretroviral therapy. *Clin Exp Immunol*. 2003;134(3):477–481. doi:10.1111/j.1365-2249.2003.02331.x.
- Garaci E, Milanese G, Vella S, et al. A randomized controlled study for the evaluation of the activity of a triple combination of zidovudine, thymosin- α 1 and interferon- α in HIV-infected individuals with CD4 counts between 200 and 500 cells/mm³. *Antivir Ther*. 1998;3(2):103–111. doi:10.1177/135965359800300203.
- Wu J, Zhou L, Liu J, et al. The efficacy of thymosin alpha 1 for severe sepsis (ETASS): a multicenter, single-blind, randomized and controlled trial. *Crit Care*. 2013;17(1):R8. doi:10.1186/cc11932.
- Gravenstein S, Duthie EH, Miller BA, et al. Augmentation of influenza antibody response in elderly men by thymosin alpha one. A double-blind placebo-controlled clinical study. *J Am Geriatr Soc*. 1989;37(1):1–8. doi:10.1111/j.1532-5415.1989.tb01561.x.
- Iino S, Toyota J, Kumada H, et al. The efficacy and safety of thymosin alpha-1 in Japanese patients with chronic hepatitis B; results from a randomized clinical trial. *J Viral Hepat*. 2005;12(3):300–306. doi:10.1111/j.1365-2893.2005.00633.x.
- Maio M, Mackiewicz A, Testori A, et al. Large randomized study of thymosin alpha 1, interferon alfa, or both in combination with dacarbazine in patients with metastatic melanoma. *J Clin Oncol*. 2010;28(10):1780–1787. doi:10.1200/JCO.2009.25.5208.