

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

Zn-thymulin modulates T-cell maturation and likely reduces proinflammatory signaling in appropriate inflammatory contexts. Thymosin alpha-1 (Tα1) is a 28-amino-acid thymic peptide that modulates innate and adaptive immunity. Both are used experimentally to counter immunosenescence and plausibly support extended healthspan. (In this review, Thymulin refers to the administered zinc-free thymic nonapeptide, whereas Zn-thymulin refers to its zinc-bound, biologically active form.)

💉 Individually both compounds are likely safe, but safety of the combined Thymulin and thymosin alpha-1 longevity regimen is unknown.

★ Plausibly effective for healthspan extension.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 4-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/thymulin-plus-thymosin-alpha-1-for-longevity.pdf>



Aging, Longevity and the Immune system

Thymic involution progressively reduces **naïve T-cell output and thymic epithelial cross-talk**, narrowing T-cell receptor diversity and impairing adaptive responses; **this is a central mechanism linking immune aging to increased infection, cancer, and poorer tissue maintenance.**

Inflammaging, a chronic, low-grade proinflammatory state, arises from senescent cells, altered innate signaling, microbiome shifts, and persistent antigenic stimulation; it promotes organ dysfunction and accelerates multimorbidity. Large cohort and deep-phenotyping studies show immune-cell signatures of aging correlate with **short-term mortality and multimorbidity**, independent of chronological age and systemic inflammation, **supporting an association between immune aging and poorer health outcomes.** A causal contribution of immune decline to healthspan loss is plausible.

Preserving **thymic output** maintains **naïve T-cell diversity**, enabling better responses to novel pathogens and vaccines, and reducing **inflammaging** lowers chronic tissue damage

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
4-Sep-26*

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

and the senescence-associated secretory phenotype (SASP), both of which plausibly extend healthy lifespan. **Preventing immune decline plausibly improves resilience, reduces inflammaging, and therefore may extend healthspan.**

Evidence strength and gaps

Mechanistic and animal data robustly link thymic function and inflammatory signaling to tissue aging and resilience, but *randomized trials that directly demonstrate lifespan or robust healthspan extension from thymic-targeted therapies are lacking.*

Thymulin and Tα1 Supporting an Aging immune system

Zn-thymulin participates in T-cell differentiation and maturation. Through these effects, support for naïve T-cell output and maintenance of a diverse T-cell repertoire is plausible. Zn-thymulin likely has anti-inflammatory effects. **Reduced production of proinflammatory cytokines by macrophages likely contributes to these anti-inflammatory effects in inflammatory contexts.** Modulation of hypothalamic-pituitary signaling is another plausible contributing mechanism.

Tα1 likely modulates antigen-presenting cells through Toll-like receptor pathways, including activation of TLR9-linked signaling and enhancement of TLR2 responses to other stimuli. Depending on the cellular and inflammatory context, these effects involve MyD88-dependent signaling and downstream regulators such as NF-κB, MAPKs, and IRF7. These pathways **likely support antigen presentation and T-cell activation through increased IL-12 production, type I interferon responses, and expression of antigen-presenting and co-stimulatory molecules.** The specific response depends on the cell type and immune stimulus. **Tα1 likely promotes regulatory T-cell development and immune tolerance** in some contexts by activating indoleamine 2,3-dioxygenase (IDO), an enzyme involved in tryptophan breakdown, in dendritic cells. At the cellular level, these signaling events **plausibly enhance natural killer cell cytotoxicity and CD8+ cytotoxic T-cell function and reduce expression of markers associated with T-cell exhaustion.** These effects could support antigen-specific responses and modulate inflammatory signaling, depending on the immune context.

Combining Thymulin and Tα1 plausibly supports complementary aspects of immune function: thymic T-cell maturation through formation of active Zn-thymulin, and peripheral antigen presentation and immune-cell responses through Tα1. Through its role in T-cell maturation, Zn-thymulin plausibly supports the availability of naïve T cells that can respond to newly encountered antigens. Tα1 likely supports dendritic-cell maturation,

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Ta1 1.6 mg day 2, once weekly)

antigen presentation, and immune signaling that helps naïve T cells become activated in response to an antigen. These effects plausibly complement Zn-thymulin's role in T-cell maturation. **This pairing plausibly improves responses to newly encountered antigens and vaccines and helps regulate chronic inflammatory signaling.** The combination is therefore a biologically plausible strategy for supporting thymic and peripheral immune function and countering aspects of age-related immune decline.

Plausible Longevity / Healthspan Benefits

- **Plausibly improved adaptive responses to new pathogens and vaccines:** Ta1 likely supports antigen presentation and T-cell activation, while Zn-thymulin's role in T-cell maturation could complement these effects.
- **Plausibly reduced susceptibility to or severity of some infections:** Complementary effects on T-cell maturation, antigen presentation, and immune-cell responses could support pathogen clearance and limit infection-related inflammatory damage.
- **Plausibly improved immune surveillance of abnormal cells:** Complementary effects on T-cell maturation, antigen presentation, and immune-cell function could support recognition and elimination of abnormal cells.
- **Plausibly reduced age-related chronic inflammation and associated tissue damage:** Zn-thymulin and Ta1 could complement one another in regulating inflammatory signaling and immune responses, potentially limiting persistent inflammation and the tissue damage it causes.
- **Plausible support for immune recovery after acute physiological stress:** Complementary effects on T-cell maturation and immune regulation could help restore immune function when acute illness or injury has disrupted it.

Cautions

- **Active autoimmune disease with unstable control,** theoretical risk of immune modulation worsening autoimmunity. (Ta1 has sometimes been explored in autoimmune contexts with mixed results.)
- **Zinc status and supplementation:** Adequate zinc availability supports Zn-thymulin activity.

Drivers of Diminished Response and Mitigation Strategies

The overall likelihood of diminished response with repeated Thymulin administration is unknown. The overall likelihood of diminished response with repeated Ta1 administration

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Ta1 1.6 mg day 2, once weekly)

is unknown. Intermittent administration is a speculative strategy for limiting diminished response; whether it preserves responsiveness to Thymulin or Ta1 is unknown.

Doses, Patterns, Sequences

Thymulin dosing

No established dosing regimen exists for Thymulin use for longevity. **Earlier rheumatoid arthritis trials tested Thymulin at 1, 5, and 10 mg/day**, but these regimens do not establish an appropriate dose or schedule for longevity. Nor is there consensus among FRU users, with doses ranging from microgram-scale to low milligram scale, and patterns ranging from short induction courses (several days to two weeks) to intermittent maintenance courses (once or several times a week.) Some users report occasional use, while others using low-dose maintenance courses report continuous use.

Thymosin alpha-1 dosing

The **1.6 mg twice-weekly** regimen is the most consistently used and is the basis for many clinical trials and regulatory approvals outside the U.S. **The risk of antibody formation against Ta1 during repeated administration is not well characterized.**

Sequencing

Administering **Thymulin one day before Ta1 is a speculative sequencing strategy intended to place potential thymic support before peripheral immune modulation.**

Zn-thymulin participates in T-cell maturation. Ta1 likely supports antigen presentation and immune-cell activation, So boost maturation, then boost activation.

Mechanistically, separating administration could plausibly alter the overlap of the peptides' biological effects, reducing excessive signaling.

Speculative Dosing Schedule

- **Thymulin, Day 1, once a week, 1 mg**
- **Thymosin alpha-1 , Day 2, once a week, 1.6 mg**

There is no evidence to support a decision to use this combined protocol just once, or more than once, per week. There is no evidence if cycling is necessary for Thymulin or Ta1, or what those cycle patterns might look like.

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

Zinc Supplementation

Zinc is an essential cofactor for Thymulin activity and a central micronutrient for innate and adaptive immunity. Adequate zinc status supports **thymic hormone function, thymocyte maturation, T-cell development, and multiple zinc-dependent enzymes and transcription factors** that regulate immune signaling. **Adequate zinc availability could also support formation of active Zn-thymulin after administration of zinc-free Thymulin.**

Population data show **zinc deficiency or insufficiency is relatively common in older adults** because of reduced intake, absorption changes, polypharmacy, and chronic disease; this makes zinc assessment and correction a logical adjunct when targeting thymic regeneration or immune support.

Serum zinc, serum copper, and ceruloplasmin may help assess zinc-copper balance and inform supplementation decisions when interpreted alongside dietary intake and clinical context.

Dietary reference / typical targets: the U.S. Recommended Dietary Allowance (RDA) is **about 11 mg/day for adult men and 8 mg/day for adult women**; requirements are higher in pregnancy and lactation. Supplemental zinc dosing should be individualized when dietary intake or zinc status is inadequate, rather than treating a particular dose range as necessary for this peptide regimen.

The adult tolerable upper intake level is 40 mg of elemental zinc daily from food, supplements, and medications combined. Chronic excessive intake can cause copper deficiency, anemia, low neutrophil counts, and neurological or immune problems.

References

Franceschi C, Bonafè M, Valensin S, Olivieri F, De Luca M, Ottaviani E, De Benedictis G. Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci.* 2000;908:244-254. *Franceschi et al 2000 is the foundational, highly cited conceptual review that coined and developed the inflammaging framework linking chronic low-grade inflammation to age-related disease and immune decline. It provides a conceptual framework linking chronic antigenic stimulation and inflammation with immune aging and age-related disease.*

Cevenini E, Monti D, Franceschi C. Inflamm-ageing. *Curr Opin Clin Nutr Metab Care.* 2013;16(1):14-20. *Cevenini et al 2013 is a concise clinical-nutrition perspective that*

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
4-Sep-26*

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

synthesizes inflammaging mechanisms (senescent cells, microbiome, nutrition) and practical modulators relevant to healthspan interventions.

Nikolich-Žugich J. The twilight of immunity: emerging concepts in aging of the immune system. *Nat Immunol.* 2018;19(1):10-19. *Nikolich-Žugich 2018 is a modern, authoritative review of immunosenescence in Nature Immunology that details thymic involution, naïve T-cell loss, repertoire narrowing, innate immune remodeling, and how these changes affect infection, vaccination, and cancer risk.*

Bach JF, Dardenne M, Pleau JM, Rosa J. Biochemical characterisation of a serum thymic factor. *Nature.* 1977;266:55–57. doi:10.1038/266055a0.

Ershler WB, Gravenstein S, Geloo ZS. Thymosin alpha-1 as an adjunct to influenza vaccination in the elderly: rationale and trial summaries. *Ann N Y Acad Sci.* 2007;1112:375-384.

Costantini C, Bellet MM, Pariano M, et al. A reappraisal of thymosin alpha1 in cancer therapy. *Front Oncol.* 2019;9:873. doi:10.3389/fonc.2019.00873.

Giunta S. Is inflammaging an auto[innate]immunity subclinical syndrome? *Immun Ageing.* 2006;3:12. doi:10.1186/1742-4933-3-12.

Gu B, Zhou Y, Nie Y, et al. Efficacy of thymosin α1 for sepsis: a systematic review and meta-analysis of randomized controlled trials. *Front Cell Infect Microbiol.* 2025;15:1673959. doi:10.3389/fcimb.2025.1673959.

Wu J, Pei F, Zhou L, et al. The efficacy and safety of thymosin α1 for sepsis (TESTS): multicentre, double blinded, randomised, placebo controlled, phase 3 trial. *BMJ.* 2025;388:e082583. doi:10.1136/bmj-2024-082583. Trial registration: NCT02867267. Correction published: *BMJ.* 2025;389:r1098. doi:10.1136/bmj.r1098.

Goldstein AL, Low TL, McAdoo M, et al. Thymosin alpha1: isolation and sequence analysis of an immunologically active thymic polypeptide. *Proc Natl Acad Sci U S A.* 1977;74(2):725–729. doi:10.1073/pnas.74.2.725.

Dardenne M, Savino W, Berrih S, Bach JF. A zinc-dependent epitope on the molecule of thymulin, a thymic hormone. *Proc Natl Acad Sci U S A.* 1985;82(20):7035–7038. doi:10.1073/pnas.82.20.7035.

Prasad AS, Meftah S, Abdallah J, Kaplan J, Brewer GJ, Bach JF, Dardenne M. Serum thymulin in human zinc deficiency. *J Clin Invest.* 1988;82(4):1202–1210. doi:10.1172/JCI113717.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
4-Sep-26*

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

Amor B, Dougados M, Mery C, Dardenne M, Bach JF. Nonathymulin in rheumatoid arthritis: two double blind, placebo controlled trials. *Ann Rheum Dis.* 1987;46(7):549–554. doi:10.1136/ard.46.7.549.

Romani L, Bistoni F, Gaziano R, et al. Thymosin alpha 1 activates dendritic cells for antifungal Th1 resistance through toll-like receptor signaling. *Blood.* 2004;103(11):4232–4239. doi:10.1182/blood-2003-11-4036.

Romani L, Bistoni F, Perruccio K, et al. Thymosin alpha1 activates dendritic cell tryptophan catabolism and establishes a regulatory environment for balance of inflammation and tolerance. *Blood.* 2006;108(7):2265–2274. doi:10.1182/blood-2006-02-004762.

Bozza S, Gaziano R, Bonifazi P, et al. Thymosin alpha1 activates the TLR9/MyD88/IRF7-dependent murine cytomegalovirus sensing for induction of anti-viral responses in vivo. *Int Immunol.* 2007;19(11):1261–1270. doi:10.1093/intimm/dxm097.

Lunin SM, Khrenov MO, Novoselova TV, Parfenyuk SB, Novoselova EG. Thymulin, a thymic peptide, prevents the overproduction of pro-inflammatory cytokines and heat shock protein Hsp70 in inflammation-bearing mice. *Immunol Invest.* 2008;37(8):858–870. doi:10.1080/08820130802447629.

National Institutes of Health, Office of Dietary Supplements. Zinc: Fact Sheet for Health Professionals. <https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/>. Accessed September 3, 2026.

Reggiani PC, Morel GR, Cónsole GM, et al. The thymus-neuroendocrine axis: physiology, molecular biology, and therapeutic potential of the thymic peptide thymulin. *Ann N Y Acad Sci.* 2009;1153:98–106. doi:10.1111/j.1749-6632.2008.03964.x.

Mishra S, Telang G, Sureshbabu A, et al. Thymosin α1 Augments CD8⁺ T-Cell Activation and Reverses Exhaustion In Vitro. *Asian Pac J Cancer Prev.* 2026;27(6):2089–2096. doi:10.31557/APJCP.2026.27.6.2089.

Zhao XH, Awaya A, Kobayashi H, Ohnuki T, Tokumitsu Y, Nomura Y. Effects of repeated administrations of facteur thymique sérique (FTS) on biochemical changes related to aging in senescence-accelerated mouse (SAM). *Jpn J Pharmacol.* 1990;53(3):311–319. doi:10.1254/jjp.53.311.

Yara S, Kawakami K, Kudeken N, et al. FTS reduces bleomycin-induced cytokine and chemokine production and inhibits pulmonary fibrosis in mice. *Clin Exp Immunol.* 2001;124(1):77–85. doi:10.1046/j.1365-2249.2001.01493.x.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026

Researcher6076

*Updated:
4-Sep-26*

Thymulin & Thymosin alpha-1 for Healthspan Available for Research Use Only

(Thymulin 1.0 mg Day 1, Tα1 1.6 mg day 2, once weekly)

Lunin SM, Khrenov MO, Novoselova TV, et al. Modulation of inflammatory response in mice with severe autoimmune disease by thymic peptide thymulin and an inhibitor of NF-kappaB signalling. *Int Immunopharmacol*. 2015;25(2):260–266. doi:10.1016/j.intimp.2015.01.021.

da Silva AL, Martini SV, Abreu SC, et al. DNA nanoparticle-mediated thymulin gene therapy prevents airway remodeling in experimental allergic asthma. *J Control Release*. 2014;180:125–133. doi:10.1016/j.jconrel.2014.02.010.

Chadwick D, Pido-Lopez J, Pires A, et al. A pilot study of the safety and efficacy of thymosin α1 in augmenting immune reconstitution in HIV-infected patients with low CD4 counts taking highly active antiretroviral therapy. *Clin Exp Immunol*. 2003;134(3):477–481. doi:10.1111/j.1365-2249.2003.02331.x.

Garaci E, Milanese G, Vella S, et al. A randomized controlled study for the evaluation of the activity of a triple combination of zidovudine, thymosin-α1 and interferon-α in HIV-infected individuals with CD4 counts between 200 and 500 cells/mm³. *Antivir Ther*. 1998;3(2):103–111. doi:10.1177/135965359800300203.

Wu J, Zhou L, Liu J, et al. The efficacy of thymosin alpha 1 for severe sepsis (ETASS): a multicenter, single-blind, randomized and controlled trial. *Crit Care*. 2013;17(1):R8. doi:10.1186/cc11932.