

## Thymulin Available for Research Use Only

### (250 mcg – 500 mcg 2–3×/week, various patterns)

**Zn-thymulin modulates T-cell maturation and likely reduces proinflammatory signaling in appropriate inflammatory contexts. Human clinical evidence is limited and comes mainly from small, older studies, while animal and in vitro research provide broader evidence of immunomodulatory activity.** (In this review, Thymulin refers to the administered zinc-free thymic nonapeptide, whereas Zn-thymulin refers to its zinc-bound, biologically active form.)

💉💉 Probably safe

★ ★ Likely to produce immunomodulatory effects

★ Plausibly effective for clinical benefits

*Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.*



*This document was updated 4-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/thymulin.pdf>*

**Zn-thymulin is the zinc-bound, biologically active form of the thymic nonapeptide Thymulin. It modulates T-cell maturation and peripheral immune function. Its use to reduce chronic inflammatory signaling or provide adjunctive benefit in autoimmune conditions is plausible.**

Human safety data are lacking for long-term use. Interactions with immunosuppressants and biologics are not well characterized.

Human research links reduced endogenous Zn-thymulin activity with aging, malnutrition, and zinc deficiency, and shows that zinc repletion can restore its activity in deficient states. These findings support the biology of Zn-thymulin. Based on mechanistic, in vitro, animal, and limited exploratory human evidence, clinical benefit from administering Thymulin for inflammatory or age-related immune conditions is plausible.

### Zn-thymulin vs Thymulin

Thymulin and Zn-thymulin are not biologically equivalent. Thymulin's established biological activity requires binding one zinc ion; therefore, the benefits discussed in this review apply specifically to zinc-bound thymulin, Zn-thymulin. Endogenous thymulin can bind available zinc, and human evidence shows that zinc supplementation can increase biologically active Zn-thymulin. Based on this biology, it is mechanistically plausible that zinc-free thymulin administered subcutaneously could bind bioavailable zinc in the body and become active Zn-

## **Thymulin Available for Research Use Only**

### **(250 mcg – 500 mcg 2–3×/week, various patterns)**

thymulin, and adequate zinc intake may support that process. However, this conversion has not been examined.

The distinction is analogous to GHK and GHK-Cu: gray-market GHK-Cu is supplied with copper already bound, whereas products labeled simply “Thymulin” are sold without bound zinc, as verified by common lab test reports.

### **Dose & Patterns**

No generally established dose or schedule exists for the uses discussed in this review. Older rheumatoid-arthritis trials tested 1, 5, and 10 mg/day, but those condition-specific regimens do not establish dosing for other uses. The schedule below is retained as speculative community practice reported by some practitioners and peptide users.

Thinking about timing and patterns for Thymulin revolves around two timescales.

**Rapid signaling:** In vitro research shows that Zn-thymulin can modulate NF-κB signaling and reduce lipopolysaccharide-induced IL-1β release in alveolar epithelial cells. A relatively rapid anti-inflammatory signaling effect is therefore plausible, but its timing and magnitude after subcutaneous administration of Thymulin in humans are unknown.

**Thymic/central effects:** Zn-thymulin participates in thymocyte maturation. Through this role, effects on central tolerance and the peripheral T-cell repertoire are plausible.

Specific thymulin binding has been demonstrated in human lymphoblastoid T-cell lines, and functional studies show effects on T-cell maturation and immune signaling. However, the molecular identity of the receptor and the complete signaling pathways are incompletely defined.

The following speculative regimen is discussed in practitioner and peptide-user communities:

- **Thymulin: 250–500 mcg, two or three times weekly**

Sometimes practitioners discuss using this dose for four to six weeks, followed by four to twelve weeks off.

### **Alternate Dose and Pattern Combining Thymulin and Thymosin Alpha-1**

Combining administered Thymulin with thymosin alpha-1 is mechanistically plausible because Zn-thymulin and thymosin alpha-1 have partly distinct immunomodulatory actions. The combination has not been studied, and their use together is based solely on biological mechanism and extrapolation.

The following speculative combined regimen is discussed in practitioner and peptide-user communities:

## **Thymulin Available for Research Use Only**

### **(250 mcg – 500 mcg 2–3×/week, various patterns)**

- **Thymulin: 1 mg once weekly on Day 1.**
- **Thymosin alpha-1: 1.6 mg once weekly on Day 2.**

Splitting these doses into twice weekly administration is also discussed.

### **Benefits**

- **Likely modulation of T-cell function and thymic signaling.** Zn-thymulin influences thymocyte maturation and peripheral T-cell markers in animal and in vitro studies.
- **Likely reduction of NF-κB-related inflammatory signaling in appropriate inflammatory contexts.** Zn-thymulin can modulate this pathway and reduce production of selected proinflammatory cytokines.
- **Plausible reduction in the frequency or duration of viral infections.** Zn-thymulin's role in T-cell maturation and immune signaling provides a mechanistic rationale, and some users report fewer or milder infections, but these reports are anecdotal.
- **Plausible support for recovery of immune function after an immune stressor.** Zn-thymulin's effects on T-cell maturation and immune signaling provide a mechanistic rationale, and short-course use after illness is reported by some practitioners and users, but these reports are anecdotal.
- **Likely anti-inflammatory effects in some experimental inflammatory models.** Benefit in particular human chronic inflammatory conditions is plausible and depends on the disease context.
- **Plausible complementary effects with thymosin alpha-1.** Zn-thymulin and thymosin alpha-1 have partly distinct immunomodulatory actions, providing a mechanistic rationale for complementary effects; reported combined use is anecdotal.

### **Indications**

- **Plausible potential application as adjunctive immune support for recurrent infections.** Zn-thymulin's role in T-cell maturation provides a mechanistic rationale; reported use is anecdotal.
- **Plausible potential application as adjunctive support during postviral recovery.** Zn-thymulin's immunomodulatory effects provide a mechanistic rationale; reported use is anecdotal.
- **Plausible support for age-related decline in T-cell function.** Endogenous Zn-thymulin activity declines with aging, and its role in thymocyte maturation provides a biological rationale for this potential application.
- **Likely reduction of age-associated inflammatory signaling in myeloid immune cells.** Zn-thymulin likely limits excessive proinflammatory cytokine production by these cells through modulation of NF-κB signaling.

## **Thymulin Available for Research Use Only**

### **(250 mcg – 500 mcg 2–3×/week, various patterns)**

- **Plausible adjunctive use with thymosin alpha-1 for complementary immunomodulation.** Their partly distinct immunomodulatory actions provide a mechanistic rationale; reported combined use is anecdotal.
- **Plausible potential adjunct in selected chronic inflammatory or autoimmune contexts.**

### **Cautions**

- **Active or recent malignancy:** use should be reviewed with the treating oncologist because the effects of Thymulin or Zn-thymulin on tumor immunity and cancer treatment are unknown.
- **Unstable or uncontrolled autoimmune disease with active severe flares:** avoid unsupervised immune-active interventions.
- **Unsupervised use in people on potent immunosuppressants or biologics** (risk of unpredictable interactions).
- **Pregnancy and breastfeeding** — insufficient safety data.
- **Uncorrected zinc deficiency when administering zinc-free Thymulin,** because inadequate zinc availability could plausibly limit formation of active Zn-thymulin.

### **Drivers of Diminished Response and Mitigation Strategies**

Receptor desensitization, neutralizing-antibody formation, downstream adaptation, and overall waning of response have not been characterized for administered Thymulin or Zn-thymulin. Diminished responsiveness during repeated exposure is theoretically possible by analogy with other peptide-signaling systems.

By analogy with other peptide and cytokine signaling systems, intermittent administration of Thymulin could plausibly reduce a theoretical risk of diminished responsiveness during continuous exposure.

### **Adequate Zinc Status and Thymulin**

Zn-thymulin is the biologically active, zinc-bound form of Thymulin. When zinc-free Thymulin is administered, adequate zinc availability could plausibly support formation of active Zn-thymulin. Zinc supplementation may be appropriate to correct inadequate intake or zinc deficiency.

Zinc intake should generally meet established dietary requirements, which are 8 mg daily for adult women and 11 mg daily for adult men. The adult tolerable upper intake level of 40 mg daily includes zinc from food, supplements, and medications.

Serum zinc, serum copper, and ceruloplasmin may help assess zinc-copper balance.

### **Thymulin “Cured” My Morning Congestion**

***I experienced chronic low-grade postnasal drip and morning congestion for as long as I can remember. Since beginning Thymulin once or twice weekly, those symptoms resolved.***

## **Thymulin Available for Research Use Only**

### **(250 mcg – 500 mcg 2–3×/week, various patterns)**

*A plausible explanation is that administered Thymulin bound available zinc and formed active Zn-thymulin. In vitro research shows that Zn-thymulin can reduce NF-κB activation and lipopolysaccharide-induced IL-1β release in alveolar epithelial cells, and airway-directed thymulin gene therapy reduced inflammation and mucus-related pathology in a mouse model of allergic asthma. Speculatively, if related immunomodulatory effects occurred in the nasal mucosa, they could plausibly reduce inflammatory signaling, mucosal swelling, vascular leakage, and mucus production, thereby improving nasal airflow and mucociliary clearance.*

### **Use with autoimmune conditions**

Zn-thymulin is best described as an immunomodulatory thymic peptide, not as an established broad immunosuppressant. It likely reduces NF-κB-related inflammatory signaling in appropriate inflammatory contexts and may indirectly create conditions favorable to regulatory T-cell stability. Potential use in autoimmune disease remains experimental.

#### **Hashimoto's thyroiditis**

Zn-thymulin's effects on T-cell maturation and NF-κB-related inflammatory signaling provide a plausible mechanistic rationale for modulating autoimmune inflammation in Hashimoto's thyroiditis.

Anecdotal users report reduced inflammatory symptoms and improved resilience.

#### **Rheumatoid arthritis (RA)**

Small, older human studies provide preliminary evidence in rheumatoid arthritis. Zn-thymulin improved abnormal T-cell subset markers in vitro in lymphocytes from eight of nine patients, an open clinical report described improvement, and two randomized, double-blind, placebo-controlled trials reported greater clinical improvement with thymulin 5 mg/day than with placebo. These findings support plausible symptomatic benefit in rheumatoid arthritis.

Some users report reduced morning stiffness or milder flares, but these reports are anecdotal. T

#### **Rheumatoid arthritis with interstitial lung disease (RA-ILD)**

Zn-thymulin's immunomodulatory and NF-κB-related effects provide a plausible rationale for modulating the inflammatory component of RA-ILD. Limiting fibrosis during inflammatory lung injury is also plausible, supported by reduced fibrosis with early FTS treatment in an experimental lung-injury model. This does not establish reversal of existing lung scarring.

### **Biological Mechanism**

Thymic epithelial cells produce the nonapeptide component of thymulin. After binding zinc, it forms biologically active Zn-thymulin, an endocrine and paracrine modulator involved in T-cell development and peripheral immune function.

Mechanistically, Zn-thymulin exerts multiple effects:

**Thymulin Available for Research Use Only**  
**(250 mcg – 500 mcg 2–3×/week, various patterns)**

- **Modulation of thymocyte maturation and peripheral T-cell phenotype.** Zn-thymulin influences thymocyte maturation and can alter peripheral T-cell markers. Effects on thymic selection, the peripheral T-cell repertoire, and central tolerance are plausible, but the specific receptor and signaling pathways remain incompletely defined.
- **Modulation of NF-κB-related inflammatory signaling.** Zn-thymulin likely reduces NF-κB-related inflammatory signaling in appropriate inflammatory contexts. In vitro research shows reduced NF-κB nuclear translocation and lipopolysaccharide-induced IL-1β release in alveolar epithelial cells, while animal studies support modulation of this pathway during inflammation. Broader anti-inflammatory effects are likely in appropriate inflammatory contexts, including reductions in elevated circulating proinflammatory cytokines.
- **Possible indirect effects on regulatory T-cell balance.** Zn-thymulin's roles in thymocyte maturation and inflammatory signaling provide a plausible rationale for indirect effects on regulatory T-cell function.
- **Neuroendocrine interactions.** Endogenous Zn-thymulin participates in communication between the thymus, immune system, and neuroendocrine system. Administered Thymulin could plausibly influence these interactions after forming active Zn-thymulin, but clinically meaningful neuroendocrine effects have not been established.

These effects provide a rationale for investigating potential applications in recurrent infections, immune dysfunction, and selected inflammatory or autoimmune conditions.

## References

- Amor B, Dougados M, Mery C, Dardenne M, Bach JF. Nonathymulin in rheumatoid arthritis: two double blind, placebo controlled trials. *Ann Rheum Dis.* 1987;46(7):549–554. doi:10.1136/ard.46.7.549. PMID: 3310925.
- Amor B, Dougados M, Mery C, de Gery A, Choay J, Dardenne M, Bach JF. Thymuline (FTS) in rheumatoid arthritis. *Arthritis Rheum.* 1984;27(1):117–118. doi:10.1002/art.1780270124. PMID: 6362675.
- Faure GC, Bene MC, Thomas P, Tamsier JN. An open trial of thymulin (FTS-Zn) in rheumatoid arthritis patients: sequential clinical and immunological follow-up. *Clin Exp Rheumatol.* 1987;5(3):291–293. PMID: 3427844.
- Israel-Biet D, Noel LH, Bach MA, Dardenne M, Bach JF. Marked reduction of DNA antibody production and glomerulopathy in thymulin (FTS-Zn) or cyclosporin A-treated (NZB × NZW) F1 mice. *Clin Exp Immunol.* 1983;54(2):359–365. PMID: 6360436. PMCID: PMC1535908.
- Bach JF, Dardenne M. Thymulin, a zinc-dependent hormone. *Med Oncol Tumor Pharmacother.* 1989;6(1):25–29. doi:10.1007/BF02985220. PMID: 2657247.
- Faure G, Bene MC, Tamsier JN, Gaucher A, Duheille J. Thymulin (FTS-Zn) induced in vitro modulation of T cell subsets markers on lymphocytes from rheumatoid arthritis and systemic

**Thymulin Available for Research Use Only**  
**(250 mcg – 500 mcg 2–3×/week, various patterns)**

lupus erythematosus patients. *Int J Immunopharmacol*. 1984;6(4):381–388.  
doi:10.1016/0192-0561(84)90058-4. PMID: 6332789.

- Dardenne M, Savino W, Berrih S, Bach JF. A zinc-dependent epitope on the molecule of thymulin, a thymic hormone. *Proc Natl Acad Sci U S A*. 1985;82(20):7035–7038.  
doi:10.1073/pnas.82.20.7035. PMID: 2413455. PMCID: PMC391304.
- Haddad JJ. Thymulin and zinc ( $Zn^{2+}$ )-mediated inhibition of endotoxin-induced production of proinflammatory cytokines and NF- $\kappa$ B nuclear translocation and activation in the alveolar epithelium: unraveling the molecular immunomodulatory, anti-inflammatory effect of thymulin/ $Zn^{2+}$  in vitro. *Mol Immunol*. 2009;47(2–3):205–214.  
doi:10.1016/j.molimm.2009.09.034. PMID: 19850345.
- Prasad AS, Meftah S, Abdallah J, Kaplan J, Brewer GJ, Bach JF, Dardenne M. Serum thymulin in human zinc deficiency. *J Clin Invest*. 1988;82(4):1202–1210. doi:10.1172/JCI113717. PMID: 3262625. PMCID: PMC442670.
- National Institutes of Health, Office of Dietary Supplements. Zinc: Fact Sheet for Health Professionals. <https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/>. Accessed September 2, 2026.
- Reggiani PC, Morel GR, Cónsole GM, Barbeito CG, Rodriguez SS, Brown OA, Bellini MJ, Pléau JM, Dardenne M, Goya RG. The thymus-neuroendocrine axis: physiology, molecular biology, and therapeutic potential of the thymic peptide thymulin. *Ann N Y Acad Sci*. 2009;1153:98–106. doi:10.1111/j.1749-6632.2008.03964.x. PMID: 19236333. PMCID: PMC2688715.
- Auger G, Blanot D, Magnin M, Gastinel LN, Pléau JM, Dardenne M, Bach JF. Synthesis and biological activity of eight thymulin analogues. *Biol Chem Hoppe Seyler*. 1987;368(5):463–470. doi:10.1515/bchm3.1987.368.1.463. PMID: 3497643.
- Bach JF, Dardenne M, Pleau JM, Rosa J. Biochemical characterisation of a serum thymic factor. *Nature*. 1977;266:55–57. doi:10.1038/266055a0.
- Lunin SM, Khrenov MO, Novoselova TV, Parfenyuk SB, Novoselova EG. Thymulin, a thymic peptide, prevents the overproduction of pro-inflammatory cytokines and heat shock protein Hsp70 in inflammation-bearing mice. *Immunol Invest*. 2008;37(8):858–870.  
doi:10.1080/08820130802447629.
- Zhao XH, Awaya A, Kobayashi H, Ohnuki T, Tokumitsu Y, Nomura Y. Effects of repeated administrations of facteur thymique sérique (FTS) on biochemical changes related to aging in senescence-accelerated mouse (SAM). *Jpn J Pharmacol*. 1990;53(3):311–319.  
doi:10.1254/jjp.53.311.
- Yara S, Kawakami K, Kudeken N, et al. FTS reduces bleomycin-induced cytokine and chemokine production and inhibits pulmonary fibrosis in mice. *Clin Exp Immunol*. 2001;124(1):77–85. doi:10.1046/j.1365-2249.2001.01493.x.



**Thymulin Available for Research Use Only**  
**(250 mcg – 500 mcg 2–3×/week, various patterns)**

- Lunin SM, Khrenov MO, Novoselova TV, et al. Modulation of inflammatory response in mice with severe autoimmune disease by thymic peptide thymulin and an inhibitor of NF-kappaB signalling. *Int Immunopharmacol.* 2015;25(2):260–266. doi:10.1016/j.intimp.2015.01.021.
- da Silva AL, Martini SV, Abreu SC, et al. DNA nanoparticle-mediated thymulin gene therapy prevents airway remodeling in experimental allergic asthma. *J Control Release.* 2014;180:125–133. doi:10.1016/j.jconrel.2014.02.010.
- Kanemaru H, Luong S, Yamamoto Y, Mizukami Y, Ito F. Thymulin restrains age-associated myeloid inflammation and enhances cancer immunotherapy. *Nat Commun.* 2026;17:6534. doi:10.1038/s41467-026-75383-0.