Thymosin Alpha-1

Healthspan: 1.0 mg SC 2× weekly, or 1.6 mg 1× weekly; Acute infection / exposure: 1.0 mg SC daily <2 weeks; Vaccination enhancement 1.6mg SC, 2× weekly for four weeks

Tα1 is a 28-amino-acid endogenous immunoregulatory peptide. [8] Tα1 is investigated for supporting cellular immunity, antiviral responses, and recovery from serious infection; healthspan applications are plausible extrapolations from immune-function studies. [2,4,6,7] Approved uses vary by jurisdiction. [21] Rather than a conventional single-receptor agonist, it influences Toll-like-receptor-associated signaling, dendritic-cell antigen presentation, T-cell and NK-cell responses, and regulatory pathways. [2,9,10]

💉 💉 Probably safe. Well tolerated in clinical studies. Common systemic toxicity appears unlikely at studied intermittent regimens over months. [7,17,24,25]

⭐ ⭐ Likely to produce immunomodulatory effects supporting recovery from infection [6,7,9,12]

⭐ Plausibly effective for supporting immune aspects of healthspan. [4,16,22]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Dose, PK, and Spacing

Human studies show short serum exposure after SC injection. A nine-volunteer study comparing formulations found a serum half-life below three hours and no accumulation after repeated administration over five days. [19] A later review describes single doses of 0.8–6.4 mg and repeated 8- and 16-mg doses twice weekly for four weeks, with approximately dose-proportional exposure and reported tolerability. [23] These higher-dose observations are review-reported PK and tolerability findings. A transient circulating pulse can initiate longer-lasting cellular signaling, so serum clearance alone cannot determine the interval between injections. [2,19,23]

Nonlinear dose-response

Tα1 does not show a simple “more dose = more immune effect” relationship. Human lymphocytes stimulated with phytohemagglutinin showed a bimodal concentration-response for high-affinity IL-2 receptor expression; responses varied between donors and required mitogenic stimulation. [20] Hepatitis B and melanoma dose-comparison findings also do not establish a consistent monotonic clinical benefit. [17,18] Tolerability at higher doses is distinct from efficacy, and neither these trials nor the laboratory curve establishes that smaller, more frequent injections are superior. [17,18,20,23]

Dosage & Patterns

In 316 patients with chronic hepatitis B, 0.8- and 1.6-mg regimens produced broadly similar overall responses over 24 weeks of treatment. The protocol used six injections weekly for two

weeks, then twice weekly for 22 weeks. The active-dose comparison did not establish equivalence; an exploratory advanced-fibrosis analysis favored 1.6 mg. [17] The proposed 0.8-mg two-times-weekly pattern is a smaller pulse than the labeled 1.6-mg twice-weekly hepatitis regimen. [21] Short serum exposure and nonlinear laboratory responses do not identify an optimal clinical schedule. [19,20]

A speculative experimental pattern for immune support during aging and for healthspan is:

- **Tα1 1.0 mg SC two times weekly, for immune support during aging and for healthspan. [17,19,20]**
- **Tα1 1.6 mg SC weekly, for immune support during aging and for healthspan is also discussed. [17,19,20]**

A speculative experimental pattern for acute infection or post-exposure immune support is:

- **Tα1 1.0 mg SC daily for less than two weeks, for acute infection or post-exposure. [7,15,17] (Higher doses have been tested in severe sepsis and covid-19.)**

Beginning early after exposure or symptom onset is a mechanistic extrapolation from the importance of early pathogen recognition. The proposed dose and timing combine evidence from different disease settings rather than reproducing an acute-infection trial. [9,11,17] ETASS used 1.6 mg SC twice daily for five days, then daily for two days; TESTS used 1.6 mg SC every 12 hours for seven days. [7,15] These severe-sepsis regimens do not validate lower-dose prophylaxis or self-treatment of minor infection, and TESTS did not show an overall survival benefit. [7]

Thes dose tested for enhancing vaccinations in older adults is

- **Tα1 1.6mg SC, twice weekly for four weeks, beginning on the day of vaccination.**

In randomized, placebo-controlled trials involving older adults, thymosin α1 (Tα1) enhanced antibody responses to influenza vaccination. It produced a fourfold antibody increase in 69% of participants versus 52% receiving placebo. The benefit was concentrated in participants aged 77 and older. A subsequent randomized study found that four weeks of Tα1 produced significantly greater antibody responses than two weeks or placebo, with reported influenza incidence of 6% versus 19% in the four-week Tα1 and placebo groups, respectively. These results support Tα1 as a potential vaccine adjuvant in older adults, although they do not establish efficacy with current vaccines. [4,16]

Benefits

- **Tα1 likely influences coordination between innate and adaptive immune responses** through antigen presentation and T-cell signaling; this is a mechanistic inference supported by cellular and animal experiments. [9,10,12]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- **Tα1 could plausibly support antiviral immune responses through pathogen recognition, antigen presentation, and T-cell function**. Direct experimental antiviral protection includes murine cytomegalovirus models. [11]
- **Tα1 likely supports selected aspects of T-cell activation and signaling**. Human CD8-cell experiments showed enhanced activation and reduced exhaustion-marker expression; these are laboratory endpoints. [12,20] In a small HIV trial, thymic-output markers increased without a significant between-group CD4-count improvement. [13]
- **Tα1 could plausibly help regulate excessive inflammatory signaling through dendritic-cell tryptophan metabolism, IL-10, and regulatory T-cell pathways**. [10]
- **Tα1 could plausibly support recovery after illness** by helping coordinate immune responses and regulate excessive inflammatory signaling. Clinical sepsis results are mixed, with a negative large phase III trial limiting a general recovery claim. [6,7,15]
- **Tα1 could plausibly offer indirect mood or cognitive support** through its immunomodulatory effects when inflammatory signaling contributes to symptoms. This is a mechanistic extrapolation from immune-regulatory pathways. [10]
- **When prolonged stress contributes to impaired immune function, Tα1 could plausibly provide support through antigen presentation and T-cell responses**. Experimental resistance to glucocorticoid-associated thymocyte apoptosis contributes to this mechanistic rationale. [9,25]

Potential applications

- antiviral immune responses. [11,17,21]
- immune aspects of aging. [4,16,22]
- Conditions involving impaired T-cell responses. [13,14,22]

Cautions

- Pregnancy and breastfeeding: human safety is insufficiently characterized. The Indonesian product information contraindicates use during pregnancy and advises against use during breastfeeding. [25]
- Combining Tα1 with cancer immunotherapies or other immunomodulators: combinations could alter immune responses and requires the treating specialist's assessment. Experience with chemotherapy combinations does not establish safety for every immunotherapy combination. [18,25]
- Autoimmune conditions: Tα1 could plausibly alter disease activity through its immunomodulatory effects. Regulatory mechanisms provide a possible benefit rationale, but the direction of the response may depend on the condition. [2,10]
- Deliberate immunosuppression, including organ transplantation, is a contraindication unless a specialist determines that the potential benefits clearly outweigh the risks. [21]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Side Effects

- Possible fatigue or malaise. Malaise occurred in hepatitis B studies, with underlying illness complicating attribution to Tα1. [17]

Duration and Diminished Response

Because multiple immune pathways are involved, conventional single-receptor downregulation is an incomplete model, but it does not rule out adaptation or diminishing response. [2,23]

Biology of Tα1 as an Adjunct Therapy for Serious Infection

Sepsis can combine excessive inflammatory signaling with impaired antigen presentation and lymphocyte dysfunction. Tα1 may help rebalance these processes through dendritic cells, monocytes, and regulatory signaling. [2,9,10,15] ETASS found improved monocyte HLA-DR and a mortality signal whose significance depended on the analysis: 28-day mortality was 26.0% versus 35.0%. [15] In the larger double-blind TESTS trial, mortality was 23.4% versus 24.1% and immune-marker differences were not significant. [7] Earlier favorable syntheses therefore need to be read alongside the negative trial and the nonsignificant high-quality-trial subgroup in the newer meta-analysis. [3,6] A benefit in selected immune phenotypes remains plausible, rather than an established effect across sepsis. [6,7]

Biology of Tα1 as an Adjunct Therapy for Minor Infection

With an ordinary viral infection, the proposed role is to help antiviral defenses work effectively. Tα1 can influence dendritic-cell/Toll-like-receptor signaling, antigen presentation, interferon-related responses, NK cells, and T-cell responses. [2,9,11] A healthy young immune system may gain little from additional modulation; older age or impaired cellular immunity provides a plausible rationale for greater benefit. [4,16] A short pattern beginning early after exposure or symptoms is a mechanistic hypothesis based on early pathogen sensing. [11]

Biology of Tα1 for Healthspan

Aging involves progressive thymic involution, reduced production of naïve T cells, diminished immune surveillance, and chronic low-grade inflammation (*inflammaging*). Tα1 may partially counter these changes by supporting T-cell survival and differentiation, enhancing dendritic-cell and macrophage function, and regulating inflammatory pathways involving IDO, IL-10, and regulatory T cells. [2,10,26] Experimental evidence also suggests that Tα1 can moderate excessive inflammatory signaling while preserving useful antimicrobial immunity. [9,10,26] Improved resistance to infection, maintenance of immune surveillance, and reduced chronic inflammation provide plausible pathways through which Tα1 could support healthspan. However, these remain mechanistic extrapolations. [23,26]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Biology of Tα1 for Vaccination Enhancement

Vaccination depends on effective antigen presentation, T-cell activation, and T-cell help for B-cell antibody production, processes that can become less effective with age. Tα1 may improve these responses by modulating dendritic-cell Toll-like-receptor signaling, enhancing interferon-related responses, and supporting T-cell function and antibody production. [2,4,9,11,26] The proposed benefit is partial restoration of age-impaired vaccine responsiveness rather than indiscriminate immune stimulation. The clinical enhancement of influenza vaccine responses is supported by human trials, while the specific cellular mechanisms remain partly extrapolated from experimental studies. [2,4,10,16]

Biology of Tα1 as an Adjunct Therapy for Autoimmune Disease

Autoimmune disease involves impaired discrimination between self and non-self, rather than simply an immune response that is "too strong." Tα1 can influence dendritic cells toward tolerogenic behavior. Experimental work links TLR9 and type I interferon signaling to indoleamine 2,3-dioxygenase (IDO), tryptophan catabolism, IL-10, and regulatory T-cell generation. [10] Regulatory T cells provide braking signals that can restrain inappropriate immune attack. Tα1 could therefore theoretically support self-tolerance while preserving useful antimicrobial immunity. This is a translational hypothesis from experimental tolerance and antifungal models. [9,10]

The relevant therapeutic aim would be context-dependent immune regulation rather than indiscriminate stimulation; autoimmune diagnoses and concomitant immunomodulators require separate assessment. [2,10,25]

Biological Mechanism

Tα1 is a thymic-derived immunoregulatory peptide whose activity involves peripheral immune cells and multiple signaling systems rather than one established conventional receptor target. [2,8,23] Experimental studies implicate TLR2/TLR9-associated signaling and downstream MyD88, p38 MAPK, NF-κB, and interferon pathways. Dendritic-cell maturation and antigen presentation can promote Th1 antimicrobial responses; TLR9/MyD88/IRF7 signaling contributes to murine antiviral responses. [9,11] IDO-dependent regulatory pathways can simultaneously support IL-10 and regulatory T cells. [10] Human lymphocyte experiments support changes in IL-2 receptor expression, while recent CD8-cell work reports activation and exhaustion-marker changes. [12,20] Together, these findings support a context-dependent immune modulator, with the balance of effector and regulatory activity depending on the biological setting. [2,9–12]

Conclusions

Tα1 has clinical and mechanistic evidence for immunomodulation, but clinical benefit depends on the disease and endpoint; the largest sepsis trial did not demonstrate reduced mortality.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[2,6,7] Short serum exposure and non-monotonic laboratory responses provide a rationale for investigating repeated pulses, without identifying an optimal schedule. [19,20] The proposed 1.0 mg SC two-times-weekly healthspan pattern and 1.0 mg SC daily short-course infection pattern remain experimental extrapolations from separate evidence streams. [7,15,17,21] Clinical experience supports generally favorable tolerability over studied courses, with liver, immune, reproductive, and long-latency risks assessed separately above rather than reduced to a single safety label. [17,24,25]

References

[1] Garaci E, Pica F, Rasi G, Favalli C. Thymosin alpha 1 in the treatment of cancer: from basic research to clinical application. *Int J Immunopharmacol*. 2000;22(12):1067-1076. doi:10.1016/s0192-0561(00)00075-8

[2] Romani L, Bistoni F, Montagnoli C, et al. Thymosin alpha1: an endogenous regulator of inflammation, immunity, and tolerance. *Ann N Y Acad Sci*. 2007;1112:326-338. doi:10.1196/annals.1415.002

[3] Liu F, Wang HM, Wang T, Zhang YM, Zhu X. The efficacy of thymosin α1 as immunomodulatory treatment for sepsis: a systematic review of randomized controlled trials. *BMC Infect Dis*. 2016;16:488. doi:10.1186/s12879-016-1823-5

[4] Ershler WB, Gravenstein S, Geloo ZS. Thymosin alpha 1 as an adjunct to influenza vaccination in the elderly: rationale and trial summaries. *Ann N Y Acad Sci*. 2007;1112:375-384. doi:10.1196/annals.1415.050

[5] Costantini C, Bellet MM, Pariano M, et al. A Reappraisal of Thymosin Alpha1 in Cancer Therapy. *Front Oncol*. 2019;9:873. doi:10.3389/fonc.2019.00873

[6] Gu B, Zhou Y, Nie Y, et al. Efficacy of thymosin α1 for sepsis: a systematic review and meta-analysis of randomized controlled trials. *Front Cell Infect Microbiol*. 2025;15:1673959. doi:10.3389/fcimb.2025.1673959

[7] Wu J, Pei F, Zhou L, et al. The efficacy and safety of thymosin α1 for sepsis (TESTS): multicentre, double blinded, randomised, placebo controlled, phase 3 trial. *BMJ*. 2025;388:e082583. doi:10.1136/bmj-2024-082583 Correction: *BMJ*. 2025;389:r1098. doi:10.1136/bmj.r1098

[8] Goldstein AL, Low TL, McAdoo M, et al. Thymosin alpha1: isolation and sequence analysis of an immunologically active thymic polypeptide. *Proc Natl Acad Sci U S A*. 1977;74(2):725-729. doi:10.1073/pnas.74.2.725

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[9] Romani L, Bistoni F, Gaziano R, et al. Thymosin alpha 1 activates dendritic cells for antifungal Th1 resistance through toll-like receptor signaling. *Blood*. 2004;103(11):4232-4239. doi:10.1182/blood-2003-11-4036

[10] Romani L, Bistoni F, Perruccio K, et al. Thymosin alpha1 activates dendritic cell tryptophan catabolism and establishes a regulatory environment for balance of inflammation and tolerance. *Blood*. 2006;108(7):2265-2274. doi:10.1182/blood-2006-02-004762

[11] Bozza S, Gaziano R, Bonifazi P, et al. Thymosin alpha1 activates the TLR9/MyD88/IRF7-dependent murine cytomegalovirus sensing for induction of anti-viral responses in vivo. *Int Immunol*. 2007;19(11):1261-1270. doi:10.1093/intimm/dxm097

[12] Mishra S, Telang G, Sureshbabu A, et al. Thymosin α1 Augments $CD8^+$ T-Cell Activation and Reverses Exhaustion In Vitro. *Asian Pac J Cancer Prev*. 2026;27(6):2089-2096. doi:10.31557/apjcp.2026.27.6.2089

[13] Chadwick D, Pido-Lopez J, Pires A, et al. A pilot study of the safety and efficacy of thymosin alpha 1 in augmenting immune reconstitution in HIV-infected patients with low CD4 counts taking highly active antiretroviral therapy. *Clin Exp Immunol*. 2003;134(3):477-481. doi:10.1111/j.1365-2249.2003.02331.x

[14] Garaci E, Milanese G, Vella S, et al. A randomized controlled study for the evaluation of the activity of a triple combination of zidovudine, thymosin-α1 and interferon-α in HIV-infected individuals with CD4 counts between 200 and 500 cells/mm^3. *Antivir Ther*. 1998;3(2):103–111. doi:10.1177/135965359800300203

[15] Wu J, Zhou L, Liu J, et al. The efficacy of thymosin alpha 1 for severe sepsis (ETASS): a multicenter, single-blind, randomized and controlled trial. *Crit Care*. 2013;17(1):R8. doi:10.1186/cc11932

[16] Gravenstein S, Duthie EH, Miller BA, et al. Augmentation of influenza antibody response in elderly men by thymosin alpha one. A double-blind placebo-controlled clinical study. *J Am Geriatr Soc*. 1989;37(1):1-8. doi:10.1111/j.1532-5415.1989.tb01561.x

[17] Iino S, Toyota J, Kumada H, et al. The efficacy and safety of thymosin alpha-1 in Japanese patients with chronic hepatitis B; results from a randomized clinical trial. *J Viral Hepat*. 2005;12(3):300-306. doi:10.1111/j.1365-2893.2005.00633.x

[18] Maio M, Mackiewicz A, Testori A, et al. Large randomized study of thymosin alpha 1, interferon alfa, or both in combination with dacarbazine in patients with metastatic melanoma. *J Clin Oncol*. 2010;28(10):1780-1787. doi:10.1200/jco.2009.25.5208

[19] Rost KL, Wierich W, Masayuki F, Tuthill CW, Horwitz DL, Herrmann WM. Pharmacokinetics of thymosin alpha1 after subcutaneous injection of three different formulations in healthy volunteers. *Int J Clin Pharmacol Ther*. 1999;37(1):51-57. PubMed PMID 10027483

[20] Leichtling KD, Serrate SA, Sztein MB. Thymosin alpha 1 modulates the expression of high affinity interleukin-2 receptors on normal human lymphocytes. *Int J Immunopharmacol*. 1990;12(1):19-29. doi:10.1016/0192-0561(90)90064-t

[21] National Drug Formulary. ZADAXIN INJECTION [SIN07483P]. Updated October 9, 2026. Accessed November 4, 2026. Product information

[22] Chen C, Wang J, Xun J, et al. Role of thymosin α1 in restoring immune response in immunological nonresponders living with HIV. *BMC Infect Dis*. 2024;24(1):97. doi:10.1186/s12879-024-08985-y

[23] Garaci E, Paci M, Matteucci C, Costantini C, Puccetti P, Romani L. Phenotypic drug discovery: a case for thymosin alpha-1. *Front Med (Lausanne)*. 2024;11:1388959. doi:10.3389/fmed.2024.1388959

[24] Wu X, Shi Y, Zhou J, et al. Combination of entecavir with thymosin alpha-1 in HBV-related compensated cirrhosis: a prospective multicenter randomized open-label study. *Expert Opin Biol Ther*. 2018;18(suppl 1):61–69. doi:10.1080/14712598.2018.1451511

[25] SciClone Pharmaceuticals. Zadaxin thymosin alfa 1 injection (thymalfasin). Product information approved by BPOM March 4, 2026. Accessed November 4, 2026. Product information

[26] Simonova MA, Ivanov I, Shoshina NS, et al. Aging and thymosin alpha-1. *Int J Mol Sci*. 2025;26(23):11470. doi:10.3390/ijms262311470

[27] Frasca D, Adorini L, Mancini C, Doria G. Reconstitution of T cell functions in aging mice by thymosin alpha 1. *Immunopharmacology*. 1986;11(3):155-163. doi:10.1016/0162-3109(86)90017-2

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076