KLOW Blend (GHK-CU / BPC-157 / TB-500(TB4) / KPV)

1.6 mg daily for skin and hair; 3.2 mg daily for tissue repair

KLOW is an experimental blend proposed to support skin quality, wound and soft-tissue repair, mucosal integrity, and inflammatory regulation through plausible complementary actions of its components. [1–3,8,13,32–38] The common formulation discussed here contains GHK-Cu, BPC-157, full-length thymosin β4 (TB4), and KPV in a 5:1:1:1 mass ratio. GHK-Cu affects extracellular-matrix production in cultured fibroblasts and experimental wounds. [2,3] BPC-157 has preclinical tendon-repair and vascular-signaling findings. [5,13,31] TB4 regulates actin dynamics and supports migration and repair in specific experimental systems. [8,9,17] KPV reduces inflammation in several mouse models and inhibits inflammatory signaling in human cell systems, providing a plausible mucosal-repair rationale for the blend. [32–38] Synergy and the proposed human SC applications remain translational inferences.

💉 💉 Probably Safe, human tolerability findings concern individual components.

⭐ Plausible human benefits for selected repair and inflammatory endpoints, inferred from component research. [1–3,8,13,32–38]

C Angiogenic – caution for subjects at elevated risk for cancers, Do NOT use with active cancer

Synthetic TB4 was tolerated in a study including 14-day IV administration, with additional observations from eye and wound treatments. [18–21] A BPC-157 IV pilot reported no adverse effects or changes in measured laboratory markers in two previously exposed adults through day 3. [40] Topical GHK-Cu findings and KPV cell and animal studies provide indirect context for injected use. [14,32–39]

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

Individual components improved repair or inflammatory outcomes in specific animal models. GHK-Cu has matrix-remodeling findings and mixed topical human cosmetic findings; the post-laser study improved satisfaction but not objective between-group skin outcomes. [1–3,14] TB4 stimulated hair growth in mice. [22] BPC-157 and TB4 improved selected tendon and wound outcomes. [5,8,13] KPV reduced intestinal inflammation in different mouse models. [32,33] These findings support plausible human applications, with dose, delivery, tissue context, and combination effects considered separately. Synergy within KLOW remains plausible.

The names TB4, Tβ4, and TB-500 are used inconsistently in the research-use-only market. Full-length TB4 and the acetylated 17–23 fragment, Ac-LKKTETQ, are different molecules; a vendor name alone does not establish a sample's identity. [6,29,43] This research survey applies to

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

KLOW Blend (GHK-CU / BPC-157 / TB-500(TB4) / KPV)

1.6 mg daily for skin and hair; 3.2 mg daily for tissue repair

KLOW containing full-length, 43-amino-acid thymosin β4. Throughout this document, "TB-500 (TB4)" refers to that full-length peptide.

Dosage & Patterns

In the 50/10/10/10 formulation, GHK-Cu, BPC-157, full-length TB4, and KPV occur in a fixed 5:1:1:1 mass ratio. The following speculative, experimental daily patterns are proposed for SC use:

- **1.6 mg daily KLOW for skin and hair**
  - ~ 1 mg GHK-Cu / daily
  - ~ 200 mcg BPC-157 / daily
  - ~ 200 mcg TB-500 (TB4) / daily
  - ~ 200 mcg KPV / daily
- **3.2 mg daily KLOW for tissue repair**
  - ~ 2 mg GHK-Cu / daily
  - ~ 400 mcg BPC-157 / daily
  - ~ 400 mcg TB-500 (TB4) / daily
  - ~ 400 mcg KPV / daily

The lower pattern is discussed for skin, hair, and healthspan goals, and the higher pattern for injury and tissue-repair goals. These are experimental patterns rather than established minimum-effective or maximum-safe doses.

Benefits

The proposed benefits below draw on component experiments.

- Smoother skin, improved hydration, elasticity, and tone are plausible cosmetic goals. [1,14]
- Plausible human wound-repair support from TB-500 (TB4). [8]
- Faster recovery from soft-tissue strains is plausible through effects on cell migration, vascular repair, and remodeling. [8,11,17]
- Possible reduction in joint discomfort. [41] Improvements in stiffness at other joints remain speculative.
- Plausible support for recovery from minor soft-tissue injuries and overuse irritation through effects on vascular repair and extracellular matrix remodeling. [3,8,11,31]
- Calmer old injury sites and fewer perceived flare-ups are anecdotal possibilities; the proposed repair rationale draws on component wound and tendon findings. [8,13]
- Human tendon healing and improved tendon or ligament resilience during training are plausible. [5,13]
- Improved human microvascular function, nutrient delivery, and tissue oxygenation under mechanical load are plausible extrapolations. [9,11]
- More even texture, fewer fine lines, and improved firmness over several weeks are plausible cosmetic goals. [1,14]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- Generalized tissue comfort, fewer aches, and smoother movement are anecdotal possibilities; mechanisms involving repair and inflammatory regulation remain plausible. [8,13,32–34]
- Intestinal and broader anti-inflammatory benefits from SC KLOW are plausible. [32–34]
- Plausible support for intestinal epithelial and mucosal-barrier integrity through reduced inflammatory signaling. [32, 36,38]

Proposed Applications

- Chronic soft-tissue irritation — plausible repair rationale. [5,8,13]
- Older injuries that flare under load — anecdotal goal with a plausible repair rationale. [8,13]
- High training volume — speculative recovery application. [27,28]
- Age-related changes in skin quality — plausible cosmetic application. [1–3,14]
- Demanding physical jobs or lifestyles — speculative tissue-resilience application. [5,8,13]
- Recovery from minor strains or overuse — plausible repair application. [5,8,13]
- Low-grade systemic inflammation — plausible extrapolation from specific cell and animal inflammatory models. [32–38,46]

Cautions

- **Copper metabolism disorders:** Impaired copper handling, particularly Wilson disease, warrants caution with a copper-containing compound. [45]
- **Liver disease:** TB4's fibrosis biology is context-dependent. Exogenous TB4 reduced chemically induced liver injury and fibrosis in rodents, whereas deleting endogenous TB4 in activated hepatic stellate cells also reduced fibrosis; restoring its expression reversed that benefit. These different interventions should not be treated as equivalent. [23,24]
- **Active cancer or elevated cancer risk:** Experimental tumor-cell studies justify precaution; the direction and magnitude of clinical risk depend on tumor biology and exposure. [25,26]

Drivers of Diminished Response and Mitigation Strategies

These components act through heterogeneous mechanisms rather than a receptor pathway. Downstream adaptation and a benefit from cycling remain mechanistic or speculative proposals.

KLOW and Plausible Healthspan Benefits

The plausible healthspan rationale for KLOW combines four complementary biological programs: GHK-Cu matrix and gene-expression effects, TB4 cytoskeletal and vascular-repair pathways, BPC-157 gastrointestinal and endothelial protection, and KPV inflammatory regulation. [1–5,8,17,31–38] These mechanisms may support tissue integrity, repair capacity, and inflammatory balance during aging. Broader human healthspan benefit from the combination remains speculative.

GHK-related gene-expression studies suggest effects on repair, inflammatory, and antioxidant programs in cultured human cell systems. [30,47] GHK-Cu directly increased collagen synthesis

and changed glycosaminoglycan and proteoglycan production in experimental systems. [2,3] Supporting collagen, elastin, glycosaminoglycan, and metalloprotein regulation is a plausible route to maintaining connective-tissue resilience during aging. [1–3]

TB-500 (TB4) contributes actin regulation, cell migration, and vascular-repair mechanisms to the healthspan rationale. [9,11,17] Wound-healing benefits were observed in specific animal models, with other results mixed. [8,27] Complementarity with GHK-Cu's matrix effects remains plausible. [2,3]

BPC-157's gastrointestinal protection and nitric-oxide-related effects are described in preclinical research. [4,48] Increased VEGFR2 signaling and improved perfusion were observed in animal model. [31] Intestinal-barrier preservation and vascular repair are biologically plausible, as are lower chronic inflammatory burden, improved nutrient delivery, and improved tissue resilience. Broad human healthspan benefit remains speculative. [4,31,48]

KPV contributes a plausible inflammatory-regulation and mucosal-repair rationale. Reduced intestinal inflammation and neutrophil recruitment have been observed in mouse models. [32–34] Human epithelial-cell experiments support mechanisms involving inflammatory signaling. [36–38]

Synergy among the four peptides is mechanistically plausible, providing a rationale for tissue resilience and inflammatory regulation, bit its contribution to human healthspan remains speculative. [1–3,8,13,31–38]

KLOW and Cancer Risk

The cancer concern involves possible support of an existing malignancy through migration and angiogenesis. Increasing TB4 expression within melanoma cells promoted tumor growth and metastasis in mice. [25] GHK-Cu combined with ascorbate reduced tumor growth in a mouse sarcoma experiment. [30] KPV has some positive and some mixed effects on tumors. [35]

TB4 regulates actin dynamics, cell migration, and repair-associated angiogenesis. [9,11,17] Its increased expression promoted vascularization and metastatic behavior in a mouse melanoma model, whereas overexpression in myeloma cells reduced proliferation and migration and prolonged survival in a mouse model. Lower expression was also associated with poorer outcomes in a human myeloma cohort. [25,26] These findings support context-specific precaution rather than a uniformly tumor-promoting or tumor-protective classification.

Experimental Exposure Alternatives

Three proposed alternatives change component exposure; their comparative cancer risk is speculative:

- GHK-Cu with KPV retains plausible matrix-repair and inflammatory-regulation rationales while removing TB4 and BPC-157 exposure. [2,3,32–38]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

- GHK-Cu with KPV and BPC-157 retains plausible soft-tissue-repair and inflammatory-regulation rationales while omitting TB4. Comparative cancer risk remains speculative. [2,3,13,25,26,32–38]
- Alternating vials of KLOW and GHK-Cu (possibly with KPV and/or BPC-157) is a speculative approach to reducing continuous exposure to TB-500 (TB4).

Zinc Supplementation

SC KLOW introduces copper directly into systemic circulation, potentially altering copper–zinc homeostasis. Intracellular copper binds strongly to metallothioneins, cysteine-rich proteins involved in metal sequestration and regulation. Copper has a higher affinity for metallothionein than zinc and can displace bound zinc, potentially altering its intracellular distribution and availability for zinc-dependent enzymes, transcription factors, and signaling pathways. Sustained copper exposure could therefore produce functional zinc insufficiency through altered metal sequestration. Unlike the well-established reduction in copper absorption caused by excessive oral zinc, clinically significant zinc depletion from subcutaneous GHK-Cu remains a mechanistically plausible but unproven effect in humans.

Adult zinc requirements are 8 mg daily for nonpregnant, nonlactating women and 11 mg daily for men; pregnancy and lactation have different requirements. The adult tolerable upper intake level is 40 mg daily from all sources. Sustained excess zinc can impair copper absorption and contribute to hematological, neurological, or immune problems.

Biological Mechanism

GHK-Cu is a copper-binding tripeptide complex with experimental matrix-remodeling actions. Copper is a cofactor in connective-tissue synthesis, antioxidant defense, and mitochondrial energy metabolism; enhancement of those functions after KLOW administration is a mechanistic hypothesis. [1,45] GHK-Cu increased collagen synthesis in fibroblast cultures and altered collagen, glycosaminoglycan, decorin, and biglycan responses in rat wounds and fibroblasts. [2,3] In a mouse inflammatory lung-injury model, it reduced NF-κB-associated signaling and inflammatory mediators, including TNF-α and IL-6. [46] Reviews also describe elastin, cell-migration, and capillary-growth effects. [1] The frequently cited “over 4,000 genes” finding is an estimate from GHK-treated cultured-cell expression profiles, not a demonstrated whole-body regenerative program after injection. [30,47] These observations support a plausible remodeling rationale while keeping copper biology, cell responses, and clinical outcomes distinct.

BPC-157 is a 15-amino-acid experimental peptide with cytoprotective and repair-related findings in preclinical systems. [4,5,13] In rat tendon-derived cultures, it promoted outgrowth, migration, spreading, and survival under oxidative stress, associated with FAK–paxillin signaling; it did not directly increase proliferation in that experiment. [5] In human endothelial cells it increased VEGFR2 expression and internalization and activated VEGFR2–Akt–eNOS signaling, without

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

increasing VEGF-A expression. Rat ischemic-limb experiments showed improved perfusion and vessel formation. [31] Gastrointestinal and inflammatory research discusses barrier preservation, cytokine modulation, and interactions with nitric oxide, dopamine, and serotonin systems. [4,48] FGF-2-related repair signaling and coordinated growth-factor effects remain mechanistic proposals. [4,31] These pathways plausibly connect vascular restoration, inflammation control, and matrix remodeling to tissue recovery. [5,13,31]

Full-length TB4 is a 43-amino-acid peptide that binds and sequesters globular actin, regulating the monomer pool available for filament assembly. [6,17] TB4 improved wound repair and cardiac injury outcomes in particular animal models. [8,9] Corneal-cell studies showed reduced TNF-α-induced NF-κB activation and increased laminin-5 production through a pathway distinct from TGF-β signaling. [7,12] Its repair-related effects include migration and local vascular responses, with outcome and tissue context determining the interpretation. [10,11,23,24,27,28]

KPV (Lys-Pro-Val) corresponds to the C-terminal tripeptide sequence of α-melanocyte-stimulating hormone. PepT1-mediated uptake in intestinal epithelial and immune-cell systems supports inhibition of NF-κB and MAPK signaling and reduced cytokine secretion. [32] Human bronchial epithelial-cell experiments support inhibition of p65RelA nuclear import, a mechanism distinct from MC3R activation by γ-MSH. [37] Several anti-inflammatory effects are at least partly independent of melanocortin receptors; calcium responses in other cell systems make universal receptor independence too strong a conclusion. [33,34,39] KPV reduced neutrophil recruitment in mouse inflammatory models. [33,34] A shift toward repair-associated immune activity and improved mucosal integrity is a plausible mechanistic extension to KPV in KLOW. [32,36,38]

References

[1] Pickart L, Margolina A. Regenerative and Protective Actions of the GHK-Cu Peptide in the Light of the New Gene Data. *Int J Mol Sci*. 2018;19(7):1987. doi:10.3390/ijms19071987

[2] Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. *FEBS Lett*. 1988;238(2):343–346. doi:10.1016/0014-5793(88)80509-x

[3] Siméon A, Wegrowski Y, Bontemps Y, Maquart FX. Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu(2+). *J Invest Dermatol*. 2000;115(6):962–968. doi:10.1046/j.1523-1747.2000.00166.x

[4] Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157: novel therapy in gastrointestinal tract. *Curr Pharm Des*. 2011;17(16):1612–1632. doi:10.2174/138161211796196954

[5] Chang CH, Tsai WC, Lin MS, Hsu YH, Pang JH. The promoting effect of pentadecapeptide BPC 157 on tendon healing involves tendon outgrowth, cell survival, and cell migration. *J Appl Physiol (1985)*. 2011;110(3):774–780. doi:10.1152/japplphysiol.00945.2010

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[6] Goldstein AL, Hannappel E, Sosne G, Kleinman HK. Thymosin β4: a multi-functional regenerative peptide. Basic properties and clinical applications. *Expert Opin Biol Ther*. 2012;12(1):37–51. doi:10.1517/14712598.2012.634793

[7] Sosne G, Qiu P, Christopherson PL, Wheater MK. Thymosin beta 4 suppression of corneal NFkappaB: a potential anti-inflammatory pathway. *Exp Eye Res*. 2007;84(4):663–669. doi:10.1016/j.exer.2006.12.004

[8] Malinda KM, Sidhu GS, Mani H, et al. Thymosin beta4 accelerates wound healing. *J Invest Dermatol*. 1999;113(3):364–368. doi:10.1046/j.1523-1747.1999.00708.x

[9] Bock-Marquette I, Saxena A, White MD, Dimaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. *Nature*. 2004;432(7016):466–472. doi:10.1038/nature03000

[10] Sosne G, Szliter EA, Barrett R, Kernacki KA, Kleinman H, Hazlett LD. Thymosin beta 4 promotes corneal wound healing and decreases inflammation in vivo following alkali injury. *Exp Eye Res*. 2002;74(2):293–299. doi:10.1006/exer.2001.1125

[11] Smart N, Risebro CA, Melville AA, et al. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. *Nature*. 2007;445(7124):177–182. doi:10.1038/nature05383

[12] Sosne G, Xu L, Prach L, et al. Thymosin beta 4 stimulates laminin-5 production independent of TGF-beta. *Exp Cell Res*. 2004;293(1):175–183. doi:10.1016/j.yexcr.2003.09.022

[13] Staresinic M, Sebecic B, Patrlj L, et al. Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth. *J Orthop Res*. 2003;21(6):976–983. doi:10.1016/s0736-0266(03)00110-4

[14] Miller TR, Wagner JD, Baack BR, Eisbach KJ. Effects of topical copper tripeptide complex on CO2 laser-resurfaced skin. *Arch Facial Plast Surg*. 2006;8(4):252–259. doi:10.1001/archfaci.8.4.252

[15] Kang EA, Han YM, An JM, et al. BPC157 as Potential Agent Rescuing from Cancer Cachexia. *Curr Pharm Des*. 2018;24(18):1947–1956. doi:10.2174/1381612824666180614082950

[16] Xing Y, Ye Y, Zuo H, Li Y. Progress on the Function and Application of Thymosin β4. *Front Endocrinol (Lausanne)*. 2021;12:767785. doi:10.3389/fendo.2021.767785

[17] Xue B, Leyrat C, Grimes JM, Robinson RC. Structural basis of thymosin-β4/profilin exchange leading to actin filament polymerization. *Proc Natl Acad Sci U S A*. 2014;111(43):E4596–E4605. doi:10.1073/pnas.1412271111

[18] Ruff D, Crockford D, Girardi G, Zhang Y. A randomized, placebo-controlled, single and multiple dose study of intravenous thymosin beta4 in healthy volunteers. *Ann N Y Acad Sci*. 2010;1194:223–229. doi:10.1111/j.1749-6632.2010.05474.x

[19] Guarnera G, DeRosa A, Camerini R, 8 European sites. The effect of thymosin treatment of venous ulcers. *Ann N Y Acad Sci*. 2010;1194:207–212. doi:10.1111/j.1749-6632.2010.05490.x

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

[20] Sosne G, Ousler GW. Thymosin beta 4 ophthalmic solution for dry eye: a randomized, placebo-controlled, Phase II clinical trial conducted using the controlled adverse environment (CAE™) model. *Clin Ophthalmol*. 2015;9:877–884. doi:10.2147/opth.s80954

[21] Sosne G, Kleinman HK, Springs C, Gross RH, Sung J, Kang S. 0.1% RGN-259 (Thymosin ß4) Ophthalmic Solution Promotes Healing and Improves Comfort in Neurotrophic Keratopathy Patients in a Randomized, Placebo-Controlled, Double-Masked Phase III Clinical Trial. *Int J Mol Sci*. 2023;24(1):554. doi:10.3390/ijms24010554

[22] Philp D, Nguyen M, Scheremeta B, et al. Thymosin beta4 increases hair growth by activation of hair follicle stem cells. *FASEB J*. 2004;18(2):385–387. doi:10.1096/fj.03-0244fje

[23] Li X, Wang L, Chen C. Effects of exogenous thymosin β4 on carbon tetrachloride-induced liver injury and fibrosis. *Sci Rep*. 2017;7(1):5872. doi:10.1038/s41598-017-06318-5

[24] Kim J, Lee C, Han J, et al. Targeted Deletion of Thymosin Beta 4 in Hepatic Stellate Cells Ameliorates Liver Fibrosis in a Transgenic Mouse Model. *Cells*. 2023;12(12):1658. doi:10.3390/cells12121658

[25] Cha HJ, Jeong MJ, Kleinman HK. Role of thymosin beta4 in tumor metastasis and angiogenesis. *J Natl Cancer Inst*. 2003;95(22):1674–1680. doi:10.1093/jnci/djg100

[26] Caers J, Hose D, Kuipers I, et al. Thymosin β4 has tumor suppressive effects and its decreased expression results in poor prognosis and decreased survival in multiple myeloma. *Haematologica*. 2010;95(1):163–167. doi:10.3324/haematol.2009.006411

[27] Gonzalez-Franquesa A, Stocks B, Borg ML, et al. Discovery of thymosin β4 as a human exerkine and growth factor. *Am J Physiol Cell Physiol*. 2021;321(5):C770–C778. doi:10.1152/ajpcell.00263.2021

[28] Tokura Y, Nakayama Y, Fukada S, et al. Muscle injury-induced thymosin β4 acts as a chemoattractant for myoblasts. *J Biochem*. 2011;149(1):43–48. doi:10.1093/jb/mvq115

[29] Janoshik Analytical. TB4/TB-500/TB4(17-23) analysis. Accessed October 22, 2026. Janoshik

[30] Pickart L, Vasquez-Soltero JM, Pickart FD, Majnarich J. GHK, the Human Skin Remodeling Peptide, Induces Anti-Cancer Expression of Numerous Caspase, Growth Regulatory, and DNA Repair Genes. *J Anal Oncol*. 2014;3(2):79–87. doi:10.6000/1927-7229.2014.03.02.2

[31] Hsieh MJ, Liu HT, Wang CN, et al. Therapeutic potential of pro-angiogenic BPC157 is associated with VEGFR2 activation and up-regulation. *J Mol Med (Berl)*. 2017;95(3):323–333. doi:10.1007/s00109-016-1488-y

[32] Dalmasso G, Charrier-Hisamuddin L, Nguyen HT, Yan Y, Sitaraman S, Merlin D. PepT1-mediated tripeptide KPV uptake reduces intestinal inflammation. *Gastroenterology*. 2008;134(1):166–178. doi:10.1053/j.gastro.2007.10.026

[33] Kannengiesser K, Maaser C, Heidemann J, et al. Melanocortin-derived tripeptide KPV has anti-inflammatory potential in murine models of inflammatory bowel disease. *Inflamm Bowel Dis*. 2008;14(3):324–331. doi:10.1002/ibd.20334

[34] Getting SJ, Schiöth HB, Perretti M. Dissection of the anti-inflammatory effect of the core and C-terminal (KPV) alpha-melanocyte-stimulating hormone peptides. *J Pharmacol Exp Ther*. 2003;306(2):631–637. doi:10.1124/jpet.103.051623

[35] Viennois E, Ingersoll SA, Ayyadurai S, et al. Critical role of PepT1 in promoting colitis-associated cancer and therapeutic benefits of the anti-inflammatory PepT1-mediated tripeptide KPV in a murine model. *Cell Mol Gastroenterol Hepatol*. 2016;2(3):340–357. doi:10.1016/j.jcmgh.2016.01.006

[36] Xiao B, Xu Z, Viennois E, et al. Orally Targeted Delivery of Tripeptide KPV via Hyaluronic Acid-Functionalized Nanoparticles Efficiently Alleviates Ulcerative Colitis. *Mol Ther*. 2017;25(7):1628–1640. doi:10.1016/j.ymthe.2016.11.020

[37] Land SC. Inhibition of cellular and systemic inflammation cues in human bronchial epithelial cells by melanocortin-related peptides: mechanism of KPV action and a role for MC3R agonists. *Int J Physiol Pathophysiol Pharmacol*. 2012;4(2):59–73. PubMed

[38] Cheng J, Wu P, Li C, et al. Inflammation-triggered self-immolative conjugates enable oral peptide delivery by overcoming gastrointestinal barriers. *Sci Adv*. 2026;12(3):eaea2989. doi:10.1126/sciadv.aea2989

[39] Elliott RJ, Szabo M, Wagner MJ, Kemp EH, MacNeil S, Haycock JW. alpha-Melanocyte-stimulating hormone, MSH 11-13 KPV and adrenocorticotropic hormone signalling in human keratinocyte cells. *J Invest Dermatol*. 2004;122(4):1010–1019. doi:10.1111/j.0022-202x.2004.22404.x

[40] Lee E, Burgess K. Safety of Intravenous Infusion of BPC157 in Humans: A Pilot Study. *Altern Ther Health Med*. 2025;31(5):20–24. PubMed

[41] Lee E, Padgett B. Intra-Articular Injection of BPC 157 for Multiple Types of Knee Pain. *Altern Ther Health Med*. 2021;27(4):8–13. PubMed

[42] Wang X, Liu L, Qi L, et al. A first-in-human, randomized, double-blind, single- and multiple-dose, phase I study of recombinant human thymosin β4 in healthy Chinese volunteers. *J Cell Mol Med*. 2021;25(17):8222–8228. doi:10.1111/jcmm.16693

[43] US Food and Drug Administration. Certain Bulk Drug Substances for Use in Compounding that May Present Significant Safety Risks. Accessed October 22, 2026. FDA

[44] National Institutes of Health, Office of Dietary Supplements. Zinc: Fact Sheet for Health Professionals. Accessed October 22, 2026. NIH ODS

[45] National Institutes of Health, Office of Dietary Supplements. Copper: Fact Sheet for Health Professionals. Accessed October 22, 2026. NIH ODS

[46] Park JR, Lee H, Kim SI, Yang SR. The tri-peptide GHK-Cu complex ameliorates lipopolysaccharide-induced acute lung injury in mice. *Oncotarget*. 2016;7(36):58405–58417. doi:10.18632/oncotarget.11168

[47] Pickart L, Vasquez-Soltero JM, Margolina A. GHK and DNA: resetting the human genome to health. *Biomed Res Int*. 2014;2014:151479. doi:10.1155/2014/151479

[48] Park JM, Lee HJ, Sikiric P, Hahm KB. BPC 157 Rescued NSAID-cytotoxicity Via Stabilizing Intestinal Permeability and Enhancing Cytoprotection. *Curr Pharm Des*. 2020;26(25):2971–2981. doi:10.2174/1381612826666200523180301

[49] US Food and Drug Administration. KPV (lysine-proline-valine): evaluation for the 503A Bulks List. Pharmacy Compounding Advisory Committee briefing document. May 12, 2026. FDA briefing document