

GHK-Cu Topical Serum

(Once or Twice Daily on Face & Scalp)

GHK-Cu is the copper complex of a naturally occurring human peptide. It is not a classical receptor agonist. By binding and transporting copper and interacting with extracellular-matrix, inflammatory, antioxidant, and repair pathways, it can influence multiple signaling processes involved in skin maintenance and wound repair. It is widely used in cosmetic formulations to improve several measures of photoaged facial skin and support hair growth and thickness.

★ ★ ★ Effective for improving several measures of photoaged facial skin.

★ ★ ★ Effective for increasing hair growth and thickness.

★ ★ ★ Effective to support local wound repair and extracellular-matrix remodeling.

★ ★ Safety is likely based on animal evidence and common use in cosmetics.

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 23-Sep-26. The most recent version of this document is available at <https://pdfs.researcher6076.com/pdfs/ghk-cu-topical.pdf>

Formulation

An ideal formulation would be 2% GHK-Cu raw powder in an aqueous cosmetic base containing 2% total mixed high- and low-molecular-weight hyaluronic acid.

Use Patterns

- **Apply twice daily.** Once-daily applications are a practical lower-frequency pattern.
- **Apply a thin, even layer to clean skin.** Avoid direct contact with the eyes and mucosal surfaces.

Benefits

- **Photoaged skin appearance:** Topical GHK-Cu improves the overall appearance of photoaged facial and periocular skin, including clarity and visible texture, in human studies using GHK-Cu creams.
- **Fine lines and wrinkles:** Topical GHK-Cu reduces facial and periocular lines and wrinkles over approximately 12 weeks in the reported human studies.
- **Skin firmness and density:** Topical GHK-Cu improves measured skin density, thickness, firmness, and laxity in the reported facial studies.

GHK-Cu Topical Serum

(Once or Twice Daily on Face & Scalp)

- **Uneven pigmentation appearance:** Topical GHK-Cu improves mottled pigmentation and visible photodamage in reported human facial studies.
- **Hair growth:** Increases hair count and diameter.
- **Local wound repair:** A 2% GHK-Cu gel supported closure of diabetic lower-extremity ulcers when used with debridement and structured wound care. This evidence applies to the studied medical formulation and setting.
- **Collagen synthesis support:** GHK-Cu likely promotes local collagen synthesis and extracellular-matrix repair based on injected and topical animal evidence, supported by fibroblast and ex vivo skin findings.
- **Matrix remodeling support:** GHK-Cu likely supports regulated remodeling of collagen, glycosaminoglycans, and other matrix components during local repair.
- **Scar appearance support:** Improvement in scar texture and appearance is plausible through effects on collagen production, matrix remodeling, inflammation, and wound repair.
- **Basement membrane support:** GHK-Cu plausibly supports the dermal-epidermal junction by increasing collagen IV, based on fibroblast and ex vivo skin research.
- **Inflammatory signal regulation:** Local modulation of excessive inflammatory signaling is plausible through effects on NF-κB-related and cytokine pathways described in preclinical research.
- **Antioxidant defense support:** Local support of antioxidant defenses is plausible through copper-dependent enzymes and GHK-Cu-associated regulation of oxidative-stress pathways.

Cautions

- **Copper allergy**, contact sensitivity, or sensitivity to another ingredient in the finished base warrants caution and patch testing.
- **Active dermatitis, a disrupted skin barrier, or recent cosmetic procedures** can increase irritation and absorption. Introduce the serum only after the skin has recovered or with clinician guidance.
- **Use on chronic ulcers, burns, infected wounds, or other significant injuries** belongs in supervised medical wound care rather than a cosmetic protocol.
- Application directly over a **known or suspected skin cancer** warrants clinical guidance because GHK-Cu participates in repair, angiogenic, and matrix-remodeling pathways.
- **Wilson disease or another disorder of copper handling** warrants clinician guidance before prolonged or large-area use.

Biological Mechanism

GHK binds copper with high affinity and forms GHK-Cu, a redox-active but biologically regulated complex. In skin models, GHK-Cu influences fibroblast and keratinocyte activity, collagen and glycosaminoglycan production, matrix metalloproteinases and their inhibitors, inflammatory

GHK-Cu Topical Serum

(Once or Twice Daily on Face & Scalp)

cytokine signaling, antioxidant defenses, angiogenesis, and recruitment of repair processes. These overlapping effects support regulated tissue remodeling rather than a single linear pathway.

Formulation Chemistry with Hyaluronic Acid

An appropriate serum combines 2% GHK-Cu with 2% total mixed-molecular-weight hyaluronic acid in an aqueous cosmetic base. Hyaluronic acid is a water-binding polymer that increases surface hydration and can improve slip and film formation. Higher-molecular-weight fractions primarily support surface hydration and film properties, while lower-molecular-weight fractions interact more readily with superficial skin compartments.

In fibroblast and ex vivo skin experiments, GHK-Cu combined with hyaluronic acid increased collagen I, IV, and VII expression, with the strongest collagen IV response reported for a 1:9 GHK-Cu to low-molecular-weight hyaluronic-acid ratio. This provides a plausible biological rationale for combining the ingredients.

References

1. Mulder GD, Patt LM, Sanders L, et al. Enhanced healing of ulcers in patients with diabetes by topical treatment with glycyl-L-histidyl-L-lysine copper. Wound Repair Regen. 1994;2(4):259-269. DOI: [10.1046/j.1524-475X.1994.20406.x](https://doi.org/10.1046/j.1524-475X.1994.20406.x).
2. Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. FEBS Lett. 1988;238(2):343-346. DOI: [10.1016/0014-5793\(88\)80509-X](https://doi.org/10.1016/0014-5793(88)80509-X).
3. Maquart FX, Bellon G, Chaqour B, et al. In vivo stimulation of connective tissue accumulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺ in rat experimental wounds. J Clin Invest. 1993;92(5):2368-2376. DOI: [10.1172/JCI116842](https://doi.org/10.1172/JCI116842).
4. Siméon A, Monier F, Emonard H, et al. Expression and activation of matrix metalloproteinases in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu²⁺. J Invest Dermatol. 2000;115(6):962-968. DOI: [10.1046/j.1523-1747.2000.00166.x](https://doi.org/10.1046/j.1523-1747.2000.00166.x).
5. Lau SJ, Sarkar B. The interaction of copper(II) and glycyl-L-histidyl-L-lysine, a growth-modulating tripeptide from plasma. Biochem J. 1981;199(3):649-656. DOI: [10.1042/bj1990649](https://doi.org/10.1042/bj1990649).
6. Conato C, Kozłowski H, Młynarz P, et al. Copper complexation by glycyl-L-histidyl-L-lysine and related peptides. Biochim Biophys Acta. 2001;1526(2):199-210. DOI: [10.1016/S0304-4165\(01\)00127-1](https://doi.org/10.1016/S0304-4165(01)00127-1).
7. Gorouhi F, Maibach HI. Role of topical peptides in preventing or treating aged skin. Int J Cosmet Sci. 2009;31(5):327-345. DOI: [10.1111/j.1468-2494.2009.00490.x](https://doi.org/10.1111/j.1468-2494.2009.00490.x).

GHK-Cu Topical Serum

(Once or Twice Daily on Face & Scalp)

8. Pickart L, Vasquez-Soltero JM, Margolina A. GHK peptide as a natural modulator of multiple cellular pathways in skin regeneration. Biomed Res Int. 2015;2015:648108. DOI: [10.1155/2015/648108](https://doi.org/10.1155/2015/648108).
9. Pickart L, Margolina A. Regenerative and protective actions of the GHK-Cu peptide in the light of the new gene data. Int J Mol Sci. 2018;19(7):1987. DOI: [10.3390/ijms19071987](https://doi.org/10.3390/ijms19071987).
10. Jiang F, Wu Y, Liu Z, Hong M, Huang Y. Synergy of GHK-Cu and hyaluronic acid on collagen IV upregulation via fibroblast and ex-vivo skin tests. J Cosmet Dermatol. 2023;22(9):2598-2604. DOI: [10.1111/jocd.15763](https://doi.org/10.1111/jocd.15763).
11. Badenhorst T, Svirskis D, Wu Z. Physicochemical characterization of native glycyl-L-histidyl-L-lysine tripeptide for wound healing and anti-aging: a preformulation study for dermal delivery. Pharm Dev Technol. 2016;21(2):152-160. DOI: [10.3109/10837450.2014.979944](https://doi.org/10.3109/10837450.2014.979944).
12. Choi HR, Kang YA, Ryoo SJ, et al. Stem cell recovering effect of copper-free GHK in skin. J Pept Sci. 2012;18(11):685-690. DOI: [10.1002/psc.2455](https://doi.org/10.1002/psc.2455).
13. Bo SL, Nararatwanchai T, Supornpun N, Poomarapan K, Thaploka S. The efficacy of 2% copper peptide (GHK-Cu) serum for eyebrow hypotrichosis: a randomized, double-blind, vehicle-controlled, split-face comparative study. Procedia of Multidisciplinary Research. 2026;4(5). [Full text](#).
14. Leyden JJ, Stevens T, Finkey MB, et al. Skin care benefits of copper peptide containing facial cream. American Academy of Dermatology Annual Meeting. 2002. Conference report summarized in peer-reviewed reviews of topical peptides.
15. Abdulghani AA, Sherr A, Shirin S, et al. Effects of topical creams containing vitamin C, a copper-binding peptide cream and melatonin compared with tretinoin on the ultrastructure of normal skin: a pilot clinical, histologic, and ultrastructural study. Dis Manag Clin Outcomes. 1998;1(4):136-141. DOI: [10.1016/S1088-3371\(98\)00011-4](https://doi.org/10.1016/S1088-3371(98)00011-4).
16. Mazurowska L, Mojski M. Biological activities of selected peptides: skin penetration ability of copper complexes with peptides. J Cosmet Sci. 2008;59(1):59-69.
17. Miller TR, Wagner JD, Baack BR, Eisbach KJ. Effects of topical copper tripeptide complex on CO2 laser-resurfaced skin. Arch Facial Plast Surg. 2006;8(4):252-259. DOI: [10.1001/archfaci.8.4.252](https://doi.org/10.1001/archfaci.8.4.252).
18. Hostynek JJ, Dreher F, Maibach HI. Human skin retention and penetration of a copper tripeptide in vitro as function of skin layer towards anti-inflammatory therapy. Inflamm Res. 2010;59(11):983-988. DOI: [10.1007/s00011-010-0214-4](https://doi.org/10.1007/s00011-010-0214-4).
19. Pavicic T, Gauglitz GG, Lersch P, et al. Efficacy of cream-based novel formulations of hyaluronic acid of different molecular weights in anti-wrinkle treatment. J Drugs Dermatol. 2011;10(9):990-1000. [Full text](#).

GHK-Cu Topical Serum

(Once or Twice Daily on Face & Scalp)

20. Muhammad P, Novianto E, Setyorini M, et al. Effectiveness of topical hyaluronic acid of different molecular weights in xerosis cutis treatment in elderly: a double-blind, randomized controlled trial. Arch Dermatol Res. 2024;316(6):329. DOI: [10.1007/s00403-024-03003-2](https://doi.org/10.1007/s00403-024-03003-2).