GLOW Blend (GHK-Cu / BPC-157 / TB-500 (TB4)) Available for Research Use Only

1.4 mg for cosmetic use or 2.8 mg for tissue repair

Glow is a peptide blend of GHK-Cu, BPC-157, and TB-500 (TB4), typically in the ratio 5:1:1. It is intended to combine GHK-Cu's collagen and repair-signaling with BPC-157 and TB-500 (TB4)'s tissue-repair and anti-inflammatory actions to support skin quality, wound healing, and connective-tissue resilience. Synergies in the combination are plausible. TB-500 (TB4)'s effects on cell migration and angiogenesis warrant caution in people with active cancer, a history of cancer, or elevated cancer risk. [1,8,9,13,25,26]

💉 💉 Probably safe (some human evidence)

⭐ ⭐ Probably effective (strong animal signals and some positive human signals)

C Angiogenic – caution for subjects at elevated risk for cancers, Do NOT use with active cancer

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 12-Oct-26. The most recent version of this document is available at https://pdfs.researcher6076.com/pdfs/glow-blend.pdf

*(The names **TB4**, **Tβ4** and **TB-500** are used inconsistently in gray market and research-use-only community. Vendor labels may use "TB-500" for either the fragment or full-length Tβ4. Labels such as "TB-500 (TB4)" or "TB-500 (frag)" can help indicate the intended identity, but the name alone does not establish what a sample contains.*

Janoshik's reporting terminology helps clarify these distinctions. "TB-500 (TB-4)" identifies full-length, 43-amino-acid Tβ4, while "TB-500 889 (Da)" identifies the acetylated fragment corresponding to residues 17–23, Ac-LKKTETQ. A report labeled only "TB-500" identifies a fragment, with its specific identity given in the comments. [29])

This research survey applies to the GLOW blend using full-length, 43-amino-acid Tβ4 which is common. Throughout the document "TB-500 (TB4) is used to refer to full-length, 43-amino-acid Tβ4.

Short term human safety is informed by the safety of the individual peptides. TB-500 (TB4) has short-term human tolerability data from intravenous studies, with additional observations from local eye and wound treatments. GHK-Cu short-term safety is likely based on animal evidence. BPC-157 has positive, albeit limited human safety signals, along with very widespread community use with few serious adverse events reported. Long-term safety of the injected GLOW blend remains uncertain. [18-21]

GLOW Blend (GHK-Cu / BPC-157 / TB-500 (TB4)) Available for Research Use Only

1.4 mg for cosmetic use or 2.8 mg for tissue repair

GHK-Cu research supports skin repair and extracellular-matrix remodeling, with some topical human findings for cosmetic improvement. Animal studies support likely hair-growth benefits from TB-500 (TB4) through hair-follicle stem-cell activity. These individual findings support GLOW's cosmetic rationale. Synergies in the combination are plausible. [1-3,14,22]

Animal studies support likely healing benefits from individual GLOW components, including TB-500 (TB4)'s effects on wound repair and BPC-157's effects on tendon healing. GHK-Cu research also supports extracellular-matrix production and remodeling. Synergies in the combination are plausible. [2,3,8,13]

Dosage & Patterns

The 50/10/10 GLOW formulation contains GHK-Cu, BPC-157, and TB-500 (TB4) in a fixed 5:1:1 mass ratio. Human studies do not establish the least effective or maximum safe dose. The doses below describe typical clinic and community practices rather than validated GLOW regimens.

- **1.4 mg / daily GLOW, for skin & hair**
  - ~ 1 mg GHK-CU / daily
  - ~ 200 mcg BPC-157 / daily
  - ~ 200 mcg TB-500 (TB4) / daily
- **2.8 mg / daily GLOW, for tissue repair**
  - ~ 2 mg GHK-CU / daily
  - ~ 400 mcg BPC-157 / daily
  - ~ 400 mcg TB-500 (TB4) / daily

With the lower dose for skin and hair repair and regeneration, and the higher dose for more aggressive injury and tissue repair. Some practitioners report using daily doses more than two times larger.

Benefits

The benefits below draw on research involving the individual peptides and reports from GLOW users. TB-500 (TB4) has human tolerability and local-treatment data alongside animal repair studies. GHK-Cu has topical human and preclinical findings, while BPC-157's repair evidence is predominantly preclinical. Synergies in the combination are plausible. Community reports describe user experiences and are identified separately from scientific findings. [1,13,18-21]

- **Smoother, healthier-looking skin**, with improvements in hydration, elasticity, and overall tone.
- **Likely support for wound repair** from TB-500 (TB4), based on animal studies.
- **Faster recovery from soft-tissue strains is plausible** through effects on cell migration, vascular repair, and tissue remodeling. [8,11,17]
- **Reduced joint stiffness and background inflammation**, often noticed in knees, shoulders, and hips as connective-tissue signaling normalizes.

- **Plausible support for recovery from minor soft-tissue injuries and overuse irritation** through effects on vascular repair and extracellular-matrix remodeling. Synergies in the combination are plausible. Relief of delayed-onset muscle soreness remains a community-reported benefit. [3,8,11,31]
- **Calmer old injury sites**, with many users describing less reactivity or "flare-ups" in previously problematic areas.
- **Likely support for tendon healing** from BPC-157, based on animal studies, with plausible complementary support from TB-500 (TB4)'s repair pathways. **Improved tendon and ligament resilience** during training remains plausible rather than demonstrated. [5,8,13]
- **Likely support for vascular repair** from TB-500 (TB4), based on animal injury studies. **Improved microvascular function, nutrient delivery, and tissue oxygenation under mechanical load** are plausible benefits. [9,11]
- **More even skin texture**, including subtle reductions in fine lines and improved firmness over several weeks.
- **Generalized tissue "comfort"**, a broad effect people describe as less tightness, fewer aches, and smoother movement patterns.

Indications

- Chronic soft-tissue irritation
- Older injuries that flare under load
- High training volume
- Age-related changes in skin quality
- Demanding physical jobs or lifestyles
- Recovering from minor strains or overuse
- Low-grade systemic inflammation
- Running GH-axis peptides

Contraindications

- Copper metabolism disorders
- Liver disease
- Active cancer, elevated cancer risk

Reported Side Effects

- Nausea, headache, and fatigue have been reported by users. Their frequency and causal relationship to GLOW are uncertain.

Drivers of Diminished Response and Mitigation Strategies

None of TB-500 (TB4), BPC-157, or GHK-Cu are receptor agonists. Adaptation within repair pathways is plausible but without evidence. [8,17]

Cycling is a speculative approach to limiting continuous exposure.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

GLOW: Plausible Longevity Benefits

The plausibly pro-longevity rationale for the GLOW blend arises from the convergence of **three distinct but complementary biological programs**: GHK-Cu's gene-level regenerative and anti-inflammatory signaling, TB-500 (TB4)'s cytoskeletal and angiogenic repair pathways, and BPC-157's gastrointestinal, endothelial, and microvascular stabilization effects. Each peptide **targets a different conserved pathway in age-related decline**, and together they form a mechanistically coherent strategy for supporting tissue integrity, repair capacity, and inflammatory balance, three domains that strongly influence healthspan.

GHK-Cu contributes a broad gene-reprogramming signal that shifts cellular behavior away from pro-inflammatory and pro-oxidative states toward regeneration, extracellular-matrix rebuilding, and antioxidant defense. Age-related declines in GHK-Cu correlate with reduced collagen synthesis, impaired wound healing, and increased inflammatory tone; restoring this signal may help maintain ECM quality, reduce chronic inflammatory burden, and preserve microcirculatory function. These actions address structural aging at the level of collagen, elastin, glycosaminoglycans, and metalloprotein regulation, supporting the connective-tissue architecture that underpins organ resilience.

TB-500 (TB4) contributes actin regulation, cell migration, and vascular repair to GLOW's proposed healthspan rationale. Animal studies support likely benefits for wound healing and selected forms of tissue repair. Its effects on fibrosis vary by tissue and experimental context, so a general antifibrotic benefit should not be assumed. These repair activities plausibly support tissue resilience during aging. Synergies with GHK-Cu's extracellular-matrix effects are plausible. [8,9,11,17,23,24]

BPC-157 complements both signals by stabilizing the gastrointestinal barrier, protecting microvasculature, and modulating nitric-oxide pathways that influence endothelial tone and perfusion. Because aging is strongly shaped by chronic low-grade inflammation originating from gut permeability, vascular instability, and impaired angiogenic balance, BPC-157's protective effects on the gut–vascular axis may reduce systemic inflammatory load and improve nutrient delivery, oxygenation, and waste clearance. Its anti-ulcer, endothelial-protective, and tendon-repair properties add a third dimension to the blend's regenerative profile, supporting both local and systemic resilience.

Synergies in combining these three peptides are plausible, providing a plausible rationale for supporting tissue resilience and healthspan. [1-3,8,13,31]

GLOW and Cancer Risk

The cancer concern with GLOW involves possible support of an existing malignancy through effects on angiogenesis and cell migration. **TB4 tumor-promotion findings in animal models make this concern more than purely theoretical**, although tumor-cell expression studies are

not equivalent to administering TB-500 (TB4). These findings do not establish that GLOW causes new cancers, and its cancer risk in humans remains unresolved. [25,26]

GHK-Cu has experimental findings relevant to protective cellular responses and cancer-related gene expression. In a mouse sarcoma model, GHK-Cu combined with ascorbic acid reduced tumor growth. This supports a protective effect of that combination in the model studied, rather than establishing that GHK-Cu alone prevents or treats cancer. [30] Although mildly pro-angiogenic, it has **not** been linked to tumor promotion in vivo. But these findings do not establish general cancer protection or show that GHK-Cu offsets potential tumor-related concerns from TB-500 (TB4). [1,30]

BPC-157 has experimental findings suggesting protective effects in selected settings. Experimental work discussed by Kang and colleagues suggests that BPC-157 may counter cancer-associated muscle wasting and related inflammatory changes, but these benefits do not establish suppression of tumor growth or metastasis. [15]

TB-500 (TB4) has direct experimental evidence relevant to cancer risk. Increased TB4 expression promoted tumor vascularization and metastasis in a mouse melanoma model. TB-500 (TB4) is **overexpressed in several cancers** and associated with metastasis and poor prognosis. [16,25] TB-500 (TB4) regulates actin dynamics and supports cell migration and angiogenesis during repair. These processes can also contribute to tumor progression. [17,25] Other studies found tumor-suppressive effects in multiple myeloma. These findings support caution around an existing malignancy and caution around an elevated cancer risk. [25,26]

Conservative Alternatives Around Cancer Risks

For someone with a desire to be cautious, three alternative patterns are available

- **Using GHK-Cu alone** offers a simpler approach to skin repair and extracellular-matrix support while avoiding exposure to TB-500 (TB4) and BPC-157.
- Combine GHK-Cu daily with BPC-157 daily or less frequently provides additional soft tissue support while avoiding the cancer risks of TB-500 (TB4).
- Alternating vials of GLOW and GHK-Cu is a speculative approach to reducing continuous exposure to BPC-157 and TB-500 (TB4).

Consider zinc Supplementation Because of GHK-Cu

People often **consider zinc supplementation alongside GHK-Cu** because zinc helps maintain balanced metal homeostasis in the body. GHK-Cu delivers copper to enzymes involved in repair, and zinc plays a complementary role by supporting antioxidant defenses, stabilizing cellular signaling, improved immune response, and helping regulate metalloproteins that manage copper–zinc balance. When overall zinc status is low, the body may have more difficulty buffering shifts in copper availability, so **maintaining adequate zinc intake can help keep these**

metal-dependent pathways stable while GHK-Cu is being used. The zinc requirement is not large, probably 20 mg daily is sufficient to support the continuous maximum dose of GHK-Cu.

Biological Mechanism

GHK-Cu is described in research as a naturally occurring **copper-binding tripeptide** that functions as a repair and remodeling signal across multiple tissues. Biologically, it binds copper(II) and delivers it to enzymes involved in **collagen cross-linking, antioxidant defense** (such as superoxide dismutase), and mitochondrial energy production, while simultaneously activating gene programs linked to **extracellular-matrix rebuilding, angiogenesis**, and **wound healing**. Physiologically, GHK-Cu reduces inflammatory signaling through suppression of **NF-κB, TNF-α**, and **IL-6**, increases the expression of structural proteins like collagen, elastin, and glycosaminoglycans, and promotes fibroblast and keratinocyte migration during tissue repair. It also improves microcirculation by supporting capillary growth and endothelial function and has been shown to modulate over 4,000 human genes toward a regenerative, anti-inflammatory profile. This combination of copper transport, gene-level repair signaling, and inflammation control is why GHK-Cu appears across research on skin quality, connective-tissue healing, and general tissue regeneration.

BPC-157 is described in research as a **cytoprotective, angiogenic**, and **anti-inflammatory** peptide that operates through several converging biological pathways. It stabilizes the gut-vascular axis by up-regulating **VEGF, FGF-2**, and other growth-repair signals while simultaneously suppressing **TNF-α, IL-6**, and **NF-κB-driven inflammation**. At the cellular level, BPC-157 enhances **fibroblast migration, collagen organization**, and **endothelial repair**, which supports tendon, ligament, and soft-tissue healing. It also interacts with the nitric-oxide system to normalize microvascular tone and protect against oxidative stress. In animal and early mechanistic studies, BPC-157 preserves tight-junction integrity in the gut, accelerates angiogenesis in injured tissue, modulates dopamine and serotonin pathways, and promotes rapid healing across muscle, nerve, and gastrointestinal models. Together, these effects position BPC-157 as a broad-spectrum repair peptide that coordinates inflammation control, vascular restoration, and extracellular-matrix rebuilding.

TB-500 (TB4) is a naturally occurring 43-amino-acid peptide that binds and sequesters globular actin, helping regulate actin availability for cell movement and tissue repair. Animal studies support likely benefits for wound healing, cardiac protection, and selected inflammatory responses. Cell studies also show effects on epithelial migration and inflammatory signaling. These findings support TB-500 (TB4)'s contribution to GLOW's repair rationale.

Synergies in the combination are plausible. [7-11,17]

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Conclusions

GLOW combines peptides with research support for skin repair, extracellular-matrix remodeling, and tissue healing. TB-500 (TB4) contributes direct animal repair evidence and limited human tolerability and local-treatment findings. Synergies in the combination are plausible. Experimental tumor findings support caution around an existing malignancy. Cycling remains speculative and has not been shown to reduce cancer risk. [1,8,13,18-21,25,26]

References

1. Pickart L, Margolina A. Regenerative and protective actions of the GHK-Cu peptide in the light of the new gene data. Int J Mol Sci. 2018;19(7):1987. https://doi.org/10.3390/ijms19071987

2. Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. FEBS Lett. 1988;238(2):343-346. https://doi.org/10.1016/0014-5793(88)80509-X

3. Siméon A, Wegrowski Y, Bontemps Y, Maquart FX. Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu(2+). J Invest Dermatol. 2000;115(6):962-968. https://doi.org/10.1046/j.1523-1747.2000.00166.x

4. Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157: novel therapy in gastrointestinal tract. Curr Pharm Des. 2011;17(16):1612-1632. https://doi.org/10.2174/138161211796196954

5. Chang CH, Tsai WC, Lin MS, Hsu YH, Pang JH. The promoting effect of pentadecapeptide BPC 157 on tendon healing involves tendon outgrowth, cell survival, and cell migration. J Appl Physiol (1985). 2011;110(3):774-780. https://doi.org/10.1152/japplphysiol.00945.2010

6. Goldstein AL, Hannappel E, Sosne G, Kleinman HK. Thymosin β4: a multi-functional regenerative peptide. Basic properties and clinical applications. Expert Opin Biol Ther. 2012;12(1):37-51. https://doi.org/10.1517/14712598.2012.634793

7. Sosne G, Qiu P, Christopherson PL, Wheater MK. Thymosin beta 4 suppression of corneal NFκB: a potential anti-inflammatory pathway. Exp Eye Res. 2007;84(4):663-669. https://doi.org/10.1016/j.exer.2006.12.004

8. Malinda KM, Sidhu GS, Mani H, et al. Thymosin beta4 accelerates wound healing. J Invest Dermatol. 1999;113(3):364-368. https://doi.org/10.1046/j.1523-1747.1999.00708.x

9. Bock-Marquette I, Saxena A, White MD, DiMaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. Nature. 2004;432(7016):466-472. https://doi.org/10.1038/nature03000

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

10. Sosne G, Szliter EA, Barrett R, Kernacki KA, Kleinman HK, Hazlett LD. Thymosin beta 4 promotes corneal wound healing and decreases inflammation in vivo following alkali injury. Exp Eye Res. 2002;74(2):293-299. https://doi.org/10.1006/exer.2001.1125

11. Smart N, Risebro CA, Melville AAD, et al. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. Nature. 2007;445(7124):177-182. https://doi.org/10.1038/nature05383

12. Sosne G, Xu L, Prach L, et al. Thymosin beta 4 stimulates laminin-5 production independent of TGF-beta. Exp Cell Res. 2004;293(1):175-183. https://doi.org/10.1016/j.yexcr.2003.09.022

13. Staresinic M, Sebecic B, Patrlj L, et al. Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth. J Orthop Res. 2003;21(6):976-983. https://doi.org/10.1016/S0736-0266(03)00110-4

14. Miller TR, Wagner JD, Baack BR, Eisbach KJ. Effects of topical copper tripeptide complex on CO2 laser-resurfaced skin. Arch Facial Plast Surg. 2006;8(4):252-259. https://doi.org/10.1001/archfaci.8.4.252

15. Kang EA, Han YM, An JM, et al. BPC157 as potential agent rescuing from cancer cachexia. Curr Pharm Des. 2018;24(18):1947-1956. https://doi.org/10.2174/1381612824666180614082950

16. Xing Y, Ye Y, Zuo H, Li Y. Progress on the function and application of thymosin β4. Front Endocrinol (Lausanne). 2021;12:767785. https://doi.org/10.3389/fendo.2021.767785

17. Xue B, Leyrat C, Grimes JM, Robinson RC. Structural basis of thymosin-β4/profilin exchange leading to actin filament polymerization. Proc Natl Acad Sci U S A. 2014;111(43):E4596-E4605. https://doi.org/10.1073/pnas.1412271111

18. Ruff D, Crockford D, Girardi G, Zhang Y. A randomized, placebo-controlled, single and multiple dose study of intravenous thymosin beta4 in healthy volunteers. Ann N Y Acad Sci. 2010;1194:223-229. https://doi.org/10.1111/j.1749-6632.2010.05474.x

19. Guarnera G, DeRosa A, Camerini R. The effect of thymosin treatment of venous ulcers. Ann N Y Acad Sci. 2010;1194:207-212. https://doi.org/10.1111/j.1749-6632.2010.05490.x

20. Sosne G, Ousler GW. Thymosin beta 4 ophthalmic solution for dry eye: a randomized, placebo-controlled, Phase II clinical trial conducted using the controlled adverse environment (CAE™) model. Clin Ophthalmol. 2015;9:877-884. https://doi.org/10.2147/OPTH.S80954

21. Sosne G, Kleinman HK, Springs C, Gross RH, Sung J, Kang S. 0.1% RGN-259 (Thymosin ß4) ophthalmic solution promotes healing and improves comfort in neurotrophic keratopathy patients in a randomized, placebo-controlled, double-masked Phase III clinical trial. Int J Mol Sci. 2023;24(1):554. Published online December 29, 2022. https://doi.org/10.3390/ijms24010554

22. Philp D, Nguyen M, Scheremeta B, et al. Thymosin beta4 increases hair growth by activation of hair follicle stem cells. FASEB J. 2004;18(2):385-387. https://doi.org/10.1096/fj.03-0244fje

23. Li X, Wang L, Chen C. Effects of exogenous thymosin β4 on carbon tetrachloride-induced liver injury and fibrosis. Sci Rep. 2017;7(1):5872. https://doi.org/10.1038/s41598-017-06318-5

24. Kim J, Lee C, Han J, et al. Targeted deletion of thymosin beta 4 in hepatic stellate cells ameliorates liver fibrosis in a transgenic mouse model. Cells. 2023;12(12):1658. https://doi.org/10.3390/cells12121658

25. Cha HJ, Jeong MJ, Kleinman HK. Role of thymosin beta4 in tumor metastasis and angiogenesis. J Natl Cancer Inst. 2003;95(22):1674-1680. https://doi.org/10.1093/jnci/djg100

26. Caers J, Hose D, Kuipers I, et al. Thymosin β4 has tumor suppressive effects and its decreased expression results in poor prognosis and decreased survival in multiple myeloma. Haematologica. 2010;95(1):163-167. https://doi.org/10.3324/haematol.2009.006411

27. Gonzalez-Franquesa A, Stocks B, Borg ML, et al. Discovery of thymosin β4 as a human exerkine and growth factor. Am J Physiol Cell Physiol. 2021;321(5):C770-C778. https://doi.org/10.1152/ajpcell.00263.2021

28. Tokura Y, Nakayama Y, Fukada S, et al. Muscle injury-induced thymosin β4 acts as a chemoattractant for myoblasts. J Biochem. 2011;149(1):43-48. https://doi.org/10.1093/jb/mvq115

29. Janoshik Analytical. TB4/TB-500/TB4(17-23) analysis. Accessed October 1, 2026. https://janoshik.com/pricelist/tb4-tb-500-tb417-23-analysis/

30. Pickart L, Vasquez-Soltero JM, Pickart FD, Majnarich J. GHK, the human skin remodeling peptide, induces anti-cancer expression of numerous caspase, growth regulatory, and DNA repair genes. J Anal Oncol. 2014;3(2):79-87. https://neoplasiaresearch.com/pms/index.php/jao/article/download/217/212/209

31. Hsieh MJ, Liu HT, Wang CN, et al. Therapeutic potential of pro-angiogenic BPC157 is associated with VEGFR2 activation and up-regulation. J Mol Med (Berl). 2017;95(3):323-333. https://doi.org/10.1007/s00109-016-1488-y