KLOW Blend (GHK-CU / BPC-157 / TB-500(TB4) / KPV) Available for Research Use Only (1.6 mg for cosmetic effects, 3.2 mg for tissue repair, SC daily, cycled)

KLOW is a peptide blend that combines GHK-Cu, BPC-157, TB-500 (TB4), and KPV, typically in a 5:1:1:1 ratio. It is intended to support skin quality, soft tissue healing, mucosal integrity, and a low-noise systemic anti-inflammatory tone. The rationale is to combine GHK-Cu gene-level regenerative signaling and copper delivery, with TB-500 (TB4) actin and migration support, BPC-157 endothelial and gut-vascular stabilization, and KPV's likely intestinal anti-inflammatory effects and plausible epithelial-barrier support.[32–34,36,38] Together these signals aim to improve extracellular matrix quality, microvascular perfusion, tissue remodeling, and mucosal integrity. [1,8,9,13,25,26] TB-500 (TB4)'s effects on cell migration and angiogenesis warrant caution in people with active cancer, a history of cancer, or elevated cancer risk.

💉 💉 Probably Safe

C Angiogenic – caution for subjects at elevated risk for cancers, Do NOT use with active cancer

⭐ ⭐ Probably effective (strong animal signals and/or some positive human signals)

Disclaimer: These notes are not authoritative and should not be interpreted as definitive scientific guidance. They are summaries, compilations and plausible extrapolations drawn from the most relevant research I was able to locate, and may omit nuances, conflicting data, or emerging evidence. The material reflects an attempt to organize and understand complex topics, not to provide clinical recommendations or expert interpretation.

This document was updated 12-Oct-26. The most recent version of this document is available at https://pdfs.researcher6076.com/pdfs/klow-blend.pdf

Short term human safety of the blend is informed by the safety of the individual peptides. TB-500 (TB4) has short-term human tolerability data from intravenous studies, with additional observations from local eye and wound treatments. GHK-Cu short-term safety is likely based on animal evidence. BPC-157 has positive, albeit limited human safety signals, along with very widespread community use with few serious adverse events reported. Long-term safety of these injected components remains uncertain. [18-21] KPV's safety assessment rests on preclinical observations and community reports. Long-term safety of the injected KLOW blend remains probable. [18–21,32–34,36,38]

Animal studies support likely healing benefits from individual KLOW components. GHK-Cu research supports skin repair and extracellular-matrix remodeling, with some topical human findings for cosmetic improvement. Animal studies support likely hair-growth benefits from TB-500 (TB4) through hair-follicle stem-cell activity. Animal studies also support likely healing benefits from TB-500 (TB4) and BPC-157. Synergies in the combination are plausible. [1-3,8,13,14,22] KPV has reproducible preclinical anti-inflammatory and barrier data and growing

experiential use. **The combined blend is mechanistically coherent but lacks controlled human subcutaneous trials.**

*The names **TB4**, **Tβ4** and **TB-500** are used inconsistently in gray market and research-use-only community. Vendor labels may use "TB-500" for either the fragment or full-length Tβ4. Labels such as "TB-500 (TB4)" or "TB-500 (frag)" can help indicate the intended identity, but the name alone does not establish what a sample contains.*

Janoshik's reporting terminology helps clarify these distinctions. "TB-500 (TB-4)" identifies full-length, 43-amino-acid Tβ4, while "TB-500 889 (Da)" identifies the acetylated fragment corresponding to residues 17–23, Ac-LKKTETQ. A Janoshik report labeled only "TB-500" identifies some fragment, with its specific identity given in the comments. [29]

This research survey applies to KLOW containing full-length, 43-amino-acid thymosin β4. Throughout this document, "TB-500 (TB4)" refers to that full-length peptide. [29]

Dosage & Patterns

In the 50/10/10/10 blend KLOW dosing ratio of GHK-Cu, TB-500 (TB4), BPC-157, and KPV are fixed to that ratio. Dose guidance relies on mechanistic extrapolation from animal studies, common practices at med spas, and reports by forum users of peptides. The following pattern emerges

- **1.6 mg / daily KLOW, for skin & hair**
  - ~ 1 mg GHK-Cu / daily
  - ~ 200 mcg BPC-157 / daily
  - ~ 200 mcg TB-500 (TB4) / daily
  - ~ 200 mcg KPV / daily
- **3.2 mg / daily KLOW, for tissue repair**
  - ~ 2 mg GHK-Cu / daily
  - ~ 400 mcg BPC-157 / daily
  - ~ 400 mcg TB-500 (TB4) / daily
  - ~ 400 mcg KPV / daily

With the lower dose for skin and hair repair and regeneration, and the higher dose for more aggressive injury and tissue repair.

Benefits

The benefits below draw on research involving the individual peptides and reports from KLOW users. TB-500 (TB4) has human tolerability and local-treatment data alongside animal repair studies. GHK-Cu has topical human and preclinical findings, while BPC-157 and KPV's evidence is predominantly preclinical. Synergies in the combination are plausible. Community reports describe user experiences and are identified separately from scientific findings. [1,13,18-21]

- **Smoother, healthier-looking skin**, with improvements in hydration, elasticity, and overall tone.
- **Likely support for wound repair,** based on animal studies.
- **Faster recovery from soft tissue strains** is plausible through effects on cell migration, vascular repair, and tissue remodeling. [8,11,17]
- **Reduced joint stiffness and background inflammation**, often noticed in knees, shoulders, and hips as connective-tissue signaling normalizes.
- **Plausible support for recovery from minor soft-tissue injuries and overuse irritation** through effects on vascular repair and extracellular matrix remodeling. Synergies in the combination are plausible. Relief of delayed-onset muscle soreness remains a community-reported benefit. [3,8,11,31]
- **Calmer old injury sites**, with many users describing less reactivity or "flare-ups" in previously problematic areas.
- **Likely support for tendon healing**, based on animal studies. Improved tendon and ligament resilience during training remains plausible rather than demon-strated. [5,8,13]
- **Likely support for vascular repair** based on animal injury studies. **Improved microvascular function, nutrient delivery, and tissue oxygenation** under mechanical load are plausible benefits. [9,11]
- **More even skin texture**, including subtle reductions in fine lines and improved firmness.
- **Generalized tissue "comfort"**, a broad effect people describe as **less tightness, fewer aches, and smoother movement patterns**.
- **Likely reduction in intestinal inflammation and neutrophil recruitment, with plausible support for broader anti-inflammatory effects. [32–34,37]**
- **Plausible support for intestinal epithelial and mucosal-barrier integrity through reduced inflammatory signaling. [32,36,38]**

Applications

- Chronic soft-tissue irritation
- Older injuries that flare under load
- High training volume
- Age-related changes in skin quality
- Demanding physical jobs or lifestyles
- Recovering from minor strains or overuse
- Low-grade systemic inflammation
- Running GH-axis peptides

Cautions

- Copper metabolism disorders
- Liver disease
- Active cancer, elevated cancer risk

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

Side Effects

- Nausea, headache, and fatigue have been reported by users. Their frequency and causal relationship to KLOW are uncertain.

Drivers of Diminished Response and Mitigation Strategies

None of TB-500 (TB4), BPC-157, GHK-Cu, or KPV are receptor agonists. Downstream adaptation within repair pathways is plausible but without evidence. [8,17]

KLOW: Plausible Longevity Benefits

The plausibly pro-longevity rationale for the KLOW blend arises from the convergence of **four distinct but complementary biological programs**: GHK-Cu's gene-level regenerative and anti-inflammatory signaling, TB-500 (TB4)'s cytoskeletal and angiogenic repair pathways, BPC-157's gastrointestinal, endothelial, and microvascular stabilization effects, and KPV's anti-inflammatory effects. Each peptide **targets a different conserved pathway in age-related decline**, and together they form a mechanistically coherent strategy for supporting tissue integrity, repair capacity, and inflammatory balance, domains that strongly influence healthspan.

GHK-Cu contributes a broad genereprogramming signal that shifts cellular behavior away from proinflammatory and prooxidative states toward regeneration, extracellularmatrix rebuilding, and antioxidant defense. Age-related declines in GHK-Cu correlate with reduced collagen synthesis, impaired wound healing, and increased inflammatory tone; restoring this signal may help maintain ECM quality, reduce chronic inflammatory burden, and preserve microcirculatory function. **These actions address structural aging at the level of collagen, elastin, glycosaminoglycans, and metalloprotein regulation, supporting the connectivetissue architecture that underpins organ resilience.**

TB-500 (TB4) contributes actin regulation, cell migration, and vascular repair to KLOW's proposed healthspan rationale. Animal studies support likely benefits for wound healing and selected forms of tissue repair. Its effects on fibrosis vary by tissue and experimental context, so a general antifibrotic benefit should not be assumed. **These repair activities plausibly support tissue resilience during aging.** Synergies with GHK-Cu's extra-cellular-matrix- effects are plausible. [8,9,11,17,23,24]

BPC-157 complements both signals by stabilizing the gastrointestinal barrier, protecting microvasculature, and modulating nitricoxide pathways that influence endothelial tone and perfusion. Because aging is strongly shaped by chronic lowgrade inflammation originating from gut permeability, vascular instability, and impaired angiogenic balance, BPC-157's protective effects on the gut–vascular axis may reduce systemic inflammatory load and improve nutrient delivery, oxygenation, and waste clearance. Its antiulcer, endothelial-protective, and tendon-repair properties add a third dimension to the blend's regenerative profile, supporting both local and systemic resilience.

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

KPV contributes to KLOW's proposed healthspan rationale **through likely reductions in intestinal inflammation and neutrophil recruitment, with plausible support for mucosal-barrier integrity and broader inflammatory regulation.** These effects plausibly complement the repair pathways of GHK-Cu, TB-500 (TB4), and BPC-157, supporting tissue resilience during aging. Synergies in the combination and benefits for healthspan remain plausible. [32–34,36–38]

Synergies in combining these four peptides are plausible, providing a rationale for supporting tissue resilience and healthspan. [1-3,8,13,31-34,36-38]

KLOW and Cancer Risk

The cancer concern with KLOW involves possible support of an existing malignancy through effects on angiogenesis and cell migration. **TB4 tumor-promotion findings in animal models make this concern more than purely *theoretical***, *although tumor-cell expression studies are not equivalent to administering TB-500 (TB4). These findings do not establish that KLOW causes new cancers, and its cancer risk* in humans remains unresolved. [25,26]

GHK-Cu has experimental findings relevant to protective cellular responses and cancer-related gene expression. In a mouse sarcoma model, GHK-Cu combined with ascorbic acid reduced tumor growth. Although mildly proangiogenic, it has not been linked to tumor promotion in vivo. But these findings do not establish general cancer protection or show that GHK-Cu offsets potential tumor-related concerns from TB-500 (TB4). [1,30]

BPC-157 has experimental findings suggesting protective effects in selected settings. Experimental work discussed by Kang and colleagues suggests that BPC-157 may counter cancer-associated muscle wasting and related inflammatory changes, but these benefits do not establish suppression of tumor growth or metastasis. [15]

KPV likely reduces inflammation-associated intestinal tumor development, based on a mouse model of colitis-associated cancer in which its protective effect depended on PepT1-mediated uptake. Broader cancer-protective effects remain plausible. [35]

TB-500 (TB4) has direct experimental evidence relevant to cancer risk. Increased TB4 expression promoted tumor vascularization and metastasis in a mouse melanoma model. TB-500 (TB4) is overexpressed in several cancers and associated with metastasis and poor prognosis. [16,25] TB-500 (TB4) regulates actin dynamics and supports cell migration and angiogenesis during repair. These processes can also contribute to tumor progression. [17,25] Other studies found tumor-suppressive effects in multiple myeloma. **These findings support caution around an existing malignancy and caution around an elevated cancer risk**. [25,26]

Conservative Alternatives Around Cancer Risks

For someone without elevated cancer risk, but with a desire to be cautious, three alternative patterns are available

- Using GHK-Cu and KPV offers a simpler approach to skin repair and extracellular-matrix support while avoiding exposure to TB-500 (TB4) and BPC-157.
- Combine GHK-Cu, KPV, and BPC-157. This provides additional soft tissue support while avoiding exposure to TB-500 (TB4).
- Alternating vials of KLOW and GHK-Cu (possibly with KPV and/or BPC-157) is a speculative approach to reducing continuous exposure to TB-500 (TB4).

Zinc Supplementation with GHK-Cu

Practitioners sometimes **consider zinc supplementation alongside GHK-Cu** because zinc helps maintain balanced metal homeostasis in the body. GHK-Cu delivers copper to enzymes involved in repair, and zinc plays a complementary role by supporting antioxidant defenses, stabilizing cellular signaling, improved immune response, and helping regulate metalloproteins that manage copper–zinc balance. When overall zinc status is low, the body may have more difficulty buffering shifts in copper availability, so **maintaining adequate zinc intake can help keep these metal-dependent pathways stable while GHK-Cu is being used**. The zinc requirement is not large, probably 20 mg daily is sufficient to support the continuous maximum dose of GHK-Cu.

Biological Mechanism

GHK-Cu is a naturally occurring copper-binding tripeptide that functions as a repair and remodeling signal across multiple tissues. Biologically, it binds copper(II) and delivers it to enzymes involved in collagen cross-linking, antioxidant defense (such as superoxide dismutase), and mitochondrial energy production, while simultaneously activating gene programs linked to extracellular-matrix rebuilding, angiogenesis, and wound healing. Physiologically, GHK-Cu reduces inflammatory signaling through suppression of NF-κB, TNF-α, and IL-6, increases the expression of structural proteins like collagen, elastin, and glycosaminoglycans, and promotes fibroblast and keratinocyte migration during tissue repair. It also improves microcirculation by supporting capillary growth and endothelial function and has been shown to modulate over 4,000 human genes toward a regenerative, anti-inflammatory profile. This combination of copper transport, gene-level repair signaling, and inflammation control is why GHK-Cu appears across research on skin quality, connective-tissue healing, and general tissue regeneration.

BPC-157 is a cytoprotective, angiogenic, and anti-inflammatory peptide that operates through several converging biological pathways. It stabilizes the gut-vascular axis by up-regulating VEGF, FGF-2, and other growth-repair signals while simultaneously suppressing TNF-α, IL-6, and NF-κB-driven inflammation. At the cellular level, BPC-157 enhances fibroblast migration, collagen organization, and endothelial repair, which supports tendon, ligament, and soft-tissue healing. It also interacts with the nitric-oxide system to normalize microvascular tone and protect against oxidative stress. In animal and early mechanistic studies, BPC-157 preserves tight-junction integrity in the gut, accelerates angiogenesis in injured tissue, modulates

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

dopamine and serotonin pathways, and promotes rapid healing across muscle, nerve, and gastrointestinal models. Together, these effects position BPC-157 as a broad-spectrum repair peptide that coordinates inflammation control, vascular restoration, and extracellular-matrix rebuilding.

TB-500 (TB4) is a naturally occurring 43-amino-acid peptide that binds and sequesters globular actin, helping regulate actin availability for cell movement and tissue repair. Animal studies support likely benefits for wound healing, cardiac protection, and selected inflammatory responses. Cell studies also show effects on epithelial migration and inflammatory signaling.

KPV is the C-terminal tripeptide (Lys-Pro-Val) of α-melanocyte-stimulating hormone. Its anti-inflammatory effects plausibly involve PepT1-mediated uptake into intestinal epithelial and immune cells, followed by inhibition of NF-κB and MAPK signaling and reduced inflammatory cytokine production. These effects are largely independent of melanocortin-receptor activation. Studies in human bronchial epithelial cells also support a plausible mechanism involving inhibition of NF-κB nuclear entry. [32–34,37] KPV plausibly supports a shift toward an M2 repair-associated macrophage phenotype; this effect was demonstrated in mice receiving the engineered oral proKPV formulation. [38] KPV likely reduces neutrophil recruitment. [33,34] It plausibly supports intestinal epithelial and mucosal barrier integrity through reduced inflammatory signaling. In mice with colitis, the engineered oral KPV formulation restored occludin and ZO-1 expression and reduced intestinal permeability. These findings support a plausible barrier-repair rationale for KPV in KLOW. [32,36,38] KPV likely reduces intestinal inflammation, with plausible support for mucosal-barrier integrity and broader anti-inflammatory applications in skin and other tissues. [32–34,37]

References

1. Pickart L, Margolina A. Regenerative and protective actions of the GHK-Cu peptide in the light of the new gene data. Int J Mol Sci. 2018;19(7):1987. https://doi.org/10.3390/ijms19071987
2. Maquart FX, Pickart L, Laurent M, Gillery P, Monboisse JC, Borel JP. Stimulation of collagen synthesis in fibroblast cultures by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu2+. FEBS Lett. 1988;238(2):343-346. https://doi.org/10.1016/0014-5793(88)80509-X
3. Siméon A, Wegrowski Y, Bontemps Y, Maquart FX. Expression of glycosaminoglycans and small proteoglycans in wounds: modulation by the tripeptide-copper complex glycyl-L-histidyl-L-lysine-Cu(2+). J Invest Dermatol. 2000;115(6):962-968. https://doi.org/10.1046/j.1523-1747.2000.00166.x
4. Sikiric P, Seiwerth S, Rucman R, et al. Stable gastric pentadecapeptide BPC 157: novel therapy in gastrointestinal tract. Curr Pharm Des. 2011;17(16):1612-1632. https://doi.org/10.2174/138161211796196954
5. Chang CH, Tsai WC, Lin MS, Hsu YH, Pang JH. The promoting effect of pentadecapeptide BPC 157 on tendon healing involves tendon outgrowth, cell survival, and cell migration. J Appl Physiol (1985). 2011;110(3):774-780. https://doi.org/10.1152/japplphysiol.00945.2010

This document provides educational summaries of limited and evolving peptide data. Nothing here is medical advice for diagnosis or treatment. © 2026 Researcher6076

6. Goldstein AL, Hannappel E, Sosne G, Kleinman HK. Thymosin β4: a multi-functional regenerative peptide. Basic properties and clinical applications. Expert Opin Biol Ther. 2012;12(1):37-51. https://doi.org/10.1517/14712598.2012.634793
7. Sosne G, Qiu P, Christopherson PL, Wheater MK. Thymosin beta 4 suppression of corneal NFκB: a potential anti-inflammatory pathway. Exp Eye Res. 2007;84(4):663-669. https://doi.org/10.1016/j.exer.2006.12.004
8. Malinda KM, Sidhu GS, Mani H, et al. Thymosin beta4 accelerates wound healing. J Invest Dermatol. 1999;113(3):364-368. https://doi.org/10.1046/j.1523-1747.1999.00708.x
9. Bock-Marquette I, Saxena A, White MD, DiMaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. Nature. 2004;432(7016):466-472. https://doi.org/10.1038/nature03000
10. Sosne G, Szliter EA, Barrett R, Kernacki KA, Kleinman HK, Hazlett LD. Thymosin beta 4 promotes corneal wound healing and decreases inflammation in vivo following alkali injury. Exp Eye Res. 2002;74(2):293-299. https://doi.org/10.1006/exer.2001.1125
11. Smart N, Risebro CA, Melville AAD, et al. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. Nature. 2007;445(7124):177-182. https://doi.org/10.1038/nature05383
12. Sosne G, Xu L, Prach L, et al. Thymosin beta 4 stimulates laminin-5 production independent of TGF-beta. Exp Cell Res. 2004;293(1):175-183. https://doi.org/10.1016/j.yexcr.2003.09.022
13. Staresinic M, Sebecic B, Patrlj L, et al. Gastric pentadecapeptide BPC 157 accelerates healing of transected rat Achilles tendon and in vitro stimulates tendocytes growth. J Orthop Res. 2003;21(6):976-983. https://doi.org/10.1016/S0736-0266(03)00110-4
14. Miller TR, Wagner JD, Baack BR, Eisbach KJ. Effects of topical copper tripeptide complex on CO2 laser-resurfaced skin. Arch Facial Plast Surg. 2006;8(4):252-259. https://doi.org/10.1001/archfaci.8.4.252
15. Kang EA, Han YM, An JM, et al. BPC157 as potential agent rescuing from cancer cachexia. Curr Pharm Des. 2018;24(18):1947-1956. https://doi.org/10.2174/1381612824666180614082950
16. Xing Y, Ye Y, Zuo H, Li Y. Progress on the function and application of thymosin β4. Front Endocrinol (Lausanne). 2021;12:767785. https://doi.org/10.3389/fendo.2021.767785
17. Xue B, Leyrat C, Grimes JM, Robinson RC. Structural basis of thymosin-β4/profilin exchange leading to actin filament polymerization. Proc Natl Acad Sci U S A. 2014;111(43):E4596-E4605. https://doi.org/10.1073/pnas.1412271111
18. Ruff D, Crockford D, Girardi G, Zhang Y. A randomized, placebo-controlled, single and multiple dose study of intravenous thymosin beta4 in healthy volunteers. Ann N Y Acad Sci. 2010;1194:223-229. https://doi.org/10.1111/j.1749-6632.2010.05474.x
19. Guarnera G, DeRosa A, Camerini R. The effect of thymosin treatment of venous ulcers. Ann N Y Acad Sci. 2010;1194:207-212. https://doi.org/10.1111/j.1749-6632.2010.05490.x
20. Sosne G, Ousler GW. Thymosin beta 4 ophthalmic solution for dry eye: a randomized, placebo-controlled, Phase II clinical trial conducted using the controlled adverse

environment (CAE™) model. Clin Ophthalmol. 2015;9:877-884. https://doi.org/10.2147/OPTH.S80954

21. Sosne G, Kleinman HK, Springs C, Gross RH, Sung J, Kang S. 0.1% RGN-259 (Thymosin ß4) ophthalmic solution promotes healing and improves comfort in neurotrophic keratopathy patients in a randomized, placebo-controlled, double-masked Phase III clinical trial. Int J Mol Sci. 2023;24(1):554. Published online December 29, 2022. https://doi.org/10.3390/ijms24010554
22. Philp D, Nguyen M, Scheremeta B, et al. Thymosin beta4 increases hair growth by activation of hair follicle stem cells. FASEB J. 2004;18(2):385-387. https://doi.org/10.1096/fj.03-0244fje
23. Li X, Wang L, Chen C. Effects of exogenous thymosin β4 on carbon tetrachloride-induced liver injury and fibrosis. Sci Rep. 2017;7(1):5872. https://doi.org/10.1038/s41598-017-06318-5
24. Kim J, Lee C, Han J, et al. Targeted deletion of thymosin beta 4 in hepatic stellate cells ameliorates liver fibrosis in a transgenic mouse model. Cells. 2023;12(12):1658. https://doi.org/10.3390/cells12121658
25. Cha HJ, Jeong MJ, Kleinman HK. Role of thymosin beta4 in tumor metastasis and angiogenesis. J Natl Cancer Inst. 2003;95(22):1674-1680. https://doi.org/10.1093/jnci/djg100
26. Caers J, Hose D, Kuipers I, et al. Thymosin β4 has tumor suppressive effects and its decreased expression results in poor prognosis and decreased survival in multiple myeloma. Haematologica. 2010;95(1):163-167. https://doi.org/10.3324/haematol.2009.006411
27. Gonzalez-Franquesa A, Stocks B, Borg ML, et al. Discovery of thymosin β4 as a human exerkine and growth factor. Am J Physiol Cell Physiol. 2021;321(5):C770-C778. https://doi.org/10.1152/ajpcell.00263.2021
28. Tokura Y, Nakayama Y, Fukada S, et al. Muscle injury-induced thymosin β4 acts as a chemoattractant for myoblasts. J Biochem. 2011;149(1):43-48. https://doi.org/10.1093/jb/mvq115
29. Janoshik Analytical. TB4/TB-500/TB4(17-23) analysis. Accessed October 1, 2026. https://janoshik.com/pricelist/tb4-tb-500-tb417-23-analysis/
30. Pickart L, Vasquez-Soltero JM, Pickart FD, Majnarich J. GHK, the human skin remodeling peptide, induces anti-cancer expression of numerous caspase, growth regulatory, and DNA repair genes. J Anal Oncol. 2014;3(2):79-87. https://neoplasiaresearch.com/index.php/jao/article/view/217
31. Hsieh MJ, Liu HT, Wang CN, et al. Therapeutic potential of pro-angiogenic BPC157 is associated with VEGFR2 activation and up-regulation. J Mol Med (Berl). 2017;95(3):323-333. https://doi.org/10.1007/s00109-016-1488-y
32. Dalmasso G, Charrier-Hisamuddin L, Nguyen HT, et al. PepT1-mediated tripeptide KPV uptake reduces intestinal inflammation. Gastroenterology. 2008;134(1):166-178. https://doi.org/10.1053/j.gastro.2007.10.026
33. Kannengiesser K, Maaser C, Heidemann J, et al. Melanocortin-derived tripeptide KPV has anti-inflammatory potential in murine models of inflammatory bowel disease. Inflamm Bowel Dis. 2008;14(3):324-331. https://doi.org/10.1002/ibd.20334

34. Getting SJ, Schiöth HB, Perretti M. Dissection of the anti-inflammatory effect of the core and C-terminal (KPV) alpha-melanocyte-stimulating hormone peptides. J Pharmacol Exp Ther. 2003;306(2):631-637. https://doi.org/10.1124/jpet.103.051623
35. Viennois E, Ingersoll SA, Ayyadurai S, et al. Critical role of PepT1 in promoting colitis-associated cancer and therapeutic benefits of the anti-inflammatory PepT1-mediated tripeptide KPV in a murine model. Cell Mol Gastroenterol Hepatol. 2016;2(3):340-357. https://doi.org/10.1016/j.jcmgh.2016.01.006
36. Xiao B, Xu Z, Viennois E, et al. Orally Targeted Delivery of Tripeptide KPV via Hyaluronic Acid-Functionalized Nanoparticles Efficiently Alleviates Ulcerative Colitis. Mol Ther. 2017;25(7):1628-1640. https://doi.org/10.1016/j.ymthe.2016.11.020
37. Land SC. Inhibition of cellular and systemic inflammation cues in human bronchial epithelial cells by melanocortin-related peptides: mechanism of KPV action and a role for MC3R agonists. Int J Physiol Pathophysiol Pharmacol. 2012;4(2):59-73. https://pubmed.ncbi.nlm.nih.gov/22837805/
38. Cheng J, Wu P, Li C, et al. Inflammation-triggered self-immolative conjugates enable oral peptide delivery by overcoming gastrointestinal barriers. Sci Adv. 2026;12(3):eaea2989. https://doi.org/10.1126/sciadv.aea2989
39. Elliott RJ, Szabo M, Wagner MJ, et al. alpha-Melanocyte-stimulating hormone, MSH 11-13 KPV and adrenocorticotropic hormone signalling in human keratinocyte cells. J Invest Dermatol. 2004;122(4):1010-1019. https://doi.org/10.1111/j.0022-202x.2004.22404.x