TME groups ranked in the top 10 in every fold

PathoTME-LR · 16-shot · five held-out folds · rank 1 = highest attribution

NSCLC

Predicted LUSC probability · 4 of 16 groups

GROUP · MEAN (pp)	FOLD RANK 0	1	2	3	4	TOP 5
Fibroblast spatial context 4.76	3	3	1	6	1	4/5
Lymphocyte spatial context 3.10	9	2	5	9	4	3/5
Granulocyte measurements 2.58	4	5	8	3	6	3/5
Endothelial-cell measurements 2.46	2	6	9	2	7	2/5

12 groups do not meet this rule

BRCA

Predicted ILC probability · 3 of 16 groups

GROUP · MEAN (pp)	FOLD RANK 0	1	2	3	4	TOP 5
Macrophage measurements 2.69	1	8	1	1	1	4/5
Stroma geometry 1.60	8	7	10	5	9	1/5
Tissue composition 1.46	7	10	5	6	5	2/5

13 groups do not meet this rule

CRC

Predicted MUCINOUS probability · 2 of 16 groups

GROUP · MEAN (pp)	FOLD RANK 0	1	2	3	4	TOP 5
Global tissue 4.48	1	9	2	8	4	3/5
Carcinoma-cell measurements 3.89	5	7	7	10	9	1/5

14 groups do not meet this rule

BLCA

Predicted PAPILLARY probability · 1 of 16 groups

GROUP · MEAN (pp)	FOLD RANK 0	1	2	3	4	TOP 5
Granulocyte measurements 3.46	10	3	5	6	3	3/5

15 groups do not meet this rule

Selection: rank ≤10 of 16 groups in all five folds. Darker cells indicate higher ranks.

Rank consistency does not establish stable effect sizes or predictive usefulness. All fold scores are retained in the CSV.

Mean attribution: absolute patient-level probability change after joint group replacement, averaged over five folds (pp).