| Study                                 | Non-responder | Responder | Standardised Mean<br>Difference | SMD   | 95%-CI        | Weight (fixed) | Weight (random) |
|---------------------------------------|---------------|-----------|---------------------------------|-------|---------------|----------------|-----------------|
| Chaput 2017                           | 17            | 9         |                                 | 0.00  | [-0.81; 0.81] | 8.4%           | 8.4%            |
| Frankel 2017                          | 15            | 24        |                                 | -0.34 | [-0.99; 0.31] | 13.0%          | 13.0%           |
| Gopalakrishnan 2017                   | 13            | 30        | <del></del>                     | 0.00  | [-0.65; 0.65] | 13.0%          | 13.0%           |
| Matson 2018                           | 26            | 16        |                                 | 0.00  | [-0.62; 0.62] | 14.2%          | 14.2%           |
| Routy NSCLC 2017                      | 46            | 41        | #                               | 0.28  | [-0.14; 0.70] | 30.8%          | 30.8%           |
| Routy RCC 2017                        | 21            | 45        |                                 | 0.08  | [-0.44; 0.60] | 20.5%          | 20.5%           |
| Fixed effect model                    | 138           | 165       |                                 | 0.06  | [-0.18; 0.29] | 100.0%         |                 |
| Random effects model                  | I             |           | <b>*</b>                        |       | [-0.18; 0.29] |                | 100.0%          |
| Heterogeneity: $I^2 = 0\%$ , $\tau^2$ | = 0, p = 0.77 |           |                                 |       | - · · -       |                |                 |