| Study | Non-responder R | esponder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|---|-----------------|----------|---------------------------------|----------|--------------------------------|----------------|--------------------|
| Chaput 2017 | 17 | 9 | - i | -0.29 [| [–1.11; 0.52] | 8.5% | 12.8% |
| Frankel 2017 | 15 | 24 | - } • - - | 0.22 [| [-0.43; 0.87] | 13.4% | 15.9% |
| Gopalakrishnan 2017 | 13 | 30 - | | -1.12 [- | -1.82; -0.43] | 11.5% | 14.9% |
| Matson 2018 | 26 | 16 | 3 - | 0.27 | [-0.36; 0.89] | 14.3% | 16.4% |
| Routy NSCLC 2017 | 46 | 41 | | -0.18 [| [-0.60; 0.24] | 31.5% | 21.2% |
| Routy RCC 2017 | 21 | 45 | 1 . | 0.27 | [-0.25; 0.79] | 20.7% | 18.8% |
| Fixed effect model | 138 | 165 | | - | -0.32; 0.15] | | |
| Random effects mode Heterogeneity: $I^2 = 61\%$, | * - | | | | [-0.51 ; 0.28] | | 100.0% |