| Study | Non-responder | Responder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|--------------------------------------|--------------------------|-----------|---------------------------------|---------------|---------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | į: — - | — 1.09 | [0.22; 1.96] | 7.4% | 10.2% |
| Frankel 2017 | 15 | 24 | <u>i</u> | 0.11 | [-0.54; 0.75] | 13.4% | 15.4% |
| Gopalakrishnan 2017 | 13 | 30 | - 1: | 0.29 | [-0.36; 0.95] | 13.0% | 15.1% |
| Matson 2018 | 26 | 16 | - ii | -0.23 | [-0.85; 0.40] | | 16.0% |
| Routy NSCLC 2017 | 46 | 41 | 1: | | [-0.66; 0.19] | | 23.7% |
| Routy RCC 2017 | 21 | 45 | | 0.15 | [-0.37; 0.66] | 20.7% | 19.6% |
| Fixed effect model | 138 | 165 | | 0.06 | [-0.18; 0.29] | 100.0% | |
| Random effects model | | | * | 0.11 | [-0.21; 0.43] | | 100.0% |
| Heterogeneity: $I^2 = 43\%$, τ | $t^2 = 0.0676, p = 0.12$ | <u> </u> | | | - | | |
| | • | -2 | -1 0 1 | 2 | | | |