| Study | Non-responder | Responder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|--------------------------------------|--------------------------|-----------|---------------------------------|---------|-----------------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | -0.09 | [-0.90; 0.72] | 8.5% | 10.0% |
| Frankel 2017 | 15 | 24 | | -0.04 | [-0.68; 0.61] | 13.3% | 14.5% |
| Gopalakrishnan 2017 | 13 | 30 | - | -0.65 | [-1.32; 0.01] | 12.5% | 13.7% |
| Matson 2018 | 26 | 16 | - | 0.29 | [-0.34; 0.91] | 14.2% | 15.2% |
| Routy NSCLC 2017 | 46 | 41 | | -0.26 | [-0.69; 0.16] | 31.0% | 26.5% |
| Routy RCC 2017 | 21 | 45 | - - | 0.26 | [-0.26; 0.78] | 20.5% | 20.1% |
| Fixed effect model | 138 | 165 | | -0.08 [| [-0.32; 0.15 <u>]</u> | 100.0% | |
| Random effects model | | | | _ | -0.35; 0.20] | | 100.0% |
| Heterogeneity: $I^2 = 24\%$, τ | $c^2 = 0.0278, p = 0.26$ | ; | 1 1 | | · • | | |