Study	Non-responder Re	sponder	Standardised Mean Difference	SMD	95%-CI	Weight (fixed)	Weight (random)
Chaput 2017	17	9		-0.64 [	_1.47; 0.19]	8.1%	11.1%
Frankel 2017	15	24	- 1 -	0.34	-0.31; 0.98]	13.2%	15.3%
Gopalakrishnan 2017	13	30	<del>-   -</del>	0.18	-0.47; 0.83]	13.1%	15.2%
Matson 2018	26	16		-0.22	-0.84; 0.41]	14.3%	16.0%
Routy NSCLC 2017	46	41	3	0.15	-0.27; 0.58]	31.4%	23.4%
Routy RCC 2017	21	45	<del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> - <del>'</del> <del>'</del> - <del>'</del> <del>'</del> <del>'</del>	-0.60 [-	-1.13; -0.07]	19.9%	19.1%
Fixed effect model	138	165		-0.09 [-	-0.32; 0.15]	100.0%	
Random effects mode	I			-0.11 [·	-0.43; 0.22]		100.0%
Heterogeneity: $I^2 = 45\%$ ,	$\tau^2 = 0.0731, p = 0.11$				_		