| Study | Non-responder Re | esponder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|--------------------------------------|---------------------------|----------|---|----------|--------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | 0.00 [- | -0.80; 0.81] | 8.6% | 12.1% |
| Frankel 2017 | 15 | 24 | | -0.03 [- | -0.68; 0.61] | 13.4% | 15.7% |
| Gopalakrishnan 2017 | 13 | 30 | | - 1.04 | 0.35; 1.73] | 11.7% | 14.5% |
| Matson 2018 | 26 | 16 | <u></u> _ | 0.07 [- | -0.55; 0.69] | 14.4% | 16.2% |
| Routy NSCLC 2017 | 46 | 41 | | 0.23 [- | -0.19; 0.65] | 31.3% | 22.4% |
| Routy RCC 2017 | 21 | 45 | - t | -0.34 [- | -0.86; 0.19] | 20.6% | 19.1% |
| Fixed effect model | 138 | 165 | | 0.13 [- | -0.10; 0.37] | 100.0% | |
| Random effects model | | | | - | -0.21; 0.50] | | 100.0% |
| Heterogeneity: $I^2 = 52\%$, τ | 2 = 0.0982, p = 0.06 | | | | · | | |