| Study | Non-responder Re | sponder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|---------------------------------------|------------------|---------|---------------------------------|--------------------|----------------------|-------------------|-----------------|
| Chaput 2017 | 17 | 9 | <u> </u> | 0.00 [- | -0.81; 0.81] | 8.4% | 8.4% |
| Frankel 2017 | 15 | 24 | | -] 00.0 | -0.65; 0.65] | 13.2% | 13.2% |
| Gopalakrishnan 2017 | 13 | 30 | | -0.17 ⁻ | -0.82; 0.48] | 12.9% | 12.9% |
| Matson 2018 | 26 | 16 | | 0.34 [- | -0.28; 0.97] | 14.0% | 14.0% |
| Routy NSCLC 2017 | 46 | 41 | | -] 00.0 | -0.42; 0.42] | 31.0% | 31.0% |
| Routy RCC 2017 | 21 | 45 | | -] 00.0 | -0.52; 0.52] | 20.5% | 20.5% |
| Fixed effect model | 138 | 165 | | 0.03 [- | -0.21; 0.26] | 100.0% | |
| Random effects model | | | * | - | -0.21; 0.26] | | 100.0% |
| Heterogeneity: $I^2 = 0\%$, τ^2 | = 0, p = 0.93 | | | | · | | |