| Study No. | on-responder Res | sponder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|---|------------------|---------|---------------------------------|----------|--------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | -0.35 [- | -1.17; 0.46] | 8.3% | 8.3% |
| Frankel 2017 | 15 | 24 | | 0.11 [- | -0.54; 0.76] | 13.2% | 13.2% |
| Gopalakrishnan 2017 | 13 | 30 | | 0.21 [- | -0.45; 0.86] | 12.9% | 12.9% |
| Matson 2018 | 26 | 16 | | 0.11 [- | -0.52; 0.73] | 14.2% | 14.2% |
| Routy NSCLC 2017 | 46 | 41 | | -0.17 [- | -0.59; 0.25] | 30.9% | 30.9% |
| Routy RCC 2017 | 21 | 45 | - | 0.00 [- | -0.52; 0.52] | 20.5% | 20.5% |
| Fixed effect model | 138 | 165 | | -0.03 [- | -0.26; 0.21] | 100.0% | |
| Random effects model Heterogeneity: $I^2 = 0\%$, $\tau^2 = 0$ | p = 0.86 | | | _ | -0.26; 0.21] | | 100.0% |