| Study | Non-responder | Responder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|---------------------------------------|--------------------------|-----------|--|-------|---------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | -0.05 | [-0.86; 0.75] | 8.5% | 8.8% |
| Frankel 2017 | 15 | 24 | | -0.28 | [-0.93; 0.37] | 13.2% | 13.5% |
| Gopalakrishnan 2017 | 13 | 30 | + - | 0.55 | [-0.12; 1.21] | 12.7% | 13.0% |
| Matson 2018 | 26 | 16 | | 0.58 | [-0.05; 1.22] | 13.7% | 14.0% |
| Routy NSCLC 2017 | 46 | 41 | | 0.15 | [-0.27; 0.57] | 31.3% | 30.4% |
| Routy RCC 2017 | 21 | 45 | | 0.34 | [-0.18; 0.86] | 20.5% | 20.4% |
| Fixed effect model | 138 | 165 | | 0.22 | [-0.01; 0.46] | 100.0% | |
| Random effects mode | I | | | 0.22 | [-0.02; 0.47] | | 100.0% |
| Heterogeneity: $I^2 = 4\%$, τ^2 | $p^2 = 0.0042, p = 0.39$ | | | | - · · - | | |