| Study | Non-responder | Responder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|--------------------------------------|--------------------------|-----------|---------------------------------|----------------------|--------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | -0.42 [· | -1.24; 0.40] | 8.3% | 9.3% |
| Frankel 2017 | 15 | 24 | | -0.37 [· | -1.02; 0.28] | 13.1% | 14.0% |
| Gopalakrishnan 2017 | 13 | 30 | - - | 0.33 [- | -0.32; 0.99] | 12.9% | 13.8% |
| Matson 2018 | 26 | 16 | | 0.01 [- | -0.61; 0.64] | 14.3% | 15.1% |
| Routy NSCLC 2017 | 46 | 41 | | 0.20 | -0.22; 0.62] | 31.1% | 27.7% |
| Routy RCC 2017 | 21 | 45 | - 1 | -0.39 [· | -0.91; 0.14] | 20.3% | 20.1% |
| Fixed effect model | 138 | 165 | ÷ | -0.06 [- | -0.29; 0.18] | 100.0% | |
| Random effects model | | | | - | -0.33; 0.20] | | 100.0% |
| Heterogeneity: $I^2 = 17\%$, τ | $p^2 = 0.0185, p = 0.30$ |) | 1 1 | | - | | |