| Study No. | on-responder R | Responder | Standardised Mean Difference | SMD | 95%-CI | Weight (fixed) | Weight (random) |
|---|----------------|-----------|---------------------------------|-------|---------------|----------------|-----------------|
| Chaput 2017 | 17 | 9 | | 0.00 | [-0.81; 0.81] | 8.4% | 8.4% |
| Frankel 2017 | 15 | 24 | - : - | -0.37 | [-1.02; 0.28] | 13.0% | 13.0% |
| Gopalakrishnan 2017 | 13 | 30 | | 0.00 | [-0.65; 0.65] | 13.0% | 13.0% |
| Matson 2018 | 26 | 16 | - i • | 0.00 | [-0.62; 0.62] | 14.2% | 14.2% |
| Routy NSCLC 2017 | 46 | 41 | | -0.23 | [-0.66; 0.19] | 30.9% | 30.9% |
| Routy RCC 2017 | 21 | 45 | | -0.17 | [-0.69; 0.35] | 20.5% | 20.5% |
| Fixed effect model | 138 | 165 | - | -0.16 | [-0.39; 0.08] | 100.0% | |
| Random effects model | | | | -0.16 | [-0.39; 0.08] | | 100.0% |
| Heterogeneity: $I^2 = 0\%$, $\tau^2 = 0$ | p = 0.95 | | | | - , - | | |
| | | 2 | 1 0 1 | 2 | | | |