

Standard Model and beyond

in Colombia and in the World



UNIVERSIDAD DE ANTIOQUIA
1803

Diego Restrepo

Instituto de Física
Universidad de Antioquia
Phenomenology Group
<http://gfif.udea.edu.co>

**XXIX CONGRESO
NACIONAL
vigesimonovenoo DE FÍSICA**

Focus on

Dark matter and neutrinos

In collaboration with


HEP Community in Colombia

Least Action

SO(3) scalar product

$$L = \frac{1}{2}m\mathbf{v}^2 - V(|\mathbf{r}|) = \frac{1}{2}m\mathbf{v} \cdot \mathbf{v} - V(|\mathbf{r}|)$$

Lie groups



$$U = \exp \left(i \sum_j T_j \theta^j \right), \quad (1)$$

where θ^j are the parameters of the transformation and T_j are the generators.

Consider the 1×1

$$K = -i, \quad (2)$$

which generates an element of dilaton group , SO(1), $R(\xi)$

$$\lambda(\xi) = e^\xi, \quad (3)$$

which are just the group of the real exponentials. Such a number can be transformed as

$$x \rightarrow x' = e^\xi x, \quad (4)$$

that corresponds to a boost by e^ξ . We can define the invariant scalar product just as the division of real numbers, such that

$$x \cdot y \rightarrow x' \cdot y' \equiv \frac{x'}{y'} = \frac{e^\xi x}{e^\xi y} = \frac{x}{y} = x \cdot y. \quad (5)$$

Queremos obtener una representación 2×2 del álgebra

$$K = \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix} \rightarrow K^2 = -\mathbf{1}, \quad (6)$$

que genera un elemento del grupo SO(1, 1) con *parámetro* ξ

$$\Lambda = \exp(i\xi K) = \begin{pmatrix} \cosh \xi & \sinh \xi \\ \sinh \xi & \cosh \xi \end{pmatrix}, . \quad (7)$$

La transformación de una coordenada temporaloide y otra espacialoide ($c = 1$)


$$\begin{pmatrix} t \\ x \end{pmatrix} = \begin{pmatrix} x^0 \\ x^1 \end{pmatrix} \rightarrow \begin{pmatrix} x'^0 \\ x'^1 \end{pmatrix} = \begin{pmatrix} \cosh \xi & \sinh \xi \\ \sinh \xi & \cosh \xi \end{pmatrix} \begin{pmatrix} x^0 \\ x^1 \end{pmatrix}$$

$$\cosh \xi = \gamma = \frac{1}{\sqrt{1 - v^2}}$$

Special: parameter ξ or v is constant, e.g, inertial system invariance: *Global* conservation of E and p (still action at a distance!)

General: parameter $\xi(t, x)$ or $v(t, x)$ is constant, e.g, accelerated system invariance: *Local* conservation of E and p

Inestability of binary particle systems




Gravitational wave discovery by LIGO



credits: science.org

Noether's paradigm → Lagrangian formulation of classical field theory

$U(1)$: From special θ to general $\theta(t, x)$



What is a *particle wavicle*?

<https://www.quantamagazine.org/what-is-a-particle-20201112/>



Is a “Quantum Excitation of a Field”



Is a “Irreducible Representation of a Group”



H




Interacción → Fuerza = $\Delta p/\Delta t$

Introducción


Campos de materia →

Campos de radiación ~~~~~




[doi:10.1088/1742-6596/1287/1/012045](https://doi.org/10.1088/1742-6596/1287/1/012045)

a)



Emisión

b)





absorción

Interacción → Fuerza = $\Delta p/\Delta t$

Introducción


Campos de materia →

Campos de radiación ↗




[doi:10.1088/1742-6596/1287/1/012045](https://doi.org/10.1088/1742-6596/1287/1/012045)

a)




b)



$$e^- \rightarrow e^{-iEt}$$


$$e^+ \rightarrow e^{-iE(-t)}$$

Single charge



$$(e^-)^* = e^+$$

↗ fotón neutro



Emisión

absorción

Under a general Lorentz transformation we have for a **pure upperscript** 4-vector

$$A^\mu(x) \rightarrow A'^\mu(x) = \Lambda^\mu{}_\nu A^\nu(\Lambda^{-1}x). \quad (8)$$

A **pure underscript** 4-vector is

$$\partial_\mu = \frac{\partial}{\partial x^\mu} = \left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right) = (\partial_0, \nabla). \quad (9)$$

From

$$\frac{1}{x'^\mu} = (\Lambda^{-1})^\nu{}_\mu \frac{1}{x^\nu}, \quad (10)$$

the transformation properties for a $\partial_\mu = \partial/\partial x^\mu$, are

$$\partial'_\mu = \partial_\nu (\Lambda^{-1})^\nu{}_\mu. \quad (11)$$

In this way, the invariant scalar product between the 4-vector field and the four-gradient is just

$$\partial_\mu A^\mu \rightarrow \partial'_\mu A'^\mu = \partial_\mu A^\mu . \quad (12)$$

Photon: Representation of the Poincaré Group which transform as a vector under $\text{SO}(1, 3)$

Name	Symbol	$\text{SO}(1, 3)$
Photon	A^μ	$\Lambda^\mu{}_\nu A^\nu$
4-gradient	∂_μ	$\partial_\nu (\Lambda^{-1})^\nu{}_\mu$

Table 1: Scalar products: $\partial_\mu A^\mu$, $A^\nu A_\nu$, $\partial_\mu \partial^\mu$

Name	Symbol	$\text{SU}(N)$
scalar N -plet	Ψ	$U\Psi$
scalar anti- N -plet	Ψ^\dagger	$\Psi^\dagger U^\dagger$

Table 2: Scalar products: $\Psi^\dagger \Psi$

Name	Symbol	$\text{SL}(2, C)$	$U(1)_Q$
e_L : electron left	ξ_α	$S_\alpha^\beta \xi_\beta$	$e^{i\theta} \xi_\alpha$
$(e_L)^\dagger$: positron right	$(\xi_\alpha)^\dagger = \xi_{\dot{\alpha}}^\dagger$	$\xi_{\dot{\beta}}^\dagger [S^\dagger]_{\dot{\alpha}}^{\dot{\beta}}$	$\xi_{\dot{\alpha}}^\dagger e^{-i\theta}$
e_R : electron right	$(\eta^\alpha)^\dagger = \eta^\dagger{}^{\dot{\alpha}}$	$[(S^{-1})^\dagger]_{\dot{\beta}}^{\dot{\alpha}} \eta^\dagger{}^{\dot{\beta}}$	$e^{i\theta} \eta^\dagger{}^{\dot{\alpha}}$
$(e_R)^\dagger$: positron left	η^α	$\eta^\beta [S^{-1}]_\beta^\alpha$	$e^{-i\theta} \eta^\alpha$

Table 3: electron **left**: $\text{SL}(2, C) \times U(1)$ and positron **left**: $\text{SL}(2, C)^* \times U(1)$

Scalar products

- $\cancel{U(1)}$ Majorana scalars: $\xi^\alpha \xi_\alpha + \xi_{\dot{\alpha}}^\dagger \xi^{\dot{\alpha}}$, $\eta^\alpha \eta_\alpha + \eta_{\dot{\alpha}}^\dagger \eta^{\dot{\alpha}}$.
- Dirac scalar: $\eta^\alpha \xi_\alpha + \xi_{\dot{\alpha}}^\dagger \eta^{\dot{\alpha}}$.
- Tensor under subgroup $\text{SL}(2, C)$ but vector under $\text{SO}(1, 3)$: $S^{\dot{\alpha}}{}_{\dot{\beta}} \bar{\sigma}^{\mu}{}^{\dot{\beta}}{}^{\dot{\alpha}} S_\beta^\alpha = \Lambda^\mu{}_\nu \bar{\sigma}^\nu{}^{\dot{\alpha}}{}^{\alpha}$

$$\bar{\sigma}^\mu = (\sigma^0, -\boldsymbol{\sigma})$$

$$\sigma^\mu = (\sigma^0, \boldsymbol{\sigma}).$$

Name	Symbol	$\text{SL}(2, C)$	$U(1)$
e_L : electron left	ξ_α	$S_\alpha^\beta \xi_\beta$	$e^{i\theta} \xi_\alpha$
$(e_L)^\dagger$: positron right	$(\xi_\alpha)^\dagger = \xi_{\dot{\alpha}}^\dagger$	$\xi_{\dot{\beta}}^\dagger [S^\dagger]_{\dot{\alpha}}^{\dot{\beta}}$	$\xi_{\dot{\alpha}}^\dagger e^{-i\theta}$
e_R : electron right	$(\eta^\alpha)^\dagger = \eta^\dagger{}^{\dot{\alpha}}$	$[(S^{-1})^\dagger]_{\dot{\beta}}^{\dot{\alpha}} \eta^\dagger{}^{\dot{\beta}}$	$e^{i\theta} \eta^\dagger{}^{\dot{\alpha}}$
$(e_R)^\dagger$: positron left	η^α	$\eta^\beta [S^{-1}]_\beta^\alpha$	$e^{-i\theta} \eta^\alpha$

Table 4: electron **left**: $\text{SL}(2, C) \times U(1)$ and positron **left**: $\text{SL}(2, C)^* \times U(1)$

General theory: QED $\rightarrow D_\mu = i\partial_\mu - ieA_\mu$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$



$$\xi^\alpha \rightarrow \xi'^\alpha = e^{i\theta(x)} \xi^\alpha \quad \eta_\alpha \rightarrow \eta'_\alpha = e^{-i\theta(x)} \eta_\alpha$$

$$D_\mu \xi^\alpha \rightarrow (D_\mu \xi^\alpha)' = e^{i\theta(x)} D_\mu \xi^\alpha \quad D_\mu \eta_\alpha \rightarrow (D_\mu \eta_\alpha)' = e^{-i\theta(x)} D_\mu \eta_\alpha$$

$$\mathcal{L} = i\xi_{\dot{\alpha}}^\dagger \bar{\sigma}^\mu{}^{\dot{\alpha}\alpha} D_\mu \xi_\alpha + i\eta^\alpha \sigma^\mu_{\alpha\dot{\alpha}} D_\mu \eta^\dagger{}^{\dot{\alpha}} - m \left(\eta^\alpha \xi_\alpha + \xi_{\dot{\alpha}}^\dagger \eta^\dagger{}^{\dot{\alpha}} \right) - \frac{1}{4} F^{\mu\nu} F_{\mu\nu}$$

Name	Symbol	$\text{SL}(2, C)$	$U(1)$
e_L : electron left	ξ_α	$S_\alpha^\beta \xi_\beta$	$e^{i\theta} \xi_\alpha$
$(e_L)^\dagger$: positron right	$(\xi_\alpha)^\dagger = \xi_{\dot{\alpha}}^\dagger$	$\xi_{\dot{\beta}}^\dagger [S^\dagger]_{\dot{\alpha}}^{\dot{\beta}}$	$\xi_{\dot{\alpha}}^\dagger e^{-i\theta}$
e_R : electron right	$(\eta^\alpha)^\dagger = \eta^\dagger{}^{\dot{\alpha}}$	$[(S^{-1})^\dagger]_{\dot{\beta}}^{\dot{\alpha}} \eta^\dagger{}^{\dot{\beta}}$	$e^{i\theta} \eta^\dagger{}^{\dot{\alpha}}$
$(e_R)^\dagger$: positron left	η^α	$\eta^\beta [S^{-1}]_\beta^\alpha$	$e^{-i\theta} \eta^\alpha$

Table 4: electron **left**: $\text{SL}(2, C) \times U(1)$ and positron **left**: $\text{SL}(2, C)^* \times U(1)$

General theory: QED $\rightarrow D_\mu = i\partial_\mu - ieA_\mu$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

Dirac spinor

$$\begin{aligned} \xi^\alpha &\rightarrow \xi'^\alpha = e^{i\theta(x)} \xi^\alpha & \eta_\alpha &\rightarrow \eta'_\alpha = e^{-i\theta(x)} \eta_\alpha \\ D_\mu \xi^\alpha &\rightarrow (D_\mu \xi^\alpha)' = e^{i\theta(x)} D_\mu \xi^\alpha & D_\mu \eta_\alpha &\rightarrow (D_\mu \eta_\alpha)' = e^{-i\theta(x)} D_\mu \eta_\alpha \end{aligned}$$

$$\begin{aligned} \mathcal{L} &= i\xi_{\dot{\alpha}}^\dagger \bar{\sigma}^\mu{}^{\dot{\alpha}\alpha} D_\mu \xi_\alpha + i\eta^\alpha \sigma^\mu_{\alpha\dot{\alpha}} D_\mu \eta^\dagger{}^{\dot{\alpha}} - m \left(\eta^\alpha \xi_\alpha + \xi_{\dot{\alpha}}^\dagger \eta^\dagger{}^{\dot{\alpha}} \right) - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} \\ \mathcal{L} &= i\bar{\psi} \gamma^\mu D_\mu \psi - m \bar{\psi} \psi - \frac{1}{4} F^{\mu\nu} F_{\mu\nu}. \end{aligned}$$


$$\begin{aligned} \psi &= \begin{pmatrix} e_L \\ e_R \end{pmatrix} \\ \gamma^\mu &= \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix} \\ \bar{\psi} &= \psi^\dagger \gamma^0. \end{aligned}$$

$$\text{SU}(2)_L \rightarrow D_\mu = \mathbf{1} \partial_\mu - ig_2 \frac{\tau_i}{2} W_\mu^i \quad : 17 \text{ years later... (stages of grief} \rightarrow 1967)$$


Not mass, not charge

Field	Lorentz	$\text{SU}(2)_L$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	2

Denial



$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \tfrac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu}$$




$SU(2)_L \times U(1)_Y \rightarrow D_\mu = \mathbf{1} \partial_\mu - ig_2 \frac{\tau_i}{2} W_\mu^i - ig_1 B_\mu$: 17 years later... (stages of grief \rightarrow 1967)


Not mass, hypercharge,

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	2	-1/2

Denial




$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu}$$




Not mass, hypercharge, not Dirac

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	$\mathbf{2}$	$-1/2$
$(e_R)^\dagger$	η^α	$\mathbf{1}$	-1

Denial



$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R$$




$SU(2)_L \times U(1)_Y \rightarrow D_\mu = \mathbf{1} \partial_\mu - ig_2 \frac{\tau_i}{2} W_\mu^i - ig_1 B_\mu : 17$ years later... (stages of grief $\rightarrow 1967$)


Higgs mechanism: tachyonic mass $\mu^2 < 0$, and condensate

Contempt

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	2	$-1/2$
$(e_R)^\dagger$	η^α	1	-1
$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \left[\frac{H(x) + v}{\sqrt{2}} \right] \exp \left[i \frac{\tau^i}{2} G_i(x) \right]$	-	2	$1/2$



$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \tfrac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \tfrac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R + (e_R)^\dagger \Phi^\dagger L - (D^\mu \Phi)^\dagger D_\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2$$



Higgs mechanism: tachyonic mass $\mu^2 < 0$, and condensate

Field	Lorentz	$SU(2)_L$	$U(1)_Y$	Contempt
$L = \begin{pmatrix} \nu_L \\ e_L \\ (e_R)^\dagger \end{pmatrix}$	ξ_α	2	$-1/2$	
$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \left[\frac{H(x) + v}{\sqrt{2}} \right] \exp \left[i \frac{\tau^i}{2} G_i(x) \right]$	η^α	1	-1	
	-	2	$1/2$	$\begin{pmatrix} W_\mu^3 \\ B_\mu \end{pmatrix} = \begin{pmatrix} \cos \theta_W & \sin \theta_W \\ -\sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix},$

$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R + (e_R)^\dagger \Phi^\dagger L - (D^\mu \Phi)^\dagger D_\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2$$

$$\Phi \rightarrow \Phi' = \exp \left[i \frac{\tau^i}{2} \theta_i(x) \right] \Phi = \frac{1}{\sqrt{2}} [H(x) + v]$$

$$\mathcal{L} = i\bar{\psi} \gamma^\mu \partial_\mu \psi - m_e \bar{\psi} \psi - i(\nu_L)^\dagger \bar{\sigma}^\mu \partial_\mu \nu_L + \frac{1}{2} \partial^\mu H \partial_\mu H + \frac{e}{\cos \theta_W \sin \theta_W} \bar{\nu}_L \nu_L Z_\mu + \dots$$

$$-\frac{1}{2} m_H^2 H^2 \left(1 + \frac{H}{v} + \frac{H^2}{4v^2} \right) + \left(m_W^2 W^\mu - W_\mu^+ + \frac{1}{2} m_Z^2 Z^\mu Z_\mu \right) \left(1 + 2 \frac{H}{v} + \frac{H^2}{v^2} \right) + \frac{m_e}{v} \bar{\psi} \psi H$$

Z and W phenomenology and discovery

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	2	$-1/2$
$(e_R)^\dagger$	η^α	1	-1
$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \left[\frac{H(x) + v}{\sqrt{2}} \right] \exp \left[i \frac{\tau^i}{2} G_i(x) \right]$	-	2	$1/2$

$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R + (e_R)^\dagger \Phi^\dagger L - (D^\mu \Phi)^\dagger D_\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2$$

$$\Phi \rightarrow \Phi' = \exp \left[i \frac{\tau^i}{2} \theta_i(x) \right] \Phi = \frac{1}{\sqrt{2}} [H(x) + v]$$

$$\mathcal{L} = \bar{\psi} \gamma^\mu \partial_\mu \psi - m_e \bar{\psi} \psi - i(\nu_L)^\dagger \bar{\sigma}^\mu \partial_\mu \nu_L + \frac{1}{2} \partial^\mu H \partial_\mu H + \frac{e}{\cos \theta_W \sin \theta_W} \bar{\nu}_L \nu_L Z_\mu + \dots$$

$$-\frac{1}{2} m_H^2 H^2 \left(1 + \frac{H}{v} + \frac{H^2}{4v^2} \right) + \left(m_W^2 W^\mu - W_\mu^+ + \frac{1}{2} m_Z^2 Z^\mu Z_\mu \right) \left(1 + 2 \frac{H}{v} + \frac{H^2}{v^2} \right) + \frac{m_e}{v} \bar{\psi} \psi H$$

Bargaining



$SU(2)_L \times U(1)_Y \rightarrow D_\mu = \mathbf{1} \partial_\mu - ig_2 \frac{\tau_i}{2} W_\mu^i - ig_1 B_\mu : 32$ years later... (stages of grief $\rightarrow 1982$)

Hierarchy problem

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	$\mathbf{2}$	$-1/2$
$(e_R)^\dagger$	η^α	$\mathbf{1}$	-1
$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \left[\frac{H(x) + v}{\sqrt{2}} \right] \exp \left[i \frac{\tau^i}{2} G_i(x) \right]$	-	$\mathbf{2}$	$1/2$


$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R + (e_R)^\dagger \Phi^\dagger L - (D^\mu \Phi)^\dagger D_\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2$$

$$\Phi \rightarrow \Phi' = \exp \left[i \frac{\tau^i}{2} \theta_i(x) \right] \Phi = \frac{1}{\sqrt{2}} [H(x) + v]$$

$$\mathcal{L} = i \bar{\psi} \gamma^\mu \partial_\mu \psi - m_e \bar{\psi} \psi - i(\nu_L)^\dagger \bar{\sigma}^\mu \partial_\mu \nu_L + \frac{1}{2} \partial^\mu H \partial_\mu H + \frac{e}{\cos \theta_W \sin \theta_W} \bar{\nu}_L \nu_L Z_\mu + \dots$$

$$-\frac{1}{2} m_H^2 H^2 \left(1 + \frac{H}{v} + \frac{H^2}{4v^2} \right) + \left(m_W^2 W^{\mu-} W_\mu^+ + \frac{1}{2} m_Z^2 Z^\mu Z_\mu \right) \left(1 + 2 \frac{H}{v} + \frac{H^2}{v^2} \right) + \frac{m_e}{v} \bar{\psi} \psi H$$

Depression



credit: quantumdiaries.org

$SU(2)_L \times U(1)_Y \rightarrow D_\mu = 1\partial_\mu - ig_2 \frac{\tau_i}{2} W_\mu^i - ig_1 B_\mu$: 62 years later... (stages of grief \rightarrow 2012)

Higgs discovery!

Field	Lorentz	$SU(2)_L$	$U(1)_Y$
$L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	ξ_α	2	$-1/2$
$(e_R)^\dagger$	η^α	1	-1
$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \left[\frac{H(x) + v}{\sqrt{2}} \right] \exp \left[i \frac{\tau^i}{2} G_i(x) \right]$	-	2	$1/2$


$$\mathcal{L} = i(L)^\dagger \bar{\sigma}^\mu D_\mu L - \frac{1}{4} W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} - i(e_R)^\dagger \sigma^\mu D_\mu e_R + (e_R)^\dagger \Phi^\dagger L - (D^\mu \Phi)^\dagger D_\mu \Phi - \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2$$





$$\Phi \rightarrow \Phi' = \exp \left[i \frac{\tau^i}{2} \theta_i(x) \right] \Phi = \frac{1}{\sqrt{2}} [H(x) + v]$$

$$\mathcal{L} = i\bar{\psi} \gamma^\mu \partial_\mu \psi - m_e \bar{\psi} \psi - i(\nu_L)^\dagger \bar{\sigma}^\mu \partial_\mu \nu_L + \frac{1}{2} \partial^\mu H \partial_\mu H + \frac{e}{\cos \theta_W \sin \theta_W} \bar{\nu}_L \nu_L Z_\mu + \dots$$

$$-\frac{1}{2} m_H^2 H^2 \left(1 + \frac{H}{v} + \frac{H^2}{4v^2} \right) + \left(m_W^2 W^{\mu-} W_\mu^+ + \frac{1}{2} m_Z^2 Z^\mu Z_\mu \right) \left(1 + 2 \frac{H}{v} + \frac{H^2}{v^2} \right) + \frac{m_e}{v} \bar{\psi} \psi H$$

Acceptance





en.wikipedia.org

Jacobus Kapteyn - Wikipedia

Prof Jacobus Cornelius Kapteyn FRS FRSE LLD (19 January 1851 – 18 June 1922) was a Dutch astronomer. He carried out extensive studies of the Milky Way and was the discoverer of evidence for galactic rotation. Kapteyn was also among the first to suggest ...







en.wikipedia.org


Fritz Zwicky - Wikipedia





Fritz Zwicky (; German: ; February 14, 1898 – February 8, 1974) was a Swiss astronomer. He worked most of his life at the California Institute of Technology in the United States of America, where he made many important contributions in theoretical and o...







$V(\phi)$










Particle physics without hep-ex in Colombia in lens.org



High energy physics in Colombia from inspirehep.org

- Andes U., Bogota (2016) (D0,CMS)
- Antonio Narino U. (1304) (ATLAS)
- Colombia, U. Natl. (949) (LHCb)
- [Antioquia U. \(596\)](#) (CMS,DUNE,LIGO)
- Valle U., Cali (146)
- Santander Industrial U. (144) (AUGER)
- U. Atlantico, Barranquilla (62) (NOVA,DUNE)
- [Narino U. \(44\)](#)
- Natl. U. of Columbia, Medellin (43)
- Medellin U. (43) (AUGER,DUNE)
- [UPTC, Tunja \(36\)](#)
- [Santiago de Cali U. \(30\)](#)
- Tolima U. (29)
- Escuela de Ingenieria de Antioquia (27) (DUNE)
- Pereira Tech. U. (25)
- [Inst. Tech. Metro., Medellin \(19\)](#)
- [Pamplona U. \(16\)](#)
- Distrital U., Bogota (16)
- Bolivar Tech. U. (15) (NOVA,DUNE)
- U. Sergio Arboleda, Bogota (14)

<https://inspirehep.net/institutions?q=colombia>




@conhep



November 28, 2022 to December 2, 2022
America/Bogota timezone

Villa de Leyva



May 31, 2022 to June 1, 2022
America/Bogota timezone

6th: Uniandes



Jul 27 – 29, 2022
America/Bogota timezone

2nd: EIA



Jul 27 – 29, 2022
America/Bogota timezone

2nd: EIA





LATIN AMERICAN ASSOCIATION FOR HIGH ENERGY, COSMOLOGY AND ASTROPARTICLE PHYSICS

LATIN AMERICAN STRATEGY FOR RESEARCH INFRASTRUCTURE - LASF4RI

Latin American Strategy for
Research Infrastructures
for High Energy, Cosmology,
Astroparticle Physics
LASF4RI for HECAP

arXiv:2104.06852v1 [hep-ex] 14 Apr 2021

LATIN AMERICAN HECAP PHYSICS BRIEFING BOOK

Preparatory Group

Hiroaki Aihara - University of Tokyo	Alfredo Aranda - University of Colima
Reina Camacho Toro - LPNHE/CNRS	Mauro Cambiaso - Universidad Andrés Bello
Marcela Carena - Fermilab/U. of Chicago	Edgar Carrera - Universidad San Francisco de Quito
Juan Carlos D'Olivo - UNAM	Alberto Gago - Pontificia Universidad Católica del Perú
Thiago Goncalves - Valongo Observatory	Gerardo Herrera - CINVESTAV
Diana López Náir - DF/IFIBA UBA-CONICET	Marta Losada - NYUAD
Jorge Molina - Universidad Nacional de Asunción	Maritijn Mulders - CERN
Diego Restrepo - Universidad de Antioquia.	Rogério Rosenfeld - IFT-UNESP & ICTP-SAIFR
Arturo Sánchez - ICTP/INFN/ U. of Udine	Federico Sánchez - U. Nacional de San Martín
Marcelle Soules-Santos - U. Michigan	Martín Subietz - U. Mayor de San Andrés
Héctor Wahnborg - U. Nacional de la Plata	Harold Yépez Ramírez - YTU
Alfonso Zerwekh - U. Técnicas Fedrico Santa María	