

# Dark matter from SM gauge extensions

## with neutrino masses

---



Diego Restrepo

July 23, 2019 - PPC2019 - Cartagena

Instituto de Física  
Universidad de Antioquia  
Phenomenology Group  
<http://gfif.udea.edu.co>



Focus on

In collaboration with

M. Hirsch (IFIC), C. Álvarez (UTFSM), A. Flórez (UniAndes), B. Dutta (Texas A&M), C. Yaguna (UPTC), J. Calle, O. Zapata, A. Rivera (UdeA), W. Tangarife (Loyola University Chicago)

## Hidden sectors

---



+

$$\frac{1}{\Lambda} L \cdot H L \cdot H$$


$$\frac{1}{\Lambda} (\nu_R)^\dagger L \cdot H S$$



+

$$\frac{1}{\Lambda} L \cdot H L \cdot H$$

$$\frac{1}{\Lambda} (\nu_R)^\dagger L \cdot H S$$



# Dark matter and unification

---

# Unification: $SO(10)$

$$16_{F_i} = \begin{pmatrix} u_R^\dagger \\ u_R^\dagger \\ u_R^\dagger \\ u_L \\ u_L \\ u_L \\ d_L \\ d_L \\ d_L \\ d_R^\dagger \\ d_R^\dagger \\ d_R^\dagger \\ \nu_L \\ e_L \\ e_R^\dagger \\ N \end{pmatrix}_i$$

$$\Rightarrow \mathcal{L}_{SM} \supset h \, 16_F \times 16_F \times 10_S + \text{h.c.}$$



# Not-susy $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_2$

Standard Model:  $Z_2$ -even

Fermions:  $16_F$


Scalars:  $10_H, 45_H \dots$

New  $Z_2$ -odd particles

$10_F, 45_F, \dots$

$16_H, \dots$

Lightest Odd Particle (LOP) may be a suitable dark matter candidate, and can improve gauge coupling unification



# Not-susy $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_2$

Standard Model:  $Z_2$ -even

Fermions:  $16_F$

Scalars:  $10_H, 45_H \dots$

New  $Z_2$ -odd particles

$10_F, 45_F, \dots$

$16_H, \dots$

Lightest Odd Particle (LOP) may be a suitable dark matter candidate, and can improve gauge coupling unification

	fermions	scalars
$SU(2)_L \times U(1)_Y$ representation	even $SO(10)$ representations	odd $SO(10)$ representations
$1_0$	45, 54, 126, 210	16, 144
$2_{\pm 1/2}$	10, 120, 126, 210, 210'	16, 144
$3_0$	45, 54, 210	144

$SU(3)_C : 3 (T), 6, 8 (\Lambda)$

$$m_{3_0} = 2.7 \text{ TeV}, \quad m_{\Lambda} \sim 10^{10} \text{ TeV}, \quad m_{\text{GUT}} \sim 10^{16} \text{ GeV}.$$



# Not-susy $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_2$

Standard Model:  $Z_2$ -even

Fermions:  $16_F$

Scalars:  $10_H, 45_H \dots$

New  $Z_2$ -odd particles

$10_F, 45_F, \dots$

$16_H, \dots$

Lightest Odd Particle (LOP) may be a suitable dark matter candidate, and can improve gauge coupling unification

	fermions	scalars
$SU(2)_L \times U(1)_Y$ representation	even $SO(10)$ representations	odd $SO(10)$ representations
$1_0$	$45, 54, 126, 210$	$16, 144$
$2_{\pm 1/2}$	$10, 120, 126, 210, 210'$	$16, 144$
$3_0$	$45, 54, 210$	$144$

$SU(3)_C : 3 (T), 6, 8 (\Lambda)$

Split-SUSY like

arXiv:1509.06313 (C. Arbelaez, R. Longas, D.R, O. Zapata)

# Not-susy $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_2$

Standard Model:  $Z_2$ -even

Fermions:  $16_F$

Scalars:  $10_H, 45_H \dots$

New  $Z_2$ -odd particles

$10_F, 45_F, \dots$

$16_H, \dots$

Lightest Odd Particle (LOP) may be a suitable dark matter candidate, and can improve gauge coupling unification

	fermions		scalars
$SU(2)_L \times U(1)_Y$ representation	even $SO(10)$ representations	odd $SO(10)$ representations	
$1_0$	$45, 54, 126, 210$	$16$	144
$2_{\pm 1/2}$ $2_{1/2}^S$	$10, 120, 126, 210, 210'$		16, 144
$3_0$	$45, 54, 210$		144

$SU(3)_C : 3 (T), 6, 8 (\Lambda)$

Radiative hybrid seesaw (Parida 1106.4137) or 1509.06313

Partial Split-SUSY-like spectrum: bino-higgsino-wino

+

↓

$10'_H$  with fermion DM or,  $16_H, \dots$  with scalar DM

# Not-susy $SO(10) \rightarrow SU(5) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_2$

Standard Model:  $Z_2$ -even

Fermions:  $16_F$

Scalars:  $10_H, 45_H \dots$

New  $Z_2$ -odd particles

$10_F, 45_F, \dots$

$16_H, \dots$

Lightest Odd Particle (LOP) may be a suitable dark matter candidate, and can improve gauge coupling unification

	fermions	scalars
$SU(2)_L \times U(1)_Y$ representation	even $SO(10)$ representations	odd $SO(10)$ representations
$1_0$	45, 54, 126, 210	$16$ 144
$2_{\pm 1/2}$ $2_{1/2}^S$	$10$ , 120, 126, 210, 210'	16, 144
$3_0$	$45$ , 54, 210	144

$SU(3)_C$ :  $3$  (T), 6,  $8$  ( $\Lambda$ )

1509.06313

SUSY-like spectrum: bino-higgsino-wino

+

↓

$10'_H$  with fermion DM or,  $16_H, \dots$  with scalar DM

## Left-Right symmetric realization

---

# Singlet-doublet fermion dark matter

Field	Multiplicity	$3_c 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$\Phi$	1	(1, 2, 2, 0)	0	10
$\chi, \chi^c$	1	(1, 2, 2, 0)	1/2	10
$N$	1	(1, 1, 1, 0)	1/2	45

**Table 1:** The relevant part of the field content. Note that, the two fermion doublets  $\chi$  and  $\chi^c$  come from an only fermionic LR bidoublet. In the third column the relevant fields are characterized by their  $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$  quantum numbers while their SO(10) origin is specified in the fourth column.

$m_{LR}$ (GeV)	$3_c 2_L 2_R 1_{B-L}$	$m_G$ (GeV)
$2 \times 10^3$	$\Phi_{1,2,2,0} + 2\Phi_{1,1,3,-2} + \Psi_{1,1,3,0} + \Phi_{1,1,3,0} + \Phi_{8,1,1,0}$	$1.65 \times 10^{16}$
$\vdots$	$\vdots$	$\vdots$

Table 2:  $\Delta_{L,R} = 2\Phi_{1,1,3,-2}$ .  $m_{LR}$  and  $m_G$  are given in GeV.

# Triplets

---

# Minimal Left-Right Symmetric Standard Model

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	$1/2$	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	$1/2$	16
$L$	3	$(1, 2, 1, -1)$	$1/2$	16
$L^c$	3	$(1, 1, 2, +1)$	$1/2$	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126



## Left-singlet right-triplet DM

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	1/2	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	1/2	16
$L$	3	$(1, 2, 1, -1)$	1/2	16
$L^c$	3	$(1, 1, 2, +1)$	1/2	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126
$\Psi_{1130}$	1	$(1, 1, 3, 0)$	1/2	45

$$\Omega h^2 = 0.1199 \pm 0.0027 \text{ at } 3\sigma$$




Figure 1: Proper relic density scan:  $0.5 < v_R/\text{TeV} < 50$

## Mixed Left-singlet right-triplet DM

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	1/2	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	1/2	16
$L$	3	$(1, 2, 1, -1)$	1/2	16
$L^c$	3	$(1, 1, 2, +1)$	1/2	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126
$\Psi_{1130}$	1	$(1, 1, 3, 0)$	1/2	45

## Mixed Left-singlet right-triplet DM

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	1/2	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	1/2	16
$L$	3	$(1, 2, 1, -1)$	1/2	16
$L^c$	3	$(1, 1, 2, +1)$	1/2	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126
$\Psi_{1130}$	1	$(1, 1, 3, 0)$	1/2	45
$\Psi_{1132}$	1	$(1, 1, 3, 2)$	1/2	126
$\Psi_{113-2}$	1	$(1, 1, 3, -2)$	1/2	$\overline{126}$

$$\Psi_{1132} = \begin{pmatrix} \Psi^+/\sqrt{2} & \Psi^{++} \\ \Psi^0 & -\Psi^+/\sqrt{2} \end{pmatrix}, \quad \bar{\Psi}_{113-2} = \begin{pmatrix} \Psi^-/\sqrt{2} & \bar{\Psi}^0 \\ \Psi^{--} & -\Psi^-/\sqrt{2} \end{pmatrix}. \quad (1)$$

$$\begin{aligned} L \supset & M_{11} \text{Tr}(\Psi_{1130} \Psi_{1130}) + M_{23} \text{Tr}(\Psi_{1132} \bar{\Psi}_{113-2}) \\ & + \lambda_{13} \text{Tr}(\Delta_R \bar{\Psi}_{113-2} \Psi_{1130}) + \lambda_{12} \text{Tr}(\Delta_R^\dagger \Psi_{1132} \Psi_{1130}), \end{aligned} \quad (2)$$

$$\tan \gamma = \frac{\lambda_{13}}{\lambda_{12}}, \quad \lambda = \sqrt{\lambda_{12}^2 + \lambda_{13}^2}. \quad (3)$$

Blind spot at

$$M_{23} \sin 2\gamma - M_{\text{DM}} = 0 \quad (4)$$




Figure 2:  $M_{11} = 50$  TeV  $2.7 < M_{23}/\text{TeV} < 3.1$  (Right:  $\tan \gamma > 5$ )

$$\Omega h^2 = 0.1199 \pm 0.0027 \text{ at } 3\sigma$$





Figure 3:

$$\Omega h^2 = 0.1199 \pm 0.0027 \text{ at } 3\sigma$$



**Figure 3:** Proper relic density scan:  $v_R : [2, 50]$  TeV,  $M_{23} : [0.2, 50]$  TeV,  $M_{11} : 50$  TeV,  $\tan \gamma = -1$  and  $\lambda = 0.14$ .



## Direct detection cross section





Figure 4:  $v_R : [2, 50]$  TeV,  $M_{23} : [0.2, 50]$  TeV,  $M_{11} : 50$  TeV,  $\tan \gamma = -1$  and  $\lambda = 0.14$ .


# Unification

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	1/2	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	1/2	16
$L$	3	$(1, 2, 1, -1)$	1/2	16
$L^c$	3	$(1, 1, 2, +1)$	1/2	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126
$\Psi_{1130}$	1	$(1, 1, 3, 0)$	1/2	45
$\Psi_{1132}$	1	$(1, 1, 3, 2)$	1/2	126
$\Psi_{113-2}$	1	$(1, 1, 3, -2)$	1/2	$\overline{126}$

# Unification

Field	Multiplicity	$3_C 2_L 2_R 1_{B-L}$	Spin	SO(10) origin
$Q$	3	$(3, 2, 1, +\frac{1}{3})$	1/2	16
$Q^c$	3	$(\bar{3}, 1, 2, -\frac{1}{3})$	1/2	16
$L$	3	$(1, 2, 1, -1)$	1/2	16
$L^c$	3	$(1, 1, 2, +1)$	1/2	16
$\Phi$	1	$(1, 2, 2, 0)$	0	10
$\Delta_R$	1	$(1, 1, 3, -2)$	0	126
$\Psi_{1130}$	1	$(1, 1, 3, 0)$	1/2	45
$\Psi_{1132}$	1	$(1, 1, 3, 2)$	1/2	126
$\Psi_{113-2}$	1	$(1, 1, 3, -2)$	1/2	$\overline{126}$
$\Psi_{1310}$	1	$(1, 3, 1, 0)$	1/2	45
$\Psi_{8110}$	1	$(1, 1, 8, 0)$	1/2	45
$\Psi_{321\frac{1}{3}}$	1	$(3, 2, 1, 1/3)$	1/2	16
$\Psi_{321-\frac{1}{3}}$	1	$(1, 2, 3, -1/3)$	1/2	$\overline{16}$

# Unification quality



In addition to accommodate usual simplified dark matter models, Left-right symmetric standard models have additional DM portals:

New  $\Delta_R$  portal for direct detection of left-singlet right-triplet mixed dark matter, in companion with left-singlets charged and doubly charged fermions.

Next: Search for them in compressed spectra scenarios at the LHC

## Dirac fermion dark matter


---

# Isosinglet dark matter candidate

$\psi$  as a isosinglet Dirac dark matter fermion charged under a local  $U(1)_X$  (SM) couples to a SM-singlet vector mediator  $X$  as

$$\mathcal{L}_{\text{int}} = -g_\psi \bar{\psi} \gamma^\mu \psi X_\mu - \sum_f g_f \bar{f} \gamma^\mu f X_\mu,$$


where  $f$  are the Standard Model fermions



# Isosinglet Dirac fermion dark matter model and Seesaw scale

Left Field	$U(1)_{B-L}$
$(\nu_{R1})^\dagger$	+1
$(\nu_{R2})^\dagger$	+1
$(\nu_{R2})^\dagger$	+1
$\psi_L$	$-r$
$(\psi_R)^\dagger$	$r$
$\phi$	2

$$\chi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}$$



Duerr et al: 1803.07462 [PRD]



## Singlet-Doublet Dirac Dark matter Model (SD<sup>3</sup>M)

---

The model extends the standard model (SM) particle content with Dirac Fermions: from SU(2) doublets of Weyl fermions:  $\Psi_L = (\Psi_L^0, \Psi_L^-)^T$ ,  $\widetilde{(\Psi_R)} = ((\Psi_R^-)^\dagger, -(\Psi_R^0)^\dagger)^T$  and singlet Weyl fermions  $\psi_{LR}$  that interact among themselves and with the SM fields


$$\mathcal{L} \supset \textcolor{red}{M}_\psi (\psi_R)^\dagger \psi_L + \textcolor{red}{M}_\Psi \widetilde{(\Psi_R)} \cdot \Psi_L + \textcolor{red}{y}_1 (\psi_R)^\dagger \Psi_L \cdot H + \textcolor{red}{y}_2 \widetilde{(\Psi_R)} \cdot \tilde{H} \psi_L + \text{h.c} \quad (5)$$

Four free parameters:


$$\textcolor{red}{M}_\psi, \textcolor{red}{M}_\Psi < 2 \text{ GeV}, \quad \textcolor{red}{y}_1, \textcolor{red}{y}_2 > 10^{-6} \quad (6)$$

Two neutral Dirac fermion eigenstates:


$$M = \begin{pmatrix} \textcolor{red}{M}_\psi & \textcolor{red}{y}_2 v / \sqrt{2} \\ \textcolor{red}{y}_1 v / \sqrt{2} & \textcolor{red}{M}_D \end{pmatrix}, \quad M_{\text{diag}} = \begin{pmatrix} M_{\chi_1} & 0 \\ 0 & M_{\chi_2} \end{pmatrix} = U_L^\dagger M U_R \quad (7)$$




Compressed spectra region




LUX - XENON1T - LZ



Compressed spectra region




LUX - XENON1T - LZ



# Neutrino masses

---

- Lepton number ( $L$ ) is an accidental discrete or Abelian symmetry of the standard model (SM).
- Without neutrino masses  $L_e$ ,  $L_\mu$ ,  $L_\tau$  are also conserved.
- The processes which violate individual  $L$  are called Lepton flavor violation (LFV) processes.
- All the neutrino mass models predict, to some extent, LFV processes
- Only models with Majorana neutrinos predict processes with total  $L = L_e + L_\mu + L_\tau$  violation, like **neutrino less doublet beta decay** (NLDBD).
- NLDBD is experimentally challenging, specially if there is a massless neutrino in the spectrum.



Total lepton number:  $L = L_e + L_\mu + L_\tau$

Majorana  $\cancel{U(1)}_L$

Field	$Z_2 \ (\omega^2 = 1)$
SM	1
$L$	$\omega$
$(e_R)^\dagger$	$\omega$
$(\nu_R)^\dagger$	$\omega$

$$\mathcal{L}_\nu = h_D (\nu_R)^\dagger L \cdot H + \textcolor{red}{M_R} \nu_R \nu_R + \text{h.c.}$$

$$h_D \sim \mathcal{O}(1)$$

Dirac  $U(1)_L$

Field	$Z_3 \ (\omega^3 = 1)$
SM	1
$L$	$\omega$
$(e_R)^\dagger$	$\omega^2$
$(\nu_R)^\dagger$	$\omega^2$

$$\mathcal{L}_\nu = h_D (\nu_R)^\dagger L \cdot H + \text{h.c.}$$

$$h_D \sim \mathcal{O}(10^{-11})$$



Total lepton number:  $L = L_e + L_\mu + L_\tau$

Majorana  $U(1)_L$

Field	$Z_2$ ( $\omega^2 = 1$ )
SM	1
$L$	$\omega$
$(e_R)^\dagger$	$\omega$
$(\nu_R)^\dagger$	$\omega$

$$\mathcal{L}_\nu = h_D (\nu_R)^\dagger L \cdot H + M_R \nu_R \nu_R + \text{h.c.}$$

$$h_D \sim \mathcal{O}(1)$$

Dirac  $U(1)_{B-L}$

Field	$Z_3$ ( $\omega^3 = 1$ )
SM	1
$L$	$\omega$
$(e_R)^\dagger$	$\omega^2$
$(\nu_R)^\dagger$	$\omega^2$

$$\mathcal{L}_\nu = h_D (\nu_R)^\dagger L \cdot H + \text{h.c.}$$

$$h_D \sim \mathcal{O}(10^{-11})$$

Explain smallness ala Peccei-Quinn:

$$U(1)_{B-L} \xrightarrow{\langle S \rangle} Z_N, \quad N \geq 3.$$

## Small Dirac neutrino masses

To explain the **smallness** of Dirac neutrino masses choose  $U(1)_{B-L}$  which:

- Forbids tree-level mass (TL) term (  $Y(H) = +1/2$  )

$$\begin{aligned}\mathcal{L}_{\text{T.L}} &= h_D \epsilon_{ab} (\nu_R)^\dagger L^a H^b + \text{h.c.} \\ &= h_D (\nu_R)^\dagger L \cdot H + \text{h.c.}\end{aligned}$$

# Small Dirac neutrino masses

To explain the **smallness** of Dirac neutrino masses choose  $U(1)_{B-L}$  which:

- Forbids tree-level mass (TL) term (  $Y(H) = +1/2$  )

$$\begin{aligned}\mathcal{L}_{T.L} &= h_D \epsilon_{ab} (\nu_R)^\dagger L^a H^b + \text{h.c.} \\ &= h_D (\nu_R)^\dagger L \cdot H + \text{h.c.}\end{aligned}$$

- Forbids Majorana term:  $\nu_R \nu_R$

# Small Dirac neutrino masses


To explain the **smallness** of Dirac neutrino masses choose  $U(1)_{B-L}$  which:

- Forbids tree-level mass (TL) term (  $Y(H) = +1/2$  )

$$\begin{aligned}\mathcal{L}_{T.L} &= h_D \epsilon_{ab} (\nu_R)^\dagger L^a H^b + \text{h.c} \\ &= h_D (\nu_R)^\dagger L \cdot H + \text{h.c}\end{aligned}$$

- Forbids Majorana term:  $\nu_R \nu_R$
- Realizes of the 5-dimension operator which conserves lepton number in  $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ :

$$\mathcal{L}_{5-D} = \frac{h_\nu}{\Lambda} (\nu_R)^\dagger L \cdot H S + \text{h.c}$$



# Small Dirac neutrino masses

To explain the **smallness** of Dirac neutrino masses choose  $U(1)_{B-L}$  which:


- Forbids tree-level mass (TL) term (  $Y(H) = +1/2$  )

$$\begin{aligned}\mathcal{L}_{T.L} &= h_D \epsilon_{ab} (\nu_R)^\dagger L^a H^b + \text{h.c.} \\ &= h_D (\nu_R)^\dagger L \cdot H + \text{h.c.}\end{aligned}$$

- Forbids Majorana term:  $\nu_R \nu_R$
- Realizes of the 5-dimension operator which conserves lepton number in  $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ :

$$\mathcal{L}_{5-D} = \frac{h_\nu}{\Lambda} (\nu_R)^\dagger L \cdot H S + \text{h.c.}$$

- Enhancement to the *effective number of degrees of freedom in the early Universe*  $\Delta N_{\text{eff}} = N_{\text{eff}} - N_{\text{eff}}^{\text{SM}}$  (see arXiv:1211.0186)



See E. Ma, Rahul Srivastava: arXiv:1411.5042 [PLB] for tree-level realization

From 1210.6350 and 1805.02025:  $\Delta N_{\text{eff}} = 3 (T_{\nu_R}/T_{\nu_L})^4$

$$\begin{aligned}\Gamma_{\nu_R}(T) &= n_{\nu_R}(T) \sum_f \langle \sigma_f(\nu_R \bar{\nu}_R \rightarrow f\bar{f}) v \rangle \\ &= \sum_f \frac{g_{\nu_R}^2}{n_{\nu_R}} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 q}{(2\pi)^3} f_{\nu_R}(p) f_{\nu_R}(q) \sigma_f(s) (1 - \cos \theta),\end{aligned}$$

$$s = 2pq(1 - \cos \theta), \quad f_{\nu_R}(k) = 1/(e^{k/T} + 1)$$

$$n_{\nu_R}(T) = g_{\nu_R} \int \frac{d^3 k}{(2\pi)^3} f_{\nu_R}(k), \quad \text{with } g_{\nu_R} = 2$$


$$\sigma_f(s) \simeq \frac{N_C^f (Q_{BL}^f)^2 Q^2 s}{12\pi} \left( \frac{g'}{M_{Z'}} \right)^4, \quad \text{In the limit } M_{Z'}^2 \gg s.$$

with three right-handed neutrinos, the Hubble parameter is

$$H(T) = \sqrt{\frac{4\pi^3 G_N [g(T) + 21/4]}{45}} T^2.$$

The right-handed neutrinos decouple when

$$\Gamma_{\nu_R}(T_{\text{dec}}^{\nu_R}) = H(T_{\text{dec}}^{\nu_R}).$$



A. Solaguren-Beascoa, M. C. Gonzalez-Garcia: arXiv:1210.6350 [PLB]

From 1210.6350 and **1805.02025**:  $\Delta N_{\text{eff}} = 3 (T_{\nu_R}/T_{\nu_L})^4$

$$\begin{aligned}\Gamma_{\nu_R}(T) &= n_{\nu_R}(T) \sum_f \langle \sigma_f(\nu_R \bar{\nu}_R \rightarrow f\bar{f}) v \rangle \\ &= \sum_f \frac{g_{\nu_R}^2}{n_{\nu_R}} \int \frac{d^3 p}{(2\pi)^3} \frac{d^3 q}{(2\pi)^3} f_{\nu_R}(p) f_{\nu_R}(q) \sigma_f(s) (1 - \cos \theta),\end{aligned}$$

$$s = 2pq(1 - \cos \theta),$$

$$f_{\nu_R}(k) = 1/(e^{k/T} + 1)$$

$$n_{\nu_R}(T) = g_{\nu_R} \int \frac{d^3 k}{(2\pi)^3} f_{\nu_R}(k), \quad \text{with } g_{\nu_R} = 2$$



$$\sigma_f(s) \simeq \frac{N_c^f (Q_{BL}^f)^2 Q^2 s}{12\pi} \left( \frac{g'}{M_{Z'}} \right)^4, \quad \text{In the limit } M_{Z'}^2 \gg s.$$

with three right-handed neutrinos, the Hubble parameter is

$$H(T) = \sqrt{\frac{4\pi^3 G_N [g(T) + 21/4]}{45}} T^2.$$

The right-handed neutrinos decouple when

$$\Gamma_{\nu_R}(T_{\text{dec}}^{\nu_R}) = H(T_{\text{dec}}^{\nu_R}).$$



Z.-L. Han, W. Wang: arXiv:1805.02025 [EJPC]

(also: Planck 1807.06209, Riess et al 1903.07603 )

One-loop realization of  $\mathcal{L}_{5-D}$  with  
total  $L$

---



Dirac neutrino masses


$U(1)_X$   
+

Dirac fermion dark matter



$L$

$\nu_R$



Dirac neutrino masses

$$\nu_R \nu_R$$

$$(\nu_R)^\dagger L H$$

$$\nu_R \psi_R$$


$$(\psi_L)^\dagger \nu_R$$


$U(1)_X$   
+  
Forbids

Dirac fermion dark matter

$$\psi_R \psi_R$$

$$\psi_L \psi_L$$





Dirac neutrino masses

$$\nu_R \nu_R$$

$$(\nu_R)^\dagger LH$$

$$\nu_R \psi_R$$

$$(\psi_L)^\dagger \nu_R$$

$$(\nu_R)^\dagger LH \textcolor{red}{S}$$

$X(L) \neq 0$



+  
Forbids

$$U(1)_{B-L}$$

$$\langle \overline{S} \rangle Z_N$$

$$N \neq 2$$


Dirac fermion dark matter

$$\psi_R \psi_R$$

$$\psi_L \psi_L$$

$$(\psi_R)^\dagger \psi_L$$


$$(\psi_R)^\dagger \psi_L S$$




Allows

normalized to  $X(L) = -1$


# One loop topologies $U(1)_{B-L} \oplus Z_2 \oplus Z_2$




T1-3-D




T1-3-E



T3-1-A




T1-2-A




T1-2-B

Chang-Yuan Yao and Gui-Jun Ding, arXiv:1802.05231 [PRD]

# One loop topologies $U(1)_{B-L}$ only!



T1-3-D




T1-3-E

$\psi_{L,R} \rightarrow$  Singlet fermions

$\Psi_{L,R} \rightarrow$  Vector-like doublet fermions


$\sigma \rightarrow$  Singlet scalar

$\eta \rightarrow$  Doublet scalar




T3-1-A

with J. Calle, C. Yaguna, and O. Zapata, arXiv:1812.05523 [PRD]




T1-2-A



T1-2-B

# One loop topologies $U(1)_{B-L}$ only! with J. Calle, C. Yaguna, and O. Zapata, arXiv:1812.05523 [PRD]




→ generalization to two and three loops: Shaikh Saad arXiv:1902.07259

T1-3-D

$\psi_{L,R} \rightarrow$  Singlet fermions (vector-like)

$\sigma \rightarrow$  Singlet scalar

$\eta \rightarrow$  Doublet scalar



T3-1-A


Fields: $f_i$	$(\nu_{R3})^\dagger$	$(\nu_{R2})^\dagger$	$(\nu_{R1})^\dagger$	$\psi_L$	$(\psi_R)^\dagger$	$S$
(A)	+4	+4	-5	-r	r	+3

Anomaly cancellation conditions

$$\sum_i f_i = 3$$

$$\sum_i f_i^3 = 3$$

# One loop topologies $U(1)_{B-L}$ only! with J. Calle, C. Yaguna, and O. Zapata, arXiv:1812.05523 [PRD]





Fields: $f_i$	$(\nu_{R3})^\dagger$	$(\nu_{R2})^\dagger$	$(\nu_{R1})^\dagger$	$\psi_L$	$(\psi_R)^\dagger$	$S$
(A)	+4	+4	-5	-r	r	+3
(B)	$+\frac{8}{5}$	$+\frac{8}{5}$	$+\frac{2}{5}$	$\frac{7}{5}$	$-\frac{10}{5}$	$+\frac{3}{5}$

$\psi_{L,R} \rightarrow$  Singlet fermions (quiral)

$\Psi_{L,R} \rightarrow$  Vector-like doublet fermions

$\sigma \rightarrow$  Singlet scalar

$\eta \rightarrow$  Doublet scalar




Anomaly cancellation conditions

$$\sum_i f_i = 3$$

$$\sum_i f_i^3 = 3$$



$\psi_{L,R} \rightarrow$  Singlet fermions (quiral)  
 $\Psi_{L,R} \rightarrow$  Vector-like doublet fermions : 10/5  
 $\sigma \rightarrow$  Singlet scalar : 15/5



Fields: $f_i$	$(\nu_{R3})^\dagger$	$(\nu_{R2})^\dagger$	$(\nu_{R1})^\dagger$	$\psi_L$	$(\psi_R)^\dagger$	$S$
(A)	+4	+4	-5	-r	r	+3
(B)	$+\frac{8}{5}$	$+\frac{8}{5}$	$+\frac{2}{5}$	$\frac{7}{5}$	$-\frac{10}{5}$	$+\frac{3}{5}$

Anomaly cancellation conditions



$$\sum_i f_i = 3$$

$$\sum_i f_i^3 = 3$$


$$M_\psi = h_1 \langle S \rangle, y_2 = 0:$$

$$\mathcal{L} = \mathcal{L}_{\text{SD}^3\text{M}} + h_3^{ia} \widetilde{(\Psi_R)} \cdot L_i \sigma_a + h_2^{\beta a} (\nu_{R\beta})^\dagger \psi_L \sigma_a^* - V(\sigma_a, S, H).$$


with A.F Rivera, W. Tangarife, arXiv:19nn.nnnnn



# Spin independent (SI) direct detection cross section



Decoupled  $Z'$  limit



Vector SI (blue points) and scalar SI (green points)

A single  $U(1)$  symmetry to explain both the smallness of Dirac neutrino masses and the stability of Dirac fermion dark matter

A single  $U(1)$  symmetry to explain both the smallness of Dirac neutrino masses and the stability of Dirac fermion dark matter

## Dirac neutrino masses and DM

- Spontaneously broken  $U(1)_{B-L}$  generates a radiative Dirac neutrino masses
- A remnant symmetry makes the lightest field circulating the loop stable and good dark matter candidate.
- For T1-2-A: Either Singlet Doublet Dirac Dark Matter or Singlet Scalar Dark Matter with extra scalar and vector portal
- With relaxed direct detection constraints

Thanks!