

Dirac dark matter, neutrino masses, and dark baryogenesis



Diego Restrepo

Instituto de Física
Universidad de Antioquia
Phenomenology Group
<http://gfif.udea.edu.co>



Focus on

[arXiv:2205.05762](https://arxiv.org/abs/2205.05762) [PRD]

In collaboration with

Andrés Rivera (UdeA), Walter Tangarife (Loyola University Chicago)

Dark sectors







Local $U(1)_\mathcal{X}$

$$\mathcal{L} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + i \sum_i \chi_i^\dagger \not{D} \chi_i - h(\chi_1 \chi_2 \Phi + \text{h.c.})$$

Anomalons: SM-singlet Dirac fermion


dark matter $m_\Psi = h\langle\Phi\rangle$

LHC production:

Gauged Symmetry: $\mathcal{X} \rightarrow B$: $q\bar{q} \rightarrow Z' \rightarrow \text{jets}$

Gauged Symmetry: $\mathcal{X} \rightarrow L$:

$$F_{\mu\nu} V^{\mu\nu}$$



$$\bar{\Psi}\Psi = \chi_1 \chi_2 + \chi_1^\dagger \chi_2^\dagger \rightarrow \chi_\alpha \chi_\beta \Phi^{(*)}, \quad \alpha = 1, \dots, N \rightarrow N > 4$$



$$F_{\mu\nu} \quad V^{\mu\nu}$$

Local $U(1)_{\mathcal{X}}$

$$\mathcal{L} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + i \sum_i \chi_i^\dagger \not{D} \chi_i - h(\chi_1 \chi_2 \Phi + \text{h.c.})$$


Anomalons: SM-singlet Dirac fermion

dark matter $m_\Psi = h\langle\Phi\rangle$

LHC production:

Gauged Symmetry: $\mathcal{X} \rightarrow B$: $q\bar{q} \rightarrow Z' \rightarrow \text{jets}$

Gauged Symmetry: $\mathcal{X} \rightarrow L$:



multi-component
dark matter

$$\bar{\Psi}\Psi = \chi_1\chi_2 + \chi_1^\dagger\chi_2^\dagger \rightarrow \chi_\alpha\chi_\beta\Phi^{(*)},$$

$$\alpha = 1, \dots, N \rightarrow N > 4$$



Local $U(1)_\chi$

$$\mathcal{L} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + i \sum_i \chi_i^\dagger \not{D} \chi_i - h(\chi_1 \chi_2 \Phi + \text{h.c.})$$

Anomalons: SM-singlet Dirac fermion


dark matter $m_\Psi = h\langle\Phi\rangle$

LHC production:

$$F_{\mu\nu} V^{\mu\nu}$$

Gauged Symmetry: $\mathcal{X} \rightarrow B$: $q\bar{q} \rightarrow Z' \rightarrow \text{jets}$

Gauged Symmetry: $\mathcal{X} \rightarrow L$:



multi-component
dark matter

$$\bar{\Psi}\Psi = \chi_1\chi_2 + \chi_1^\dagger\chi_2^\dagger \rightarrow \chi_\alpha\chi_\beta\Phi^{(*)},$$

$$\alpha = 1, \dots, N \rightarrow N > 4$$



Local $U(1)_{\mathcal{X}}$

$$\mathcal{L} = -\frac{1}{4} V_{\mu\nu} V^{\mu\nu} + i \sum_i \chi_i^\dagger \not{D} \chi_i - y (\chi_1 \chi_2 S + \text{h.c.})$$

Anomalons: SM-singlet Dirac fermion


CP violation Yukawa y

LHC production:

$$F_{\mu\nu} V^{\mu\nu}$$

Gauged Symmetry: $\mathcal{X} \rightarrow B$: $q\bar{q} \rightarrow Z' \rightarrow \text{jets}$

Gauged Symmetry: $\mathcal{X} \rightarrow L$:



multi-component
dark matter

$$\bar{\Psi}\Psi = \chi_1 \chi_2 + \chi_1^\dagger \chi_2^\dagger \rightarrow \chi_\alpha \chi_\beta \Phi^{(*)},$$

$$\alpha = 1, \dots, N \rightarrow N > 4$$

Any local Abelian extension of the Standard Model can be reduced to a set of integers which must satisfy the gravitational anomaly, $[SO(1,3)]^2 U(1)_Y$, and the cubic anomaly, $[U(1)_X]^3$ conditions:

$$\sum_{\alpha=1}^N z_{\alpha} = 0, \quad \sum_{\alpha=1}^N z_{\alpha}^3 = 0, \quad (1)$$

- From a list of $N - 2$ integers, e.g., for N even

$$\mathbf{q} = [l_1, l_2, \dots, l_n, k_1, k_2, \dots, k_n], \quad n = (N - 2)/2. \quad (2)$$

in the range $[-m, m]$, build two vector-like solutions of N integers,

$$\mathbf{x} = [l_1, k_1, \dots, k_n, -l_1, -k_1, \dots, -k_n], \quad \mathbf{y} = [0, 0, l_1, \dots, l_n, -l_1, \dots, -l_n] \quad (3)$$

- From a list of $N - 2$ integers, e.g., for N even

$$\mathbf{q} = [l_1, l_2, \dots, l_n, k_1, k_2, \dots, k_n], \quad n = (N - 2)/2. \quad (2)$$

in the range $[-m, m]$, build two vector-like solutions of N integers,

$$\mathbf{x} = [l_1, k_1, \dots, k_n, -l_1, -k_1, \dots, -k_n], \quad \mathbf{y} = [0, 0, l_1, \dots, l_n, -l_1, \dots, -l_n] \quad (3)$$

- Obtain a (some times) **non vector-like** solution with $z_{\max} = 2m$

$$\mathbf{z} = \mathbf{x} \oplus \mathbf{y} = \left(\sum_{i=1}^N x_i y_i^2 \right) \mathbf{x} + \left(\sum_{i=1}^N x_i^2 y_i \right) \mathbf{y}, \quad (4)$$

- From a list of $N - 2$ integers, e.g., for N even

$$\mathbf{q} = [l_1, l_2, \dots, l_n, k_1, k_2, \dots, k_n], \quad n = (N - 2)/2. \quad (2)$$

in the range $[-m, m]$, build two vector-like solutions of N integers,

$$\mathbf{x} = [l_1, k_1, \dots, k_n, -l_1, -k_1, \dots, -k_n], \quad \mathbf{y} = [0, 0, l_1, \dots, l_n, -l_1, \dots, -l_n] \quad (3)$$

- Obtain a (some times) **non vector-like** solution with $z_{\max} = 2m$

$$\mathbf{z} = \mathbf{x} \oplus \mathbf{y} = \left(\sum_{i=1}^N x_i y_i^2 \right) \mathbf{x} + \left(\sum_{i=1}^N x_i^2 y_i \right) \mathbf{y}, \quad (4)$$

The parameter space to be explored with $z_{\max} = 20$ ($m = 10$) has **96 153 non vector-like** solutions

$$\# \text{ of } \mathbf{q} \text{ lists} = (2m + 1)^{N-2} = \begin{cases} 9261 \rightarrow 3 & N = 5 \\ 194841 \rightarrow 38 & N = 6 \\ \vdots & \vdots \\ 1.6 \times 10^{13} \rightarrow 65910 & N = 12, \quad \text{instead } 10^{19} \end{cases} \quad (5)$$

- From a list of $N - 2$ integers, e.g., for N even

$$\mathbf{q} = [l_1, l_2, \dots, l_n, k_1, k_2, \dots, k_n], \quad n = (N - 2)/2. \quad (2)$$

in the range $[-m, m]$, build two vector-like solutions of N integers,

$$\mathbf{x} = [l_1, k_1, \dots, k_n, -l_1, -k_1, \dots, -k_n], \quad \mathbf{y} = [0, 0, l_1, \dots, l_n, -l_1, \dots, -l_n] \quad (3)$$

- Obtain a (some times) **non vector-like** solution with $z_{\max} = 2m$

$$\mathbf{z} = \mathbf{x} \oplus \mathbf{y} = \left(\sum_{i=1}^N x_i y_i^2 \right) \mathbf{x} + \left(\sum_{i=1}^N x_i^2 y_i \right) \mathbf{y}, \quad (4)$$

The parameter space to be explored with $z_{\max} = 20$ ($m = 10$) has **96 153 non vector-like** solutions

$$\# \text{ of } \mathbf{q} \text{ lists} = (2m + 1)^{N-2} = \begin{cases} 9261 \rightarrow 3 & N = 5 \rightarrow [1, 5, -7, -8, 9] \\ 194841 \rightarrow 38 & N = 6 \\ \vdots & \vdots \\ 1.6 \times 10^{13} \rightarrow 65910 & N = 12, \quad \text{instead } 10^{19} \end{cases} \quad (5)$$

Simplest secluded model with SM-singlet massive chiral fermions from SSB

$$\mathcal{L} = i\psi_i^\dagger \not{D}\psi_i - \frac{1}{4}V_{\mu\nu}V'^{\mu\nu} + \sum_{i<j} h_{ij}\psi_i\psi_j\phi^{(*)} + \text{h.c} \quad (6)$$

96 153 \rightarrow 5 196 multi-component DM ($N = 8, 12$) \rightarrow 28 with two Dirac-fermion DM

Simplest secluded model with SM-singlet massive chiral fermions from SSB

$$\mathcal{L} = i\psi_i^\dagger \not{D} \psi_i - \frac{1}{4} V_{\mu\nu} V'^{\mu\nu} + \sum_{i < j} h_{ij} \psi_i \psi_j \phi^{(*)} + \text{h.c} \quad (6)$$

96 153 \rightarrow 5 196 multi-component DM ($N = 8, 12$) \rightarrow 28 with two Dirac-fermion DM

$$\mathbf{z} = [1, 2, 2, 4, -5, -5, -7, 8] \rightarrow \phi = 3 \rightarrow [(1, 2), (2, -5), (-5, 8), (4, -7)] \quad (7)$$

Simplest secluded model with SM-singlet massive chiral fermions from SSB

$$\mathcal{L} = i\psi_i^\dagger \not{D} \psi_i - \frac{1}{4} V_{\mu\nu} V'^{\mu\nu} + \sum_{i < j} h_{ij} \psi_i \psi_j \phi^{(*)} + \text{h.c} \quad (6)$$

96 153 \rightarrow 5 196 multi-component DM ($N = 8, 12$) \rightarrow 28 with two Dirac-fermion DM

$$\mathbf{z} = [1, 2, 2, 4, -5, -5, -7, 8] \rightarrow \phi = 3 \rightarrow [(1, 2), (2, -5), (-5, 8), (4, -7)] \quad (7)$$

$$\mathcal{L} \subset \Psi^T \begin{array}{c} \begin{array}{cccccc} & 1 & 2 & 2 & -5 & -5 & 8 \\ \begin{array}{l} 1 \\ 2 \\ 2 \\ -5 \\ -5 \\ 8 \end{array} & \left[\begin{array}{cccccc} 0 & h_{(1,2)} & h_{(1,2)} & 0 & 0 & 0 \\ h_{(1,2)} & 0 & 0 & h_{(2,-5)} & h_{(2,-5)} & 0 \\ h_{(1,2)} & 0 & 0 & 0 & 0 & 0 \\ 0 & h_{(2,-5)} & 0 & 0 & 0 & h_{(-5,8)} \\ 0 & h_{(2,-5)} & 0 & 0 & 0 & h_{(-5,8)} \\ 0 & 0 & 0 & h_{(-5,8)} & h_{(-5,8)} & 0 \end{array} \right] \end{array} \Psi \phi^{(*)} + h_{(4,-7)} \psi_4 \psi_{-7} \phi^* \end{array} \quad (8)$$

Standard model extended with $U(1)_{\mathcal{X}=L \text{ or } B}$ gauge symmetry

Fields	$SU(2)_L$	$U(1)_Y$	$U(1)_{\mathcal{X}=B \text{ or } L}$
Q_i^\dagger	2	$-1/6$	Q
d_{Ri}	1	$-1/2$	d
u_{Ri}	1	$+2/3$	u
L_i^\dagger	2	$+1/2$	L
e_{Ri}	1	-1	e
H	2	$1/2$	$h = 0$
χ_α	1	0	z_α
$(L'_L)^\dagger$	2	$1/2$	$-x'$
L''_R	2	$-1/2$	x''
e'_R	1	-1	x'
$(e''_L)^\dagger$	1	1	$-x''$
Φ	1	0	ϕ
S	1	0	s

Table 1: A minimal set of new fermion content: $L = e = 0$ for $\mathcal{X} = B$. Or $Q = u = d = 0$ for $\mathcal{X} = L$.
 $i = 1, 2, 3, \alpha = 1, 2, \dots, N'$

$$\chi_1 \rightarrow \nu_{R1}, \dots, \chi_{N_\nu} \rightarrow \nu_{R N_\nu}, \quad 2 \leq N_\nu \leq 3, \quad (9)$$

$$\mathcal{L}_{\text{eff}} = h_\nu^{\alpha i} (\nu_{R\alpha})^\dagger \epsilon_{ab} L_i^a H^b \left(\frac{\Phi^*}{\Lambda} \right)^\delta + \text{H.c.}, \quad \text{with } i = 1, 2, 3,$$

S is the complex singlet scalar responsible for the SSB of the anomaly-free gauge symmetry with D or X -charge

$$\phi = -(\nu + L)/\delta, \quad (10)$$

Anomaly cancellation I

The anomaly-cancellation conditions on $[SU(3)_c]^2 U(1)_X$, $[SU(2)_L]^2 U(1)_X$, $[U(1)_Y]^2 U(1)_X$, allow us to express three of the X -charges in terms of the others

$$u = -e - \frac{2}{3}L - \frac{1}{9}(x' - x'') , \quad d = e + \frac{4}{3}L - \frac{1}{9}(x' - x'') , \quad Q = -\frac{1}{3}L + \frac{1}{9}(x' - x'') , \quad (11)$$

while the $[U(1)_X]^2 U(1)_Y$ anomaly condition reduces to

$$(e + L)(x' - x'') = 0 . \quad (12)$$

- Previously: $x' = x''$
- We choose instead ($h = 0$):

$$e = -L , \quad (13)$$

so that (L is still a free parameter)

$$Q = -u = -d = -\frac{1}{3}L + \frac{1}{9}(x' - x'') . \quad (14)$$

If $B = 0 \rightarrow U(1)_L$

Anomaly cancellation II

The gravitational anomaly, $[\mathrm{SO}(1,3)]^2 \mathrm{U}(1)_Y$, and the cubic anomaly, $[\mathrm{U}(1)_X]^3$, can be written as the following system of Diophantine equations, respectively,

$$\sum_{\alpha=1}^N z_{\alpha} = 0, \quad \sum_{\alpha=1}^N z_{\alpha}^3 = 0, \quad (15)$$

where $N = N' + 5$ and

$$\begin{aligned} z_{N'+1} &= -x', & z_{N'+2} &= x'', \\ z_{N'+2+i} &= L, \quad i = 1, 2, 3 \end{aligned} \quad (16)$$

→

$$9Q = - \sum_{\alpha=N'+1}^{N'+5} z_{\alpha} = -x' + x'' + L + L + L, \quad (17)$$

$$Q = 0 \rightarrow \mathrm{U}(1)_L$$

September 24, 2021

Dataset

Open Access

Set of N integers between -30 and 30 with sum and cubic sum up to zero for $4 < N < 13$

Diego Restrepo

Anomalies

Solutions obtained with the python package: [anomalies](#) based on the method to find anomaly free solutions of the standard model extended with an Abelian Dark Symmetry with N right-handed singlet chiral fields described in [arXiv:1905.13729](#) [PRL].

Data scheme

- 'l': integer lists → input to obtain the 'solution' by using the [anomalies](#) package
- 'k': integer lists → input to obtain the 'solution' by using the [anomalies](#) package

- 'solution': list → of integers, Z_i which satisfy $\sum_{i=1}^N Z_i = 0$ and $\sum_{i=1}^N Z_i^3 = 0$.

- 'n': integer → number of integers in 'solution', N .

USAGE

#Example of JSON file usage in Python with pandas (see also json module)

```
>>> import pandas as pd
>>> df=pd.read_json('solutions.json')
>>> df[:2]
```

	1	k	solution	gcd	n
0	[1, 2]	[0, -3]	[1, 5, -7, -8, 9]	1	5
1	[-2, -1]	[0, -1]	[2, 4, -7, -9, 10]	1	5

Data:

390074 solutions with $5 \leq N \leq 12$ integers until '[32]' [JSON]

17

views

4

downloads

[See more details...](#)

Indexed in

OpenAIRE

Publication date:

September 24, 2021

DOI:

DOI: [10.5281/zenodo.5526707](https://doi.org/10.5281/zenodo.5526707)

Keyword(s):

[Anomaly free](#) [Diophantine equations](#) [Abelian symmetry](#)
[Gauge Symmetry](#)

License (for files):

[Creative Commons Attribution 4.0 International](#)

Versions

Version 1

Sep 24, 2021

[10.5281/zenodo.5526707](https://doi.org/10.5281/zenodo.5526707)

- $L = 0$

$$(5, 5, -3, -2, 1, -6)$$

$U(1)_B$ selection

- $L = 0$
- Effective neutrino mass: $\phi = -\nu = -5$

$$(5, 5, -3, -2, 1, -6)$$

- $L = 0$
- Effective neutrino mass: $\phi = -\nu = -5$
- Electroweak-scale vector-like fermions:

$$(L'_L)^\dagger L''_R \Phi^* \rightarrow x' = -1, x'' = 6$$

$$(5, 5, -3, -2, 1, -6)$$

- $L = 0$
- Effective neutrino mass: $\phi = -\nu = -5$
- Electroweak-scale vector-like fermions:
 $(L'_L)^\dagger L''_R \Phi^* \rightarrow x' = -1, x'' = 6$
- Dirac-fermionic DM: $(\chi_L)^\dagger \chi''_R \Phi^* \rightarrow z_3 = -3, z_4 = -2$



$$(5, 5, -3, -2, 1, -6)$$

$U(1)_B$ selection

- $L = 0$
- Effective neutrino mass: $\phi = -\nu = -5$
- Electroweak-scale vector-like fermions:
 $(L'_L)^\dagger L''_R \Phi^* \rightarrow x' = -1, x'' = 6$
- Dirac-fermionic DM: $(\chi_L)^\dagger \chi''_R \Phi^* \rightarrow z_3 = -3, z_4 = -2$


959 solutions from $\sim 400,000$

$(5, 5, -3, -2, 1, -6)$




Scotogenic realization

Any realization which does not affect anomaly cancellation is allowed




Scotogenic realization

Any realization which does not affect anomaly cancellation is allowed



Scotogenic realization

Any realization which does not affect anomaly cancellation is allowed




Field	$SU(2)_L$	$U(1)_Y$	$U(1)_B$
u_{Ri}	1	2/3	$u = 1/3$
d_{Ri}	1	-1/3	$d = 1/3$
$(Q_i)^\dagger$	2	-1/6	$Q = -1/3$
$(L_i)^\dagger$	2	1/2	$L = 0$
e_R	1	-1	$e = 0$
$(L'_L)^\dagger$	2	1/2	$-x' = -3/5$
e'_R	1	-1	$x' = 3/5$
L''_R	2	-1/2	$x'' = 18/5$
$(e'_L)^\dagger$	1	1	$-x'' = -18/5$
$\nu_{R,1}$	1	0	-3
$\nu_{R,2}$	1	0	-3
χ_R	1	0	6/5
$(\chi_L)^\dagger$	1	0	9/5
H	2	1/2	0
S	1	0	3
Φ	1	0	3
σ^-_α	1	1	3/5
σ'^-_α	1	-1	-12/5

Electroweak baryogenesis

- Standard model (SM) $m_h \sim 125$ GeV. 😞
- Beyond the SM: Source of CP contains fields charged under SM
→ too large electric dipole moments 😞

- Inert SM-singlet complex scalar field which acquires vev with temperature to have strong electroweak phase transition 😊
- CP violation (CPV) triggered in dark sectors through SM gauge singlets
→ CPV Yukawa between SM-singlet complex scalar and SM-singlet quiral fermions 😊




$U(1)_L$ or $U(1)_B$
 Z'
 transfer CP to
 SM

1811.09719 [PRL]
 Carena...


Anomaly
 cancellation

Dirac
 neutrinos

Anomalons:
 DM



Dark sector baryogenesis



CP violation occurs in the dark sector and is transmitted to SM sector by the new Z' gauge boson.

- High scale fields: Φ , $(\langle\Phi\rangle \rightarrow L'_L, L''_R, e'_L, e''_R)$: EW-scale vector-like anomalous
- Electroweak scale (EW) fields: $Z'_\mu, S, \chi_L, \chi_R$
- CP-violation

$$\begin{aligned}\mathcal{L}_{\text{Dirac DM}} &= h(\chi_L)^\dagger \chi_R \Phi^* + y(\chi_L)^\dagger \chi_R S^* + \text{h.c.}, & y \in \mathbb{C} \\ &\supset \left(m_\chi + |y| e^{i\theta} |S|\right) (\chi_L)^\dagger \chi_R + \text{h.c.}\end{aligned}$$

- CP-violation Portal

$$\mathcal{L}_{\text{anomalous}} \supset g' Z'_\mu \left[3\bar{\chi}_L \gamma^\mu \chi_L - 2\bar{\chi}_R \gamma^\mu \chi_R + \bar{Q}_i \gamma^\mu Q_i + \bar{q}_{Ri} \gamma^\mu q_{Ri} \right]$$

- Strong electroweak phase transition (EWPT) portal

$$\mathcal{L}_{\text{first order EWPT}} \supset -\lambda_{SH} H^\dagger H S^* S.$$


First-order phase transition: Effective potential ($T \neq 0$)

$h = H/\sqrt{2}$, $s = |S|$ with vevs: $v(T)$ and $w(T)$ such that $v(T_c) = w(T_c)$

$$V_T(h, s) = \frac{\lambda_H v_c^4}{4} \left(\frac{h^2}{v_c^2} + \frac{s^2}{w_c^2} - 1 \right)^2 + \frac{\lambda_H v_c^2}{m_{s,c}^2 w_{0,c}^4} h^2 s^2 + (T^2 - T_c^2)(c_h h^2 + c_s s^2), \quad (18)$$

where

$$c_h = \frac{1}{48} (9g_2^2 + 3g_1^2 + 12y_t^2 + 24\lambda_H + \lambda_{HS}), \quad c_s = \frac{1}{12} (3\lambda_S + 2\lambda_{HS}). \quad (19)$$



First-order phase transition: Effective potential ($T \neq 0$)



$h = H/\sqrt{2}$, $s = |S|$ with vevs: $v(T)$ and $w(T)$ such that $v(T_c) = w(T_c)$

$$V_T(h, s) = \frac{\lambda_H v_c^4}{4} \left(\frac{h^2}{v_c^2} + \frac{s^2}{w_c^2} - 1 \right)^2 + \frac{\lambda_H v_c^2}{m_{s,c}^2 w_{0,c}^4} h^2 s^2 + (T^2 - T_c^2)(c_h h^2 + c_s s^2), \quad (18)$$

where

$$c_h = \frac{1}{48} (9g_2^2 + 3g_1^2 + 12y_t^2 + 24\lambda_H + \lambda_{HS}), \quad c_s = \frac{1}{12} (3\lambda_S + 2\lambda_{HS}). \quad (19)$$

arXiv: Sec. 4.1 arXiv:1107.5451



First-order phase transition: Effective potential ($T \neq 0$)



$h = H/\sqrt{2}$, $s = |S|$ with vevs: $v(T)$ and $w(T)$ such that $v(T_c) = w(T_c)$

$$V_T(h, s) = \frac{\lambda_H v_c^4}{4} \left(\frac{h^2}{v_c^2} + \frac{s^2}{w_c^2} - 1 \right)^2 + \frac{\lambda_H v_c^2}{m_{s,c}^2 w_{0,c}^4} h^2 s^2 + (T^2 - T_c^2)(c_h h^2 + c_s s^2), \quad (18)$$

where

$$c_h = \frac{1}{48} (9g_2^2 + 3g_1^2 + 12y_t^2 + 24\lambda_H + \lambda_{HS}), \quad c_s = \frac{1}{12} (3\lambda_S + 2\lambda_{HS}). \quad (19)$$

arXiv: Sec. 4.1 arXiv:1107.5451



First-order phase transition: Effective potential ($T \neq 0$)



$h = H/\sqrt{2}$, $s = |S|$ with vevs: $v(T)$ and $w(T)$ such that $v(T_c) = w(T_c)$

$$V_T(h, s) = \frac{\lambda_H v_c^4}{4} \left(\frac{h^2}{v_c^2} + \frac{s^2}{w_c^2} - 1 \right)^2 + \frac{\lambda_H v_c^2}{m_{s,c}^2 w_{0,c}^4} h^2 s^2 + (T^2 - T_c^2)(c_h h^2 + c_s s^2), \quad (18)$$

where

$$c_h = \frac{1}{48} (9g_2^2 + 3g_1^2 + 12y_t^2 + 24\lambda_H + \lambda_{HS}), \quad c_s = \frac{1}{12} (3\lambda_S + 2\lambda_{HS}). \quad (19)$$

arXiv: Sec. 4.1 arXiv:1107.5451



Using the thin wall approximation for the nucleation bubbles, we use the ansatz in which the space dependence of the fields is given by

$$h(z) = \frac{1}{2}v(T_n)(1 - \tanh(z/L_w)) , \quad s(z) = \frac{1}{2}w_0(T_n)(1 + \tanh(z/L_w)) ,$$

where z is the direction normal to the wall and L_w is the wall width.

The nucleation temperature, T_n , is defined by the condition

$$\exp(-S_3/T_n) = \frac{3}{4\pi} \left(\frac{H(T_n)}{T_n} \right)^4 \left(\frac{2\pi T_n}{S_3} \right)^{\frac{3}{2}} ,$$

where S_3 is the Euclidean action of the bubble and $H(T)$ is the Hubble rate.

Boltzmann equation i

$$\xi_i(z) \equiv \mu_i(z)/T = 6(n_i - \bar{n}_i)/T^3,$$

$$-D_L \xi''_{\chi_L} - v_w \xi'_{\chi_L} + \Gamma_L (\xi_{\chi_L} - \xi_{\chi_R}) = S_{\mathcal{CP}},$$

where D_L is the diffusion constant for χ_L , which is related to the scattering rate Γ_L by

$$D_L = \frac{3x+2}{x^2+3x+2} \frac{1}{3\Gamma_L}, \quad x \equiv m_\chi/T \quad (20)$$

and

$$S_{\mathcal{CP}} = -\frac{\lambda}{2} \frac{v_w D_L}{\frac{3x+2}{x^2+3x+2} T} \frac{(1-x)e^{-x} + x^2 E_1(x)}{4m_\chi^2 K_2(x)} \frac{m_\chi w_0(T_n) \lambda \left(-2 + \cosh\left(\frac{2z}{L_w}\right) \right) \sin \theta}{L_w^3 \cosh^4\left(\frac{z}{L_w}\right)}, \quad (21)$$

where v_w is the wall's velocity $E_1(x)$ is the error function and $K_2(x)$ is the modified Bessel function of the second kind. $y = \lambda e^{i\theta - i\pi/2}$

Transfer DM asymmetry to SM quarks

The chiral particle give rise to a non-zero $U(1)_B$ charge density in the proximity of the wall. This results in a Z' background that couples to the SM fields with $U(1)_B$ charge,

$$\langle Z'_0(z) \rangle = \frac{g_B (q_{\chi_L} - q_{\chi_R}) T_n^3}{6M_{Z'}} \int_{-\infty}^{\infty} dz_1 \xi_{\chi_L}(z_1) e^{-M_{Z'}|z-z_1|},$$

which generates a chemical potential for the SM quarks,

$$\mu_Q(z) = \mu_{d_R, u_R}(z) = 3 \times \frac{5}{9} \times g_B \langle Z'_0(z) \rangle.$$

This chemical potential sources a thermal-equilibrium asymmetry in the quarks, $\Delta n_Q^{\text{EQ}}(z) \sim T_n^2 \mu_Q(z)$.

From [1]

If the Z' is sufficiently light, it mediates a long range force that extends into the region outside the bubble wall with unbroken electroweak symmetry.

Finally, the baryon-number asymmetry is then given by

$$n_B = \frac{\Gamma_{\text{sph}}}{v_w} \int_0^\infty dz n_Q^{\text{EQ}}(z) \exp\left(-\frac{\Gamma_{\text{sph}}}{v_w} z\right),$$

where Γ_{sph} is the sphaleron rate. The baryon-to-photon-number ratio is then obtained by

$$\eta_B = \frac{n_B}{s(T)}, \quad s(T) \equiv \frac{2\pi^2}{45} g_{*S}(T) T^3,$$

where $g_{*S}(T)$ is the effective number of relativistic degrees of freedom.

Our goal is to find what regions of the parameter space yield

$$0.82 \times 10^{-10} < \eta_B < 0.92 \times 10^{-10}. \quad (22)$$

- SARAH→SPheno→MicroMegas
- η_B calculation code
- Python notebook with the scan

arXiv:1810.08055

Ten Simple Rules for Reproducible Research in Jupyter Notebook Fernando Pérez, *et al*


[...] In this paper, we address several questions about reproducibility [...] Combined with software repositories and open source licensing, notebooks are powerful tools for transparent, collaborative, reproducible, and reusable data analyses.

We vary the typical Dirac-fermion DM parameter space and for each point that satisfy neutrino oscillation data, relic density and DM direct detection constraints. For each point we ...

Parameter	Range
θ	$(-\pi/2, \pi/2)$
$w_0(T_n)/\text{GeV}$	100 – 500
T_n/GeV	100 – 200
L_w/GeV^{-1}	$1/T_n - 10/T_n$
v_w	0.05 – 0.5

Table 2: Scan ranges for the free parameters that are involved in the baryogenesis mechanism.

Black points: Dirac neutrinos with proper DM and baryon assymetry



A $U(1)_B$ is presented as an example of models where all new fermions required to cancel out the anomalies are used to solve phenomenological problems of the standard model (SM):

- EW-scale fermion vector-like doublets and iso-singlet charged singlets, in conjunction with right-handed neutrinos with repeated Abelian charges, participate in the generation of small neutrino masses through the Dirac-dark Zee mechanism
- The other SM-singlets are used to explain the dark matter in the universe, while their coupling to an inert singlet scalar is the source of the CP violation.

In the presence of a strong first-order electroweak phase transition, this “dark” CP violation allows for successful electroweak baryogenesis by using long range force mediated by a sufficiently light Z' which transfers the asymmetry from the Dark sector into the SM.