

Dark matter stability and Dirac neutrinos using only Standard Model symmetries

#1

Cesar Bonilla (Munich, Tech. U.), Salvador Centelles-Chuliá (Valencia U., IFIC), Ricardo Cepedello (Valencia U., IFIC), Eduardo Peinado (Mexico U.), Rahul Srivastava (Valencia U., IFIC) (Dec 4, 2018)

Published in: *Phys.Rev.D* 101 (2020) 3, 033011 • e-Print: [1812.01599](#) [hep-ph]

Minimal radiative Dirac neutrino mass models

#1

Julian Calle (Antioquia U.), Diego Restrepo (Antioquia U.), Carlos E. Yaguna (UPTC, Tunja), Óscar Zapata (Antioquia U.) (Dec 13, 2018)

Published in: *Phys.Rev.D* 99 (2019) 7, 075008 • e-Print: [1812.05523](#) [hep-ph]

 pdf  DOI  cite  claim  reference search  46 citations

$$\mathcal{L}_5 = \frac{h}{\Lambda} (\nu_R)^\dagger L \cdot HS^* + \text{H.c.},$$




FIG. 6. T3-1-A-I ($\alpha = 0$): $B - L$ flux in the Dirac radiative seesaw.

	$(\nu_{Ri})^\dagger$	$(\nu_{Rj})^\dagger$	$(\nu_{Rk})^\dagger$						S
T3-1-A-I				$(\psi_R)^\dagger$	ψ_L	η_a	σ_a	—	
L	+4	+4	−5	r	− r	$1 - r$	$4 - r$	—	3